

Genomic Structure, Chromosomal Mapping, and Expression Pattern of Human *DCAMKL1* (*KIAA0369*), a Homologue of *DCX* (*XLIS*)

Naomichi Matsumoto, Daniela T. Pilz, and David H. Ledbetter¹

Department of Human Genetics, The University of Chicago, 924 E. 57th Street, Chicago, Illinois 60637

Received July 27, 1998; accepted October 30, 1998

Human *DCAMKL1*, also known as *KIAA0369*, is a homologue of *DCX* (Xq22.3), a gene associated with X-linked lissencephaly and subcortical band heterotopia. This suggests that *DCAMKL1* may play a role in neuronal migration. The gene also shows similarity to Ca^{2+} /calmodulin-dependent protein kinases. We have determined its genomic structure, regional mapping, and expression pattern in human tissues. *DCAMKL1* consists of at least 18 exons ranging from 58 to 3359 bp in length. We have characterized the exon/intron borders, and primers were designed to amplify each individual exon for mutation analysis. *DCAMKL1* was mapped to chromosome 13q13 by fluorescence *in situ* hybridization. Northern blot analysis showed *DCAMKL1* to be predominantly expressed in human fetal brain as a major transcript of about 5.8 kb. © 1999 Academic Press

INTRODUCTION

Classical lissencephaly (LIS) is a neuronal migration disorder characterized by gyral abnormalities ranging from agyria (absent gyri) to pachygyria (abnormal broad gyri). Its spectrum also includes subcortical band heterotopia (bands of gray matter located in the white matter) (Dobyns and Truwit, 1995). Genetic causes of LIS include deletions or intragenic mutations of *LIS1* located on chromosome 17p13.3 (Reiner *et al.*, 1993; Chong *et al.*, 1997; Lo Nigro *et al.*, 1997) and mutations of *DCX* (*Doublecortin*, *X-linked*, aka *XLIS*) on Xq22.3 (des Portes *et al.*, 1998; Gleeson *et al.*, 1998). We have found that deletions and intragenic mutations of *LIS1* or *DCX* account for 76% of sporadic LIS (Pilz *et al.*, 1998). However, in 24% of these patients, the cause of their lissencephaly remains unexplained, and additional genes involved in neuronal migration are implicated.

DCAMKL1 (doublecortin and CaM kinase-like 1) has been noted to have similarity to *DCX* (des Portes *et al.*, 1998; Gleeson *et al.*, 1998). *DCX* has a 1080-bp open

reading frame encoding a 10-kb brain-specific message. At the nucleotide level, *DCAMKL1* shows 73% identity to *DCX* over 825 bp; however the open reading frame of *DCAMKL1* extends an additional 1104 bp in the 3' direction. The *DCX* protein, which appears to have potential MAP kinase family phosphorylation sites and to function as part of an Abl-dependent signal transduction pathway, is 75% identical at the amino acid level to the predicted protein of *DCAMKL1* (GenBank Accession No. AB002367). However the C-terminus of *DCAMKL1* is considered to be similar to the family of calcium-calmodulin-dependent protein kinases (CaM kinases) and shows high similarity to the rat CaM kinase CPG16 (GenBank U78857). It is thought to be involved in cell signaling/communication (Gleeson *et al.*, 1998).

Ohara *et al.* (1997) found that more than 80% of large proteins (>1000 amino acid residues) in public databases have biologically important functions. They isolated the human *DCAMKL1* gene, originally known as *KIAA0369*, in a project sequencing entire cDNA clones derived from relatively long transcripts. The *DCAMKL1* cDNA was derived from a set of size-fractionated cDNA libraries from human brain, was selected by analysis of the protein-coding potential in an *in vitro* system, and was entirely sequenced. It was mapped to chromosome 13 using a radiation hybrid panel and found to be expressed in human brain by RT-PCR analysis (Nagase *et al.*, 1997).

Here we report the genomic organization of *DCAMKL1*. The exon/intron borders of *DCAMKL1* were identified, and sizes of the exons and introns were determined. Primers were developed for PCR amplification of the individual exons, which will enable analysis of this gene as a candidate gene for human neuronal migration disorders. We regionally mapped *DCAMKL1* to 13q13 using fluorescence *in situ* hybridization (FISH). Its expression pattern in human tissues demonstrated that *DCAMKL1* is predominantly expressed in human fetal brain. While this paper was under review, Omori *et al.* (1998) reported similar data on the expression pattern and chromosomal localization of this gene.

¹ To whom correspondence and reprint requests should be addressed. Telephone: (773) 834-0525. Fax: (773) 834-0505. E-mail: dhl@genetics.uchicago.edu.




FIG. 1. Genomic structure of the human *DCAMKL1* gene with localization of the exons and size of the introns. The position of the overlapping genomic clones is indicated at the bottom; there is a gap between GS8J4 and GS157C18.

MATERIALS AND METHODS

Isolation of *DCAMKL1* genomic clones. A human BAC library (Genome Systems, St. Louis, MO) was screened by PCR according to the manufacturer's protocol using the following primers designed from the *DCAMKL1* cDNA sequence (Accession No. AB002367): SCR-A1(5'-AGACATGGAGCTGGAGCACT-3', nucleotides 15-34)/SCR-A2(5'-CTTCTTGGCCTTCTTCTCG-3', 156-175), SCR-B1(5'-TGATGTGCCTTCAGGACTTTT-3', 725-745)/SCR-B2(5'-TCCTGGTAACGGAACTTCTCC-3', 780-800), SCR-C1(5'-TCAATTTAATCATCACCACCG-3', 5050-5070)/SCR-C2(5'-AAAAGAAAGACGGTCCCAT-3', 5130-5149). PCR was performed for 35 cycles at 94°C for 30 s, 55°C for 30 s, and 72°C for 30 s.

Pulsed-field gel electrophoresis (PFGE) analysis. The Chef Mapper Pulsed-Field Electrophoresis System (Bio-Rad, Richmond, CA) was used for PFGE analysis. BAC DNA was extracted using an automatic nucleic acid isolation system (AutoGen 740, Integrated Separation Systems, Natick, MA). One microgram of DNA was digested with *NotI* for 3 h and electrophoresed in a 1.0% SeaKem GTG agarose gel (FMC Bioproducts, Rockland, ME) with an autoalgorithm mode ranging from 5 to 300 kb at 14°C. The gel was stained with ethidium bromide.

DNA sequencing. A primer-walking strategy was adopted with primers designed using *DCAMKL1* cDNA sequence and isolated BACs as templates. Purified BAC DNA was isolated using Qiagen Midi-Prep columns (Chatsworth, CA) according to the manufacturer's protocol, with the following modifications: Qiagen-Tip 100 columns were used for a 250-ml overnight BAC culture with 10 ml of P1, 10 ml of P2, and 10 ml of P3 solutions as described elsewhere (Hubert *et al.*, 1997). One microgram of BAC DNA and 40 pmol of primers were used for sequencing with the ABI Prism Dye or dRhodamine Terminator Cycle Sequencing Ready Reaction Kits (PE Applied Biosystems, Inc., Foster, CA) followed by analysis on an ABI 377 automated DNA sequencer. The sequencing reaction was cycled 50 times at 94°C for 30 s, 55°C for 30 s, and 72°C for 60 s. PCR products were also used for direct sequencing after purification using Microcon 100 (Amicon, Inc., Beverly, MA). Thirty nanograms of DNA from PCR products and 4 pmol of primers were used for the sequencing reaction with the above mentioned kits. The sequencing reaction was performed for 25 cycles at 96°C for 10 s, 50°C for 5 s, and 60°C for 4 min.

Analysis of intron lengths (intron analysis). Standard PCR or XL-PCR (Perkin-Elmer/Roche, Branchburg, NJ) was performed in a volume of 15 μ l containing 100 ng of BAC DNA as a template, 1 μ M of each primer, 200 μ M of each dNTP, 0.033 U/ μ l *Taq* Gold (standard PCR) or 0.025 U/ μ l *rTth* (XL-PCR) polymerase, and 1 \times standard PCR buffer (1.5 mM MgCl₂) or 1 \times XL-PCR buffer [1.25 mM Mg(OAc)₂]. Initial denaturation at 95°C for 10 min was followed by 35 cycles of denaturation at 95°C for 45 s, annealing at 55°C for 30 s, and extension at 72°C for 2.5 min (standard PCR) or at 72°C for 10 min (XL-PCR). Furthermore, for XL-PCR, 35 cycles of denaturation

at 95°C for 15 s and annealing and extension at 60-62°C for 10 min were used. Products were analyzed in a 0.8 or 0.4% agarose gel.

Fluorescence in situ hybridization. Chromosome preparations were made from normal lymphoblastoid cell lines by conventional methods. Slides were incubated in 2 \times SSC at 37°C for 30 min, serially dehydrated in 70, 80, and 95% ethanol at room temperature, denatured in 70% formamide/2 \times SSC at 72°C for 2 min, and then serially dehydrated at -20°C in 70, 80, 90, and 100% ethanol. BAC DNAs were labeled with digoxigenin-11-dUTP (Boehringer Mannheim, Indianapolis, IN) by nick-translation, precipitated in ethanol with a 50 \times excess of human Cot-1 and herring testis DNA (Gibco BRL, Gaithersburg, MD), and resuspended to a final concentration of 20 ng/ μ l in hybridization solution (50% formamide, 2 \times SSC, 10% dextran sulfate). Probes were denatured at 76°C for 10 min and preassociated at 37°C for 15 min prior to hybridization. The probe-hybridization mix (10 μ l) was applied under a 22 mm \times 22 mm coverslip, and slides were incubated in a moist chamber for 16 h at 37°C and then washed as previously described (Chong *et al.*, 1997). Probes were detected in 50 μ l of rhodamine anti-digoxigenin (Boehringer Mannheim) at 1 μ g/ml. Slides were washed three times in 4 \times SSC, 0.1% Tween 20 at 45°C and mounted in antifade solution

TABLE 1
Exon/Intron Organization of the Human *DCAMKL1* Gene

Exon	Size (bp)	5' splice donor	Intron size (kb)	3' splice acceptor
1	(ND)	ND	(ND)	cgtagGTCCTA
2	(395)	TGGAAGgtgac	(18)	cttagGAGAGT
3	(347)	AAACAGgtaag	(ND)	tgtagGTGATG
4	(100)	AAAGTGgtaag	(3.8)	cccagAATGTC
5	(117)	GCTCAGgtaaa	(15)	tctagTTAATG
6	(95)	CAGAGGgtaag	(20)	tacagATCTCT
7	(85)	GAGAAGgtgat	(2.9)	ttcagAAGTGT
8	(109)	AGAAAGgtgag	(1.8)	cgcagATCGAC
9	(58)	GGCAAgtatg	(0.8)	gacagGAGCAC
10	(120)	GTAAGgtgtg	(4.0)	ctcagGGGGGA
11	(147)	CTGCTGgtaag	(15)	tttagGTGTAT
12	(172)	GACTGGgtaaa	(2.0)	tccagATACGG
13	(78)	CCGTGGgtatg	(0.8)	aacagAAGTGG
14	(97)	GCAAAGgtagg	(2.4)	cacagGAGCTC
15	(81)	GTTAATgtaag	(0.7)	tccagGATGAT
16	(114)	ATAGCAgtgtaag	(1.6)	ctcagCTGGAC
17	(74)	GATAAGgtatg	(18)	gctagACCACC
18	(>3359)			

Note. Intron sequences are in lowercase letters, and exons are in uppercase letters. ND, not determined.

TABLE 2
Intron Primers for Amplification of *DCAMKL1* Exons

Exon	Primer sequences	Product size (bp)	Annealing temperature (°C)
2	5'-AAC CTA CAT CAA TAT GTG TGT CTG C-3'	484	55
	5'-TGG AAC CTGTAG GGT CAC G-3'		
3	5'-AGT AGG AAA AAA GAA ATG GGC A-3'	449	55
	5'-TTG CCT TAA AGT CAA AGT CAC A-3'		
4	5'-CGT GGG GTT GCA TTG TTG T-3'	219	50
	5'-TTT CGA AAG CTG CCT CAA AT-3'		
5	5'-CCC CAT AAT GAT GTG TGG TTT-3'	194	55
	5'-GGA ACA AGG CAA TTT CTT CAA-3'		
6	5'-CAT GAA AAT TAT TTA ACG TGC CA-3'	194	55
	5'-GGC TGA GAC AGG TCT GAA ACA-3'		
7	5'-AAA CTC AGG CAT TCT GAC CC-3'	180	55
	5'-CAT CCT CTG CCT CCT CAG AT-3'		
8	5'-TGT GTT CCC TTT TCA ACC AA-3'	213	55
	5'-GGA AAA CAT AAC GCC AAA TG-3'		
9	5'-TGA TCC CTG ACA AAA TAA TTC AA-3'	168	55
	5'-AAC ACT CTT ATC TCA TAA GAA C-3'		
10	5'-TAT GGA AGC TGA TGA AGA TAA C-3'	200	55
	5'-CGG TGC CAT CAA TAA AGA CT-3'		
11	5'-TGT GCT GCT TCA GCT TCT GT-3'	215	55
	5'-GGA ACT CAG GAT GCC CTG TA-3'		
12	5'-TGG TTT TCA GAC AAT TGT GAT TTT-3'	211	55
	5'-TTT GCA AGC ATA GCA CTT AAA CA-3'		
13	5'-CGA GCC TGA CCC TCC TAA CT-3'	166	55
	5'-TAG AGA AAT GCT TTG TCG GC-3'		
14	5'-AGT GCT GTT TTT CCT TGT TGT C-3'	181	55
	5'-CGT TAA GAC ACA CAG AAT ATA GGG A-3'		
15	5'-CAG CTT TTC ACC TTA AAT TCA GA-3'	180	55
	5'-AAA AGG AAT GTT AGG AGA GAA CAA A-3'		
16	5'-CAT TTT TCA TTT ATC TCA TTT TCT TTC-3'	205	50
	5'-AAA AGT TAT AGG TGG ATA TTT CAG ATG-3'		
17	5'-GCT TGG TCT GGA GTC TTA ATT CTT T-3'	209	50
	5'-GGT TAA TGG AGA CTG CCA GC-3'		
18	5'-CCT CGC CCT AAT TGT CCT CT-3'	216	50
	5'-GGG TCT TAT TAA AAG GGC GA-3'		

(Vector) containing DAPI. Analysis of more than 50 cells was performed using a Zeiss Axiophot microscope equipped with filters to detect DAPI and rhodamine separately, as well as a triple-bandpass filter (Chroma Technology Corp.) to detect signals simultaneously. Images were collected and merged using a cooled CCD camera (KAF 1400, Photometrics) and IP Lab Spectrum software (Signal Analytics Corp.) or Quips mFISH Imaging software (Vysis, Inc.).

Northern analysis. Expression of *DCAMKL1* mRNA was examined by Northern hybridization using a 330-bp PCR product of the coding region of *DCAMKL1* as a probe corresponding to nucleotides 15–344. Human multiple tissue blots (2 µg of poly(A)⁺ RNA loaded per lane) were obtained from commercial sources (Clontech Laboratories, Inc., Palo Alto, CA). The blots were hybridized with the ³²P-labeled probe in ExpressHyb Hybridization Solution (Clontech), washed in 0.1× SSC/0.1× SDS at 50°C, and analyzed on a Storm 860 imaging system and with ImageQuant software from the manufacturer (Molecular Dynamics, Inc., Sunnyvale, CA).

RESULTS AND DISCUSSION

Genomic Structure of *DCAMKL1*

We isolated the BAC clones GS8J4, GS157P10, and GS112N1 with primer SCR-A1/SCR-A2, GS157C18 with SCR-B1/SCR-B2, and GS53H5 and GS228G20 with SCR-C1/SCR-C2 (Fig. 1). PFGE analysis showed that the sizes of the BAC clone inserts ranged from 90 to

180 kb (data not shown). Of these clones, GS8J4 (130 kb), GS157C18 (130 kb), and GS53H5 (180 kb) were used for further analysis because of their large inserts. Sequence analysis with primers designed from the cDNA sequence (Nagase *et al.*, 1997) and BACs as templates revealed that the human *DCAMKL1* gene consists of at least 18 exons and 17 introns. The organization of the gene is depicted in Fig. 1. The size of each exon and intron and the sequence of the exon/intron junctions are shown in Table 1. The size of these exons range from 58 bp (exon 9) to more than 3359 bp (exon 18). The exon/intron junctions conform to the 5'-donor and 3'-acceptor consensus (GT...AG) (Table 1). Exon 1 and part of exon 2 encode the 5'-untranslated region of the *DCAMKL1* cDNA. The *DCAMKL1* coding region starts at base 21 of exon 2, spans exons 3 to 17, and extends to the first 58 bp of exon 18. Exon 18 is the largest among the exons, and most of it constitutes the 3'- untranslated region of the *DCAMKL1* cDNA. The length of 15 of the 16 introns from introns 2 to 17 in the *DCAMKL1* gene was determined by PCR; the smallest is intron 15 (0.7 kb). The size of intron 3 could not be determined due to a gap in the BAC contigs. The primers designed to amplify the individual

exons and the appropriate PCR conditions are listed in Table 2.

Regional Chromosomal Mapping and Expression Pattern of *DCAMKL1*

The regional chromosomal localization of *DCAMKL1* was determined by FISH using BACs GS8J4 (covering exons 2 and 3) and GS53H5 (covering the 3'-UTR). Two signals were observed on chromosome 13, band q13, with both BACs (Fig. 2), and no signal was seen on any other chromosomes.

Northern blot analysis showed that *DCAMKL1* is expressed as a major transcript of about 5.8 kb. Weak hybridization signals of about 6.0 and 9.0 kb were also observed (Fig. 3). These signals could be due to hybridization of the *DCAMKL1* probe to unprocessed *DCAMKL1* transcripts or minor splicing variants. Omori *et al.* (1998) showed four splicing variants of *DCAMKL1*. Two variants contained both the *DCX*-like domain and the CAM kinase-like domain of the gene, and the other two contained only the CAM kinase-like domain. The former was expressed mainly in fetal brain, and the latter showed similar expression in fetal and adult brain. We also found expression of *DCAMKL1* mRNA at high levels only in the fetal brain using a probe in the *DCX*-like domain. Low levels of expression were observed in adult brain, ovary, and prostate.

Two neurological disorders, nocturnal enuresis 1 (ENUR1) (OMIM600631) and Moebius syndrome (OMIM157900), have been previously mapped to this region. ENUR1 has been linked to the markers D13S291 and D13S263 at 13q13–q14.3 (Eiberg *et al.*, 1995). Some Moebius syndrome patients have shown




FIG. 2. Regional mapping of the *DCAMKL1* gene by fluorescence *in situ* hybridization. The *DCAMKL1* gene is localized to human chromosome 13q13 using a digoxigenin-dUTP-labeled GS53H5 clone containing the *DCAMKL1* gene as a probe. Reverse DAPI banding pattern of both chromosomes 13 is shown in bottom right corner.




FIG. 3. Northern analysis of human *DCAMKL1* showing predominant expression in the fetal brain as a major transcript of 5.8 kb and minor transcripts of 6.0 and 9.0 kb. The image was obtained after overnight exposure (12 h) to a storage phosphor screen.

structural chromosomal abnormalities including 13q12.2–q13 (Slee *et al.*, 1991; Ziter *et al.*, 1977). Although abnormal neuronal migration phenotypes have so far not been mapped to 13q13, sequence similarity of *DCAMKL1* to *DCX*, their coexpression in developing brain, and simultaneous temporal expression pattern in the mouse embryo after day E-11 (Omori *et al.*, 1998) place *DCAMKL1* as a potential candidate gene for neuronal migration disorders.

ACKNOWLEDGMENTS

We are grateful to Dr. Naohiko Seki at the Kazusa DNA Institute, Japan, and Dr. Elizabeth M. Ross at the Laboratory of Molecular Neurobiology and Development, University of Minnesota for their helpful comments, and we thank Dr. Judy A. Fantes and Ms. Stephanie Mewborn for expert technical assistance with the FISH experiments. N.M. was supported in part by a research fellowship of the Uehara Memorial Foundation.

REFERENCES

- Chong, S. S., Pack, S. D., Roschke, A. V., Tanigami, A., Carrozzo, R., Smith, A. C. M., Dobyns, W. B., and Ledbetter, D. H. (1997). A revision of the lissencephaly and Miller-Dieker syndrome critical regions in chromosome 17p13.3. *Hum. Mol. Genet.* **6**: 147–155.
- des Portes, V., Pinard, J. M., Billuart, P., Vinet, M. C., Koulakoff, A., Carrie, A., Gelot, A., Dupuis, E., Motte, J., Berwald-Netter, Y., Catala, M., Kahn, A., Beldjord, C., and Chelly, J. (1998). A novel CNS gene required for neuronal migration and involved in X-linked subcortical laminar heterotopia and lissencephaly syndrome. *Cell* **92**: 51–61.
- Dobyns, W. B., and Truwit, C. L. (1995). Lissencephaly and other malformations of cortical development: 1995 update. *Neuropediatrics* **23**: 132–147.
- Eiberg, H., Berendt, I., and Mohr, J. (1995). Assignment of dominant inherited nocturnal enuresis (ENUR1) to chromosome 13q. *Nat. Genet.* **10**: 354–356.
- Gleeson, J. G., Allen, K. M., Fox, J. W., Lamperti, E. D., Berkovic, S., Scheffer, I., Cooper, E. C., Dobyns, W. B., Minnerath, S. R., Ross, M. E., and Walsh, C. A. (1998). Doublecortin, a brain-specific gene mutated in human X-linked lissencephaly and double cortex syndrome, encodes a putative signaling protein. *Cell* **92**: 63–72.
- Hubert, R. S., Mitchell, S., Chen, X.-N., Ekmekji, K., Gadowski, C., Sun, Z., Noya, D., Kim, U.-J., Chen, C., Shizuya, H., Simon, M., De

- Jong, P. J., and Korenberg, J. R. (1997). BAC and PAC contigs covering 3.5 Mb of the Down syndrome congenital heart disease region between D21S55 and MX1 on chromosome 21. *Genomics* **41**: 218–226.
- Lo Nigro, C., Chong, S. S., Smith, A. C. M., Dobyns, W. B., and Ledbetter, D. H. (1997). Point mutations and an intragenic deletion in *LIS1*, the lissencephaly causative gene in isolated lissencephaly sequence and Miller–Dieker syndrome. *Hum. Mol. Genet.* **6**: 157–164.
- Nagase, T., Ishikawa, K., Nakajima, D., Ohira, M., Seki, N., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N., and Ohara, O. (1997). Prediction of the coding sequences of unidentified human genes. VII. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNA Res.* **4**: 141–150.
- Ohara, O., Nagase, T., Ishikawa, K., Nakajima, D., Ohira, M., Seki, N., and Nomura, N. (1997). Construction and characterization of human brain cDNA libraries suitable for analysis of cDNA clones encoding relatively large proteins. *DNA Res.* **4**: 53–59.
- Omori, Y., Suzuki, M., Ozaki, K., Harada, Y., Nakamura, Y., Takahashi, E.-I., and Fujiwara, T. (1998). Expression and chromosomal localization of KIAA0369, a putative kinase structurally related to doublecortin. *J. Hum. Genet.* **43**: 169–177.
- Pilz, D. T., Matsumoto, N., Minnerath, S., Mills, P., Gleeson, J. G., Allen, K. M., Walsh, C. A., Barkovich, A. J., Dobyns, W. B., Ledbetter, D. H., Ross, M. E. (1998). *LIS1* and *XLIS (DCX)* mutations cause most classical lissencephaly, but different patterns of malformation. *Hum. Mol. Genet.* **7**: 2029–2037.
- Reiner, O., Carrozzo, R., Shen, Y., Wehnert, M., Faustinella, F., Dobyns, W. B., Caskey, C. T., and Ledbetter, D. H. (1993). Isolation of a Miller–Dieker lissencephaly gene containing G protein β -subunit-like repeats. *Nature* **364**: 717–721.
- Slee, J. J., Smart, R. D., and Viljoen, D. L. (1991). Deletion of chromosome 13 in Moebius syndrome. *J. Med. Genet.* **28**: 413–414.
- Ziter, F. A., Wiser, W. C., and Robinson, A. (1977). Three-generation pedigree of a Moebius syndrome variant with chromosome translocation. *Arch. Neurol.* **34**: 437–442.