

**Problem**

---

Let  $\mathbf{v}_1$ ,  $\mathbf{v}_2$ , and  $\mathbf{c}$  be  $n \times 1$  column vectors, and let  $\mathbf{M}_1$  and  $\mathbf{M}_2$  be  $n \times n$  matrices. Suppose

$$\begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{M}_1 \\ \mathbf{M}_2 \end{bmatrix} \mathbf{c}.$$

Show that if  $\det(\mathbf{M}_1) \neq 0$ , then

$$\mathbf{v}_2 = \mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{v}_1.$$

**Solution**

---

Left-multiply both sides of the given equation by the  $n \times 2n$  block matrix  $\begin{bmatrix} \mathbf{M}_2 \mathbf{M}_1^{-1} & -\mathbf{I}_{n \times n} \end{bmatrix}$ , where  $\mathbf{I}_{n \times n}$  is the  $n \times n$  identity matrix. On the left-hand side of the equation, we obtain

$$\begin{aligned} \begin{bmatrix} \mathbf{M}_2 \mathbf{M}_1^{-1} & -\mathbf{I}_{n \times n} \end{bmatrix} \begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \end{bmatrix} &= \mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{v}_1 - \mathbf{I}_{n \times n} \mathbf{v}_2 \\ &= \mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{v}_1 - \mathbf{v}_2. \end{aligned}$$

On the right-hand side of the equation, we obtain

$$\begin{aligned} \begin{bmatrix} \mathbf{M}_2 \mathbf{M}_1^{-1} & -\mathbf{I}_{n \times n} \end{bmatrix} \begin{bmatrix} \mathbf{M}_1 \\ \mathbf{M}_2 \end{bmatrix} \mathbf{c} &= \mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{M}_1 \mathbf{c} - \mathbf{I}_{n \times n} \mathbf{M}_2 \mathbf{c} \\ &= \mathbf{M}_2 \mathbf{c} - \mathbf{M}_2 \mathbf{c} \\ &= \mathbf{0}. \end{aligned}$$

Therefore,

$$\mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{v}_1 - \mathbf{v}_2 = \mathbf{0};$$

i.e.,

$$\mathbf{v}_2 = \mathbf{M}_2 \mathbf{M}_1^{-1} \mathbf{v}_1,$$

as asserted. ■

**Remark**

---

The interesting thing about this problem is that the equation

$$\begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{M}_1 \\ \mathbf{M}_2 \end{bmatrix} \mathbf{c}$$

looks like the equation

$$\frac{a}{b} = \frac{c}{d}.$$

The solution for  $b$  in the latter equation is

$$b = \frac{d}{c}a = dc^{-1}a,$$

while the solution for  $\mathbf{v}_2$  in the former equation is

$$\mathbf{v}_2 = \mathbf{M}_2\mathbf{M}_1^{-1}\mathbf{v}_1.$$

Strangely enough, you can solve for  $\mathbf{v}_2$  almost by pretending that you have fractions instead of block matrices.