

# DATA605 - Assignment 3

Nick Oliver

IS 605 FUNDAMENTALS OF COMPUTATIONAL MATHEMATICS - 2014

1. Problem set 1

(1) What is the rank of the matrix A?

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -1 & 0 & 1 & 3 \\ 0 & 1 & -2 & 1 \\ 5 & 4 & -2 & -3 \end{bmatrix}$$

Convert A to RREF

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -1 & 0 & 1 & 3 \\ 0 & 1 & -2 & 1 \\ 5 & 4 & -2 & -3 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rank is 4. There are no linear independent rows

```
A = matrix(c(1,2,3,4,-1,0,1,3,0,1,-2,1,5,4,-2,-3),nrow=4,byrow = TRUE)
A
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    1    2    3    4
## [2,]   -1    0    1    3
## [3,]    0    1   -2    1
## [4,]    5    4   -2   -3
```

```
pracma::rref(A)
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    1    0    0    0
## [2,]    0    1    0    0
## [3,]    0    0    1    0
## [4,]    0    0    0    1
```

```
Matrix::rankMatrix(A)
```

```
## [1] 4
## attr(,"method")
## [1] "tolNorm2"
## attr(,"useGrad")
## [1] FALSE
## attr(,"tol")
## [1] 8.881784e-16
```

- (2) Given an  $m \times n$  matrix where  $m > n$ , what can be the maximum rank? The minimum rank, assuming that the matrix is non-zero?

The maximum rank would be  $n$  as it is possible as in problem set one that no rows are linearly independent. The minimum rank is 1

- (3) What is the rank of matrix B?

$$B = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \\ 2 & 4 & 2 \end{bmatrix}$$

Convert B to reduced row echelon form. Count number of non-zero rows

$$B = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \\ 2 & 4 & 2 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

```
B = matrix(c(1,2,1,3,6,3,2,4,2),nrow=3,byrow=TRUE)
B
```

```
##      [,1] [,2] [,3]
## [1,]    1    2    1
## [2,]    3    6    3
## [3,]    2    4    2
```

```
pracma::rref(B)
```

```
##      [,1] [,2] [,3]
## [1,]    1    2    1
## [2,]    0    0    0
## [3,]    0    0    0
```

```
Matrix::rankMatrix(B)
```

```
## [1] 1
## attr(,"method")
## [1] "tolNorm2"
## attr(,"useGrad")
## [1] FALSE
## attr(,"tol")
## [1] 6.661338e-16
```

2. Problem set 2 Compute the eigenvalues and eigenvectors of the matrix A. You'll need to show your work. You'll need to write out the characteristic polynomial and show your solution.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

$$\lambda I_3 = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$\lambda I_3 - A = \begin{bmatrix} \lambda - 1 & -2 & -3 \\ 0 & \lambda - 4 & -5 \\ 0 & 0 & \lambda - 6 \end{bmatrix}$$

Apply rule of Sarrus

$$\lambda I_3 - A = \begin{bmatrix} \lambda - 1 & -2 & -3 \\ 0 & \lambda - 4 & -5 \\ 0 & 0 & \lambda - 6 \end{bmatrix} \begin{bmatrix} \lambda - 1 & -2 \\ 0 & \lambda - 4 \\ 0 & 0 \end{bmatrix}$$

$$(\lambda - 1)(\lambda - 4)(\lambda - 6) + (-2)(-5)(0) + (-3)(0)(0) - (-2)(0)(\lambda - 6) - (\lambda - 1)(-5)(0) - (-3)(\lambda - 4)(0)$$

Everything drops out except the first terms

$$(\lambda - 1)(\lambda - 4)(\lambda - 6)$$

This leaves us with three eigenvalues

$$\lambda = 1$$

$$\lambda = 4$$

$$\lambda = 6$$

Eigenvector for

$$\lambda = 1$$

```
A = matrix(c(1,2,3,0,4,5,0,0,6),nrow=3,byrow=TRUE)
```

```
A
```

```
##      [,1] [,2] [,3]
## [1,]    1    2    3
## [2,]    0    4    5
## [3,]    0    0    6
```

```
eigen(A)
```

```
## eigen() decomposition
```

```
## $values
```

```
## [1] 6 4 1
```

```
##
```

```
## $vectors
```

```
##      [,1]      [,2] [,3]
```

```
## [1,] 0.5108407 0.5547002    1
```

```
## [2,] 0.7981886 0.8320503    0
```

```
## [3,] 0.3192754 0.0000000    0
```