## DATA605 - Assignment 3

## Nick Oliver

## IS 605 FUNDAMENTALS OF COMPUTATIONAL MATHEMATICS - 2014

- 1. Problem set 1
- (1) What is the rank of the matrix A?

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -1 & 0 & 1 & 3 \\ 0 & 1 & -2 & 1 \\ 5 & 4 & -2 & -3 \end{bmatrix}$$

Convert A to RREF

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -1 & 0 & 1 & 3 \\ 0 & 1 & -2 & 1 \\ 5 & 4 & -2 & -3 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rank is 4. There are no linear independent rows

```
A = matrix(c(1,2,3,4,-1,0,1,3,0,1,-2,1,5,4,-2,-3),nrow=4,byrow = TRUE)
A
```

```
## [,1] [,2] [,3] [,4]
## [1,] 1 2 3 4
## [2,] -1 0 1 3
## [3,] 0 1 -2 1
## [4,] 5 4 -2 -3
```

pracma::rref(A)

Matrix::rankMatrix(A)

```
## [1] 4
## attr(,"method")
## [1] "tolNorm2"
## attr(,"useGrad")
## [1] FALSE
## attr(,"tol")
## [1] 8.881784e-16
```

(2) Given an mxn matrix where m > n, what can be the maximum rank? The minimum rank, assuming that the matrix is non-zero?

The maximum rank would be m as it is possible as in problem set one that no rows are linearly independent. The minimum rank is 1

(3) What is the rank of matrix B?

$$B = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \\ 2 & 4 & 2 \end{bmatrix}$$

Convert B to reduced row echelon form. Count number of non-zero rows

$$B = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \\ 2 & 4 & 2 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

```
B = matrix(c(1,2,1,3,6,3,2,4,2),nrow=3,byrow=TRUE)
B
```

```
## [,1] [,2] [,3]
## [1,] 1 2 1
## [2,] 3 6 3
## [3,] 2 4 2
```

pracma::rref(B)

Matrix::rankMatrix(B)

```
## [1] 1
## attr(,"method")
## [1] "tolNorm2"
## attr(,"useGrad")
## [1] FALSE
## attr(,"tol")
## [1] 6.661338e-16
```

2. Problem set 2 Compute the eigenvalues and eigenvectors of the matrix A. You'll need to show your work. You'll need to write out the characteristic polynomial and show your solution.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

$$\lambda I_3 = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$\lambda I_3 - A = \begin{bmatrix} \lambda - 1 & -2 & -3 \\ 0 & \lambda - 4 & -5 \\ 0 & 0 & \lambda - 6 \end{bmatrix}$$

Apply rule of Sarrus