

Term 3 Practice Activity: C4 Variance and Standard Deviation

Contents

1	Uniform Distribution	2
2	Triangular Distribution 1	4
3	Triangular Distribution 2	7
4	Normal Distribution	10
5	Comparing Uniform Distributions	13
6	Comparing Triangular Distributions	14
7	Comparing Normal Distributions	15
8	Uniform with Parameters	15
9	Triangular with Parameters	16
10	Normal with Parameters	17
11	Uniform with Parameters	18
12	Triangular with Parameters	18
13	Normal with Parameters	19
14	Uniform with Parameters	20
15	Triangular with Parameters	21
16	Normal with Parameters	22

Evaluated criteria

C4 Interprets the concept of variance and standard deviation of a probability function.

1 Uniform Distribution

Problem. A student's daily streaming time X (minutes) is modeled by

$$X \sim \text{Uniform}(2, 14).$$

Questions/Tasks.

- a) Find $E[X]$.
- b) Find $\text{Var}(X)$ and the standard deviation.
- c) Write the PDF of X .
- d) Draw the graph of the PDF, indicate $E[X]$, and draw an interval of one standard deviation around $E[X]$.
- e) Calculate the probability of the highlighted interval.
- f) Compare the probabilities of intervals of the same length around 4 and around 12. What is the difference between these probabilities?

C4

For $X \sim \text{Uniform}(a, b)$:

$$E[X] = \frac{a+b}{2}, \quad \text{Var}(X) = \frac{(b-a)^2}{12}.$$

- a) With $a = 2$ and $b = 14$,

$$E[X] = \frac{2+14}{2} = 8.$$

- b) Using the same endpoints,

$$\text{Var}(X) = \frac{(14-2)^2}{12} = \frac{144}{12} = 12,$$

so

$$\sigma = \sqrt{12} = 2\sqrt{3} \approx 3.46.$$

- c) Since $X \sim \text{Uniform}(2, 14)$, the PDF is constant on the support $[2, 14]$. Its height is


$$\frac{1}{14-2} = \frac{1}{12}.$$

Therefore,

$$f(x) = \begin{cases} \frac{1}{12}, & 2 \leq x \leq 14, \\ 0, & \text{otherwise.} \end{cases}$$

- d) From parts a and b, the mean is $E[X] = 8$ and the standard deviation is $\sigma = 2\sqrt{3}$. Therefore, one standard deviation around the mean is

$$[E[X] - \sigma, E[X] + \sigma] = [8 - 2\sqrt{3}, 8 + 2\sqrt{3}] \approx [4.54, 11.46].$$



e) The highlighted interval is the interval found in part d:

$$\left[8 - 2\sqrt{3}, 8 + 2\sqrt{3}\right].$$

So the probability of the highlighted interval is

$$P\left(8 - 2\sqrt{3} \leq X \leq 8 + 2\sqrt{3}\right).$$

Using the PDF from part c, the height of the rectangle is $\frac{1}{12}$. The base of the rectangle is the length of the highlighted interval:

$$\left(8 + 2\sqrt{3}\right) - \left(8 - 2\sqrt{3}\right) = 4\sqrt{3}.$$

Therefore,

$$P\left(8 - 2\sqrt{3} \leq X \leq 8 + 2\sqrt{3}\right) = (\text{base})(\text{height}) = (4\sqrt{3})\left(\frac{1}{12}\right).$$

Thus,


$$P\left(8 - 2\sqrt{3} \leq X \leq 8 + 2\sqrt{3}\right) = \frac{4\sqrt{3}}{12} = \frac{\sqrt{3}}{3} \approx 0.577.$$

f) Since this is a continuous model, the probability of any exact single value is 0. So we compare intervals of the same length around 4 and around 12.

Using the PDF from part c, any interval inside $[2, 14]$ with length L has probability

$$P(\text{interval}) = L\left(\frac{1}{12}\right) = \frac{L}{12}.$$

The following graph shows the 8 intervals. Intervals of length 1 are the darkest, and intervals of length 4 are the lightest.



Interval length	Around 4	Probability	Around 12	Probability	Difference
1	[3.5, 4.5]	$\frac{1}{12}$	[11.5, 12.5]	$\frac{1}{12}$	$\frac{0}{12}$
2	[3, 5]	$\frac{2}{12}$	[11, 13]	$\frac{2}{12}$	$\frac{0}{12}$
3	[2.5, 5.5]	$\frac{3}{12}$	[10.5, 13.5]	$\frac{3}{12}$	$\frac{0}{12}$
4	[2, 6]	$\frac{4}{12}$	[10, 14]	$\frac{4}{12}$	$\frac{0}{12}$

In each row, the two intervals have the same length and lie completely inside the support $[2, 14]$. Since the distribution is uniform, their probabilities are equal.

The difference between the probabilities around 4 and around 12 is always

0.

2 Triangular Distribution 1

Problem. A bus wait time W (minutes) is modeled by a triangular distribution with minimum $a = 2$, mode $c = 2$, and maximum $b = 14$:

$$W \sim \text{Triangular}(2, 2, 14).$$

Questions/Tasks.

- Find $E[W]$.
- Find $\text{Var}(W)$ and the standard deviation.
- Write the PDF of W .
- Draw the graph of the PDF, indicate $E[W]$, and draw an interval of one standard deviation around $E[W]$.
- Calculate the probability of the highlighted interval.
- Compare the probabilities of intervals of the same length around 4 and around 12. What is the difference between these probabilities?

C4

For $\text{Triangular}(a, c, b)$:

$$E[W] = \frac{a + b + c}{3}, \quad \text{Var}(W) = \frac{a^2 + b^2 + c^2 - ab - ac - bc}{18}.$$

- a) With $a = 2$, $c = 2$, and $b = 14$,

$$E[W] = \frac{2 + 14 + 2}{3} = \frac{18}{3} = 6.$$

- b) Using the same parameters,

$$\text{Var}(W) = \frac{2^2 + 14^2 + 2^2 - (2)(14) - (2)(2) - (14)(2)}{18} = \frac{4 + 196 + 4 - 28 - 4 - 28}{18} = \frac{144}{18} = 8.$$

Therefore,

$$\sigma = \sqrt{8} = 2\sqrt{2} \approx 2.83.$$

- c) Since the mode is at the left endpoint, the PDF is a decreasing straight line from $w = 2$ to $w = 14$. The total area under the PDF must be 1. Therefore, the maximum height is found from the area of the full triangle:

$$A_{\Delta} = \frac{1}{2}(\text{base})(\text{height}) = 1 \quad , \quad \text{base} = 14 - 2 = 12.$$

So

$$A_{\Delta} = \frac{1}{2}(12)(\text{height}) = 1 \quad \longrightarrow \quad 6(\text{height}) = 1 \quad \longrightarrow \quad \text{height} = \frac{1}{6}.$$

The PDF has height $\frac{1}{6}$ at $w = 2$ and height 0 at $w = 14$. Since this is one decreasing line segment, use the basic line form

$$f(w) = m(w - c),$$

To find c , use the fact that the line has height 0 at $w = 14$:

$$f(14) = 0 \quad \longrightarrow \quad m(14 - c) = 0 \quad \longrightarrow \quad 14 - c = 0 \quad \longrightarrow \quad c = 14.$$

Therefore, the line has the form

$$f(w) = m(w - 14), \quad 2 \leq w \leq 14.$$

To find m , use the fact that the line has height $\frac{1}{6}$ at the mode $w = 2$:

$$f(2) = \frac{1}{6} \quad \longrightarrow \quad m(2 - 14) = \frac{1}{6} \quad \longrightarrow \quad -12m = \frac{1}{6} \quad \longrightarrow \quad m = -\frac{1}{72}.$$

Therefore,


$$f(w) = -\frac{1}{72}(w - 14) = \frac{14 - w}{72}, \quad 2 \leq w \leq 14.$$

Thus,

$$f(w) = \begin{cases} \frac{14-w}{72}, & 2 \leq w \leq 14, \\ 0, & \text{otherwise.} \end{cases}$$

- d) From parts a and b, the mean is $E[W] = 6$ and the standard deviation is $\sigma = 2\sqrt{2}$. Therefore, one standard deviation around the mean is

$$[E[W] - \sigma, E[W] + \sigma] = [6 - 2\sqrt{2}, 6 + 2\sqrt{2}] \approx [3.17, 8.83].$$



- e) The highlighted interval is the interval found in part d:

$$[6 - 2\sqrt{2}, 6 + 2\sqrt{2}].$$

The probability of this interval is represented by the area under the PDF above the highlighted interval:

$$P(6 - 2\sqrt{2} \leq W \leq 6 + 2\sqrt{2}).$$

Since the PDF is a straight line over this interval, the highlighted region is a trapezoid. Therefore,

$$P(6 - 2\sqrt{2} \leq W \leq 6 + 2\sqrt{2}) = \frac{1}{2}(\text{width})(\text{left height} + \text{right height}).$$

Using the PDF from part c, the height at the left endpoint of the highlighted interval is

$$\text{left height} = f(6 - 2\sqrt{2}) = \frac{14 - (6 - 2\sqrt{2})}{72} = \frac{8 + 2\sqrt{2}}{72} = \frac{4 + \sqrt{2}}{36}.$$

The height at the right endpoint is

$$\text{right height} = f(6 + 2\sqrt{2}) = \frac{14 - (6 + 2\sqrt{2})}{72} = \frac{8 - 2\sqrt{2}}{72} = \frac{4 - \sqrt{2}}{36}.$$

The width of the trapezoid is

$$\text{width} = (6 + 2\sqrt{2}) - (6 - 2\sqrt{2}) = 4\sqrt{2}.$$

Substituting the width and the two heights into the trapezoid area formula,

$$P(6 - 2\sqrt{2} \leq W \leq 6 + 2\sqrt{2}) = \frac{1}{2}(4\sqrt{2}) \left(\frac{4 + \sqrt{2}}{36} + \frac{4 - \sqrt{2}}{36} \right).$$

Since

$$\frac{4 + \sqrt{2}}{36} + \frac{4 - \sqrt{2}}{36} = \frac{8}{36} = \frac{2}{9},$$

we get


$$P(6 - 2\sqrt{2} \leq W \leq 6 + 2\sqrt{2}) = \frac{1}{2}(4\sqrt{2}) \left(\frac{2}{9} \right) = \frac{4\sqrt{2}}{9} \approx 0.629.$$

- f) Since this is a continuous model, the probability of any exact single value is 0. So we compare intervals of the same length around 4 and around 12.

Using the PDF from part c, each interval probability is found as the area of a trapezoid:

$$P(r \leq W \leq s) = \frac{1}{2}(\text{interval length})(f(r) + f(s)).$$

The following graph shows the 8 intervals. Intervals of length 1 are the darkest, and intervals of length 4 are the lightest.



Interval length	Around 4	Probability	Around 12	Probability	Difference
1	[3.5, 4.5]	$\frac{5}{36}$	[11.5, 12.5]	$\frac{1}{36}$	$\frac{4}{36}$
2	[3, 5]	$\frac{10}{36}$	[11, 13]	$\frac{2}{36}$	$\frac{8}{36}$
3	[2.5, 5.5]	$\frac{15}{36}$	[10.5, 13.5]	$\frac{3}{36}$	$\frac{12}{36}$
4	[2, 6]	$\frac{20}{36}$	[10, 14]	$\frac{4}{36}$	$\frac{16}{36}$

The intervals around 4 have larger probability than the intervals around 12. This happens because the density is higher near the left side of the distribution and decreases as W increases.

3 Triangular Distribution 2

Problem. A bus wait time W (minutes) is modeled by a triangular distribution with minimum $a = 2$, mode $c = 6$, and maximum $b = 14$:

$$W \sim \text{Triangular}(2, 6, 14).$$

Questions/Tasks.

- Find $E[W]$.
- Find $\text{Var}(W)$ and the standard deviation.
- Write the PDF of W .
- Draw the graph of the PDF, indicate $E[W]$, and draw an interval of one standard deviation around $E[W]$.
- Calculate the probability of the highlighted interval.
- Compare the probabilities of intervals of the same length around 4 and around 12. What is the difference between these probabilities?

C4

For $\text{Triangular}(a, c, b)$:

$$E[W] = \frac{a + b + c}{3}, \quad \text{Var}(W) = \frac{a^2 + b^2 + c^2 - ab - ac - bc}{18}.$$

- a) With $a = 2$, $c = 6$, and $b = 14$,

$$E[W] = \frac{2 + 14 + 6}{3} = \frac{22}{3} \approx 7.33.$$

- b) Using the same parameters,

$$\text{Var}(W) = \frac{2^2 + 14^2 + 6^2 - (2)(14) - (2)(6) - (14)(6)}{18} = \frac{4 + 196 + 36 - 28 - 12 - 84}{18} = \frac{112}{18} = \frac{56}{9}.$$

Therefore,

$$\sigma = \sqrt{\frac{56}{9}} = \frac{2\sqrt{14}}{3} \approx 2.49.$$

- c) Since the mode is at $w = 6$, the PDF increases linearly from $w = 2$ to $w = 6$, then decreases linearly from $w = 6$ to $w = 14$. The total area under the PDF must be 1. Therefore, the maximum height is found from the area of the full triangle:

$$A_{\Delta} = \frac{1}{2}(\text{base})(\text{height}) = 1 \quad , \quad \text{base} = 14 - 2 = 12.$$

So

$$A_{\Delta} = \frac{1}{2}(12)(\text{height}) = 1 \quad \longrightarrow \quad 6(\text{height}) = 1 \quad \longrightarrow \quad \text{height} = \frac{1}{6}.$$

The graph has two line segments. For each segment, use the basic line form

$$f(w) = m(w - c).$$

For the left side, the line has height 0 at $w = 2$. To find c , use

$$f(2) = 0 \quad \longrightarrow \quad m_1(2 - c) = 0 \quad \longrightarrow \quad 2 - c = 0 \quad \longrightarrow \quad c = 2.$$

Therefore, the left side has the form

$$f(w) = m_1(w - 2), \quad 2 \leq w \leq 6.$$

To find m_1 , use the fact that the line has height $\frac{1}{6}$ at the mode $w = 6$:

$$f(6) = \frac{1}{6} \quad \longrightarrow \quad m_1(6 - 2) = \frac{1}{6} \quad \longrightarrow \quad 4m_1 = \frac{1}{6} \quad \longrightarrow \quad m_1 = \frac{1}{24}.$$

Therefore,

$$f(w) = \frac{1}{24}(w - 2) = \frac{w - 2}{24}, \quad 2 \leq w \leq 6.$$

For the right side, the line has height 0 at $w = 14$. To find c , use

$$f(14) = 0 \quad \longrightarrow \quad m_2(14 - c) = 0 \quad \longrightarrow \quad 14 - c = 0 \quad \longrightarrow \quad c = 14.$$

Therefore, the right side has the form

$$f(w) = m_2(w - 14), \quad 6 \leq w \leq 14.$$

To find m_2 , use the fact that the line has height $\frac{1}{6}$ at the mode $w = 6$:

$$f(6) = \frac{1}{6} \quad \longrightarrow \quad m_2(6 - 14) = \frac{1}{6} \quad \longrightarrow \quad -8m_2 = \frac{1}{6} \quad \longrightarrow \quad m_2 = -\frac{1}{48}.$$

Therefore,


$$f(w) = -\frac{1}{48}(w - 14) = \frac{14 - w}{48}, \quad 6 \leq w \leq 14.$$

Thus,

$$f(w) = \begin{cases} \frac{w-2}{24}, & 2 \leq w \leq 6, \\ \frac{14-w}{48}, & 6 \leq w \leq 14, \\ 0, & \text{otherwise.} \end{cases}$$

- d) From parts a and b, the mean is $E[W] = \frac{22}{3}$ and the standard deviation is $\sigma = \frac{2\sqrt{14}}{3}$. Therefore, one standard deviation around the mean is

$$[E[W] - \sigma, E[W] + \sigma] = \left[\frac{22 - 2\sqrt{14}}{3}, \frac{22 + 2\sqrt{14}}{3} \right] \approx [4.84, 9.83].$$



e) The highlighted interval is the interval found in part d:

$$\left[\frac{22 - 2\sqrt{14}}{3}, \frac{22 + 2\sqrt{14}}{3} \right].$$

The probability of this interval is represented by the area under the PDF above the highlighted interval:

$$P\left(\frac{22 - 2\sqrt{14}}{3} \leq W \leq \frac{22 + 2\sqrt{14}}{3}\right).$$

Instead of calculating the highlighted area directly, use the complement rule. The total area under the PDF is 1, so the highlighted probability is

$$1 - (\text{left extreme triangle}) - (\text{right extreme triangle}).$$

Using the PDF from part c, the height at the left endpoint of the highlighted interval is

$$\text{left height} = f\left(\frac{22 - 2\sqrt{14}}{3}\right) = \frac{\frac{22 - 2\sqrt{14}}{3} - 2}{24} = \frac{8 - \sqrt{14}}{36}.$$

The base of the left extreme triangle is

$$\text{left base} = \frac{22 - 2\sqrt{14}}{3} - 2 = \frac{16 - 2\sqrt{14}}{3}.$$

Therefore, the area of the left extreme triangle is

$$A_{L\Delta} = \frac{1}{2} \left(\frac{16 - 2\sqrt{14}}{3} \right) \left(\frac{8 - \sqrt{14}}{36} \right) = \frac{13}{18} - \frac{4\sqrt{14}}{27}.$$

Using the PDF from part c, the height at the right endpoint of the highlighted interval is

$$\text{right height} = f\left(\frac{22 + 2\sqrt{14}}{3}\right) = \frac{14 - \frac{22 + 2\sqrt{14}}{3}}{48} = \frac{10 - \sqrt{14}}{72}.$$

The base of the right extreme triangle is

$$\text{right base} = 14 - \frac{22 + 2\sqrt{14}}{3} = \frac{20 - 2\sqrt{14}}{3}.$$

Therefore, the area of the right extreme triangle is

$$A_{R\Delta} = \frac{1}{2} \left(\frac{20 - 2\sqrt{14}}{3} \right) \left(\frac{10 - \sqrt{14}}{72} \right) = \frac{19}{36} - \frac{5\sqrt{14}}{54}.$$

By the complement rule,

$$P\left(\frac{22-2\sqrt{14}}{3} \leq W \leq \frac{22+2\sqrt{14}}{3}\right) = 1 - A_{L\Delta} - A_{R\Delta}$$

Substituting the two triangle areas,

$$P\left(\frac{22-2\sqrt{14}}{3} \leq W \leq \frac{22+2\sqrt{14}}{3}\right) = 1 - \left(\frac{13}{18} - \frac{4\sqrt{14}}{27}\right) - \left(\frac{19}{36} - \frac{5\sqrt{14}}{54}\right).$$

Thus,


$$P\left(\frac{22-2\sqrt{14}}{3} \leq W \leq \frac{22+2\sqrt{14}}{3}\right) = -\frac{1}{4} + \frac{13\sqrt{14}}{54} \approx 0.651.$$

- f) Since this is a continuous model, the probability of any exact single value is 0. So we compare intervals of the same length around 4 and around 12.

Using the PDF from part c, each interval probability is found as the area of a trapezoid:

$$P(r \leq W \leq s) = \frac{1}{2}(\text{interval length})(f(r) + f(s)).$$

The following graph shows the 8 intervals. Intervals of length 1 are the darkest, and intervals of length 4 are the lightest.



Interval length	Around 4	Probability	Around 12	Probability	Difference
1	[3.5, 4.5]	$\frac{2}{24}$	[11.5, 12.5]	$\frac{1}{24}$	$\frac{1}{24}$
2	[3, 5]	$\frac{4}{24}$	[11, 13]	$\frac{2}{24}$	$\frac{2}{24}$
3	[2.5, 5.5]	$\frac{6}{24}$	[10.5, 13.5]	$\frac{3}{24}$	$\frac{3}{24}$
4	[2, 6]	$\frac{8}{24}$	[10, 14]	$\frac{4}{24}$	$\frac{4}{24}$

The intervals around 4 have larger probability than the intervals around 12. This happens because 4 is closer to the mode than 12, so the density is higher around 4.

4 Normal Distribution

Problem. Quiz scores S are modeled by

$$S \sim N(6, 4^2).$$

Questions/Tasks.

- a) State $E[S]$.
- b) State $\text{Var}(S)$ and the standard deviation.
- c) Write the PDF of S .
- d) Give the intervals $[\mu - \sigma, \mu + \sigma]$ and $[\mu - 2\sigma, \mu + 2\sigma]$, draw the graph of the PDF, indicate $E[S]$, and draw the interval of one standard deviation around $E[S]$.
- e) Find the probability of the highlighted interval $[\mu - \sigma, \mu + \sigma]$.
- f) Compare the probabilities of intervals of the same length around 6 and around -2 . What is the difference between these probabilities?

C4

For $N(\mu, \sigma^2)$,

$$E[S] = \mu, \quad \text{Var}(S) = \sigma^2.$$

- a) Since

$$S \sim N(6, 4^2),$$

the mean is

$$E[S] = \mu = 6.$$

- b) Using the same parameters,

$$\text{Var}(S) = \sigma^2 = 4^2 = 16.$$

Therefore,

$$\sigma = 4.$$

- c) For a normal distribution, the PDF is

$$f(s) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(s-\mu)^2}{2\sigma^2}}.$$

Using $\mu = 6$ and $\sigma = 4$, the PDF is

$$f(s) = \frac{1}{4\sqrt{2\pi}} e^{-\frac{(s-6)^2}{2(4)^2}}.$$

Equivalently,


$$f(s) = \frac{1}{4\sqrt{2\pi}} e^{-\frac{(s-6)^2}{32}}, \quad -\infty < s < \infty.$$

- d) From parts a and b, the mean is $E[S] = 6$ and the standard deviation is $\sigma = 4$. Therefore,

$$[\mu - \sigma, \mu + \sigma] = [6 - 4, 6 + 4] = [2, 10].$$

Also,

$$[\mu - 2\sigma, \mu + 2\sigma] = [6 - 2(4), 6 + 2(4)] = [-2, 14].$$



e) The highlighted interval is the interval found in part d:

$$[2, 10].$$

The probability of this interval is represented by the area under the normal curve above the highlighted interval:

$$P(2 \leq S \leq 10).$$

To use the standard normal distribution, standardize with

$$Z = \frac{S - \mu}{\sigma} = \frac{S - 6}{4}.$$

Then

$$P(2 \leq S \leq 10) = P\left(\frac{2 - 6}{4} \leq Z \leq \frac{10 - 6}{4}\right).$$

So

$$P(2 \leq S \leq 10) = P(-1 \leq Z \leq 1).$$

Using the standard normal distribution,

$$P(-1 \leq Z \leq 1) \approx 0.6827.$$

Therefore,

$$P(2 \leq S \leq 10) \approx 0.6827.$$


f) Since this is a continuous model, the probability of any exact single value is 0. So we compare intervals of the same length around 4 and around 12.

Using the standardization formula from part e,

$$Z = \frac{S - 6}{4},$$

each interval probability can be found by converting the endpoints to z -scores and using the standard normal distribution.

The following graph shows the 8 intervals. Intervals of length 1 are the darkest, and intervals of length 4 are the lightest.



Interval length	Around 4	Probability	Around 12	Probability	Difference
1	[3.5, 4.5]	0.0878	[11.5, 12.5]	0.0325	0.0553
2	[3, 5]	0.1747	[11, 13]	0.0656	0.1091
4	[2, 6]	0.3413	[10, 14]	0.1359	0.2054
8	[0, 8]	0.6247	[8, 16]	0.3023	0.3223

The intervals around 4 have larger probability than the intervals around 12. This happens because 4 is closer to the mean 6, where the normal density is highest. The value 12 is farther from the mean, so intervals around 12 have less area under the curve.

5 Comparing Uniform Distributions

Problem. Compare these uniform models for download times (seconds):

$$X_1 \sim \text{Uniform}(4, 10), \quad X_2 \sim \text{Uniform}(6, 12), \quad X_3 \sim \text{Uniform}(4, 14).$$

Questions/Tasks.

- Find $E[X_i]$ and $\text{Var}(X_i)$ for $i = 1, 2, 3$.
- Compare X_1 and X_2 : only location changes. How do center and spread change?
- Compare X_1 and X_3 : only width changes from 6 to 10. How does variance change?
- Which model has the widest spread, and why?


C4

Using $E = \frac{a+b}{2}$, $\text{Var} = \frac{(b-a)^2}{12}$:

$$E[X_1] = 7, \quad \text{Var}(X_1) = \frac{6^2}{12} = 3;$$

$$E[X_2] = 9, \quad \text{Var}(X_2) = \frac{6^2}{12} = 3;$$

$$E[X_3] = 9, \quad \text{Var}(X_3) = \frac{10^2}{12} = \frac{25}{3}.$$



Shifting both endpoints by the same amount changes the mean but not spread. Increasing interval width increases variance.

6 Comparing Triangular Distributions

Problem. A packing process time T (minutes) follows triangular models:

$$T_1 \sim \text{Triangular}(2, 5, 10), \quad T_2 \sim \text{Triangular}(2, 7, 10), \quad T_3 \sim \text{Triangular}(4, 7, 12).$$


Questions/Tasks.

- Compute $E[T_i]$ and $\text{Var}(T_i)$ for each model.
- Compare T_1 and T_2 : only mode changes. How do center and shape change?
- Compare T_2 and T_3 : all three parameters shift by $+2$. What stays the same and what changes?
- Which model is most concentrated near larger times? Explain using the peak location.

C4

Using triangular formulas:

$$\begin{aligned} E[T_1] &= \frac{2 + 5 + 10}{3} = \frac{17}{3}, & \text{Var}(T_1) &= \frac{2^2 + 5^2 + 10^2 - 2 \cdot 5 - 2 \cdot 10 - 5 \cdot 10}{18} = \frac{49}{18}. \\ E[T_2] &= \frac{2 + 7 + 10}{3} = \frac{19}{3}, & \text{Var}(T_2) &= \frac{2^2 + 7^2 + 10^2 - 2 \cdot 7 - 2 \cdot 10 - 7 \cdot 10}{18} = \frac{49}{18}. \\ E[T_3] &= \frac{4 + 7 + 12}{3} = \frac{23}{3}, & \text{Var}(T_3) &= \frac{4^2 + 7^2 + 12^2 - 4 \cdot 7 - 4 \cdot 12 - 7 \cdot 12}{18} = \frac{49}{18}. \end{aligned}$$



Changing only the mode shifts peak and mean but here keeps variance unchanged; shifting all parameters by +2 shifts location but not spread.

7 Comparing Normal Distributions

Problem. Daily step counts (thousands) are modeled as:


$$N_1 \sim N(9, 1.5^2), \quad N_2 \sim N(11, 1.5^2), \quad N_3 \sim N(9, 2.5^2).$$

Questions/Tasks.

- State mean and variance for each distribution.
- Compare N_1 and N_2 : only mean changes. What changes on the graph?
- Compare N_1 and N_3 : only standard deviation changes. What happens to spread and peak height?
- Which distribution gives a higher chance of being far from its mean? Explain.

C4

$$E[N_1] = 9, \text{ Var}(N_1) = 2.25; \quad E[N_2] = 11, \text{ Var}(N_2) = 2.25; \quad E[N_3] = 9, \text{ Var}(N_3) = 6.25.$$



Changing μ moves the center left or right without changing shape width. Increasing σ makes the curve wider and less peaked, so far-from-mean values become more likely.

8 Uniform with Parameters

Problem. A random time X (minutes) is

$$X \sim \text{Uniform}(p, p + 12), \quad p \in \mathbb{R}.$$


Questions/Tasks.

- Express $E[X]$ in terms of p .

- b) Express $\text{Var}(X)$ in terms of p .
- c) Which parameter controls location and which controls spread?
- d) If p increases by 5, how do mean and variance change?

C4

$$E[X] = \frac{p + (p + 12)}{2} = p + 6, \quad \text{Var}(X) = \frac{((p + 12) - p)^2}{12} = \frac{12^2}{12} = 12.$$



So p controls location; spread is fixed by width 12 and does not depend on p .

9 Triangular with Parameters

Problem. A project completion time Y (days) is modeled by

$$Y \sim \text{Triangular}(4, 4 + m, 14), \quad 0 \leq m \leq 10.$$

Questions/Tasks.


- a) Write $E[Y]$ and $\text{Var}(Y)$ as functions of m .
- b) State the condition on m that makes the model valid.
- c) How does the mean change as m increases?
- d) How does moving the mode affect the shape (left/right skewness)?

C4

Here $a = 4$, $c = 4 + m$, $b = 14$.

$$E[Y] = \frac{a + b + c}{3} = \frac{4 + 14 + (4 + m)}{3} = \frac{22 + m}{3}.$$

$$\text{Var}(Y) = \frac{a^2 + b^2 + c^2 - ab - ac - bc}{18} = \frac{16 + 196 + (4 + m)^2 - 56 - 4(4 + m) - 14(4 + m)}{18} = \frac{m^2 - 10m + 100}{18}.$$



Validity needs $a \leq c \leq b \Rightarrow 4 \leq 4 + m \leq 14 \Rightarrow 0 \leq m \leq 10$. Mean increases linearly with m . As mode moves right, the distribution becomes less right-skewed and can become left-skewed when the mode passes midpoint 9.

10 Normal with Parameters

Problem. A test score variable is


$$Z \sim N(60 + k, (2k)^2), \quad k > 0.$$

Questions/Tasks.

- Express mean and variance in terms of k .
- Determine valid values of k .
- As k increases, how do center and spread change?
- Identify which part of the model controls location and which controls spread.

C4

$$E[Z] = 60 + k, \quad \text{Var}(Z) = (2k)^2 = 4k^2, \quad \sigma = 2k.$$



For a valid non-degenerate normal spread, $k > 0$. Increasing k shifts mean right and also increases spread quadratically through variance.

11 Uniform with Parameters

Problem. Commute time C (minutes) is modeled by


$$C \sim \text{Uniform}(10 - r, 10 + r), \quad r > 0.$$

Questions/Tasks.

- a) Find $E[C]$ and $\text{Var}(C)$ in terms of r .
- b) What condition on r is needed for a valid interval?
- c) As r increases, what happens to interval width and spread?
- d) For what r is the distribution narrower, and why?

C4

$$E[C] = \frac{(10 - r) + (10 + r)}{2} = 10,$$
$$\text{Var}(C) = \frac{((10 + r) - (10 - r))^2}{12} = \frac{(2r)^2}{12} = \frac{r^2}{3}.$$



Need $r > 0$. Width is $2r$, so larger r means wider interval and larger variance; smaller positive r means narrower distribution.

12 Triangular with Parameters

Problem. A completion score Q is modeled by

$$Q \sim \text{Triangular}(a, a + 3, a + 9), \quad a \in \mathbb{R}.$$


Questions/Tasks.

- a) Express $E[Q]$ and $\text{Var}(Q)$ in terms of a .
- b) Is this model always valid for any real a ? Explain.
- c) Which parameter controls location? Which controls spread?
- d) If a increases, what happens to center, spread, and shape?

C4

$$E[Q] = \frac{a + (a + 3) + (a + 9)}{3} = a + 4.$$

$$\text{Var}(Q) = \frac{a^2 + (a + 3)^2 + (a + 9)^2 - a(a + 3) - a(a + 9) - (a + 3)(a + 9)}{18} = \frac{7}{2}.$$



Since $a \leq a + 3 \leq a + 9$ always holds, the distribution is valid for all real a . Parameter a shifts location; spread and shape remain unchanged.

13 Normal with Parameters

Problem. Reaction time R (seconds) is modeled by


$$R \sim N(m, (1 + s)^2), \quad s > -1.$$

Questions/Tasks.

- Express $E[R]$ and $\text{Var}(R)$ in terms of m and s .
- Give the condition(s) needed for a valid nonzero spread.
- Explain how increasing m changes the distribution.
- Explain how increasing s changes the distribution width and likelihood of values far from the mean.

C4

$$E[R] = m, \quad \text{Var}(R) = (1 + s)^2, \quad \sigma = 1 + s.$$



A valid standard deviation needs $\sigma > 0 \Rightarrow 1 + s > 0 \Rightarrow s > -1$. Increasing m shifts the center only. Increasing s increases spread, making the bell wider and less concentrated around the mean.

14 Uniform with Parameters

Problem. A delivery time U (minutes) for a local route is modeled by

$$U \sim \text{Uniform}(h - 2w, h + w), \quad w > 0.$$

Questions/Tasks.

- Derive $E[U]$, $\text{Var}(U)$, and σ_U in terms of h and w .
- Identify which parameter controls the center and which controls the interval width.
- Compare Case A: $(h, w) = (30, 4)$ and Case B: $(h, w) = (35, 4)$. Does the distribution shift, widen, or both?
- Compare Case C: $(h, w) = (30, 4)$ and Case D: $(h, w) = (30, 7)$. How do center, width, and spread change?
- If both h and w increase by 2 at the same time, explain whether the model shifts, widens, or both.

C4


For $U \sim \text{Uniform}(a, b)$, $E[U] = \frac{a+b}{2}$, $\text{Var}(U) = \frac{(b-a)^2}{12}$. Here $a = h - 2w$, $b = h + w$, so

$$\begin{aligned} E[U] &= \frac{(h - 2w) + (h + w)}{2} = h - \frac{w}{2}, \\ \text{Var}(U) &= \frac{((h + w) - (h - 2w))^2}{12} = \frac{(3w)^2}{12} = \frac{3w^2}{4}, \\ \sigma_U &= \sqrt{\frac{3w^2}{4}} = \frac{\sqrt{3}}{2} w \quad (w > 0). \end{aligned}$$

The center $h - \frac{w}{2}$ depends on both h and w , while the width $b - a = 3w$ depends only on w . Case A $\rightarrow$ B keeps w fixed, so width and spread stay the same, and the model shifts right by 5. Case C $\rightarrow$ D keeps h fixed and increases w , so width grows from 12 to 21, spread increases, and the center moves from 28 to 26.5 (left). If h and w both increase by 2, then width increases by $3(2) = 6$ (widens) and

$$E[U]' - (E[U]) = \left(h + 2 - \frac{w + 2}{2}\right) - \left(h - \frac{w}{2}\right) = 1,$$

so it also shifts right by 1.



15 Triangular with Parameters

Problem. A machine-cycle time T (seconds) is modeled by

$$T \sim \text{Triangular}(k, k + r, k + 3r), \quad r > 0.$$

Questions/Tasks.

- Derive $E[T]$, $\text{Var}(T)$, and σ_T in terms of k and r .
- State the validity conditions on k and r , and explain why they are sufficient.
- How do k and r separately affect center and spread?
- Compare shape with the symmetric triangular model having the same endpoints k and $k + 3r$. Is this model symmetric, left-skewed, or right-skewed?
- For fixed k , compare $r = 2$ and $r = 5$: discuss changes in mean, variance, and skewness direction.

C4

For $\text{Triangular}(a, c, b)$:

$$E[T] = \frac{a + b + c}{3}, \quad \text{Var}(T) = \frac{a^2 + b^2 + c^2 - ab - ac - bc}{18}.$$

With $a = k$, $c = k + r$, $b = k + 3r$:

$$E[T] = \frac{k + (k + r) + (k + 3r)}{3} = k + \frac{4r}{3}.$$

For variance,

$$\text{Var}(T) = \frac{k^2 + (k + 3r)^2 + (k + r)^2 - k(k + 3r) - k(k + r) - (k + 3r)(k + r)}{18} = \frac{7r^2}{18}.$$


Thus

$$\sigma_T = \sqrt{\frac{7r^2}{18}} = r\sqrt{\frac{7}{18}}.$$

Validity requires $a \leq c \leq b$, i.e.

$$k \leq k + r \leq k + 3r,$$

which holds whenever $r \geq 0$; with $r > 0$, spread is nonzero, so the model is valid and non-degenerate for any real k . Parameter k shifts location only. Parameter r increases both mean offset ($\frac{4r}{3}$) and spread ($\propto r^2$). For endpoints k and $k + 3r$, symmetry would need mode at midpoint $k + \frac{3r}{2}$, but here mode is $k + r$, left of midpoint; therefore the model is right-skewed. Comparing $r = 2$ and $r = 5$ (fixed k): mean increases by $\frac{4}{3}(5 - 2) = 4$, variance scales from $\frac{28}{18}$ to $\frac{175}{18}$, and skewness direction stays right-skewed.



16 Normal with Parameters

Problem. The daily amount N (GB) of network data used by a class is modeled by

$$N \sim N(50 + 3p, (2 + q)^2), \quad p \in \mathbb{R}, \quad q > -2.$$

Questions/Tasks.

- Derive $E[N]$, $\text{Var}(N)$, and σ_N in terms of p and q .
- Explain the role of p and q : which controls location and which controls spread?
- Compare Case A: $(p, q) = (0, 1)$ and Case B: $(p, q) = (-2, 1)$. What changes in center, concentration around the mean, and tail likelihood?
- Compare Case C: $(p, q) = (0, 1)$ and Case D: $(p, q) = (0, 4)$. What changes in center, spread, concentration near the mean, and likelihood of values far from the mean?
- Between models $M_1 : (p, q) = (1, 0)$ and $M_2 : (p, q) = (-1, 3)$, which model is more likely to produce values at least 15 GB above its own mean? Justify conceptually.

C4

For $N(\mu, \sigma^2)$, $E[N] = \mu$, $\text{Var}(N) = \sigma^2$, $\sigma > 0$. So

$$E[N] = 50 + 3p, \quad \text{Var}(N) = (2 + q)^2, \quad \sigma_N = 2 + q \quad (q > -2).$$

Parameter p changes only location (horizontal shift). Parameter q changes spread: larger q gives larger σ , wider curve, lower peak, and heavier tails. Case A $\rightarrow$ B keeps $q = 1$, so $\sigma = 3$ is unchanged: concentration and tail behavior relative to each mean stay the same, but mean shifts from 50 to 44 (left by 6). Case C $\rightarrow$ D keeps $p = 0$, so mean stays 50, but σ increases from 3 to 6: values are less concentrated near 50 and far-from-mean values become more likely. For M_1 , $\sigma = 2$; for M_2 , $\sigma = 5$. A value 15 GB above its own mean is $15/\sigma$ standard deviations above mean: 7.5σ for M_1 versus 3σ for M_2 . The 3σ event is much less extreme, so M_2 is more likely.

