

Package ‘BANMA’

August 26, 2022

Type Package

Title Bayesian Estimation and Prediction for Network Meta-Analysis with Non-Informative Priors

Version 1.1-1

Date 2022-07-26

Author Hisashi Noma

Maintainer Hisashi Noma <noma@ism.ac.jp>

Description

Bayesian estimation and prediction for network meta-analysis with non-informative prior distributions are implementable. Various proper and improper prior distributions are available. The posterior sampling does not require formal convergence checks of MCMC.

Depends R (>= 3.5.0)

Imports stats, MASS, invgamma

License GPL-3

Encoding UTF-8

LazyData true

RoxygenNote 7.1.1

R topics documented:

BANMA-package	2
data.edit	2
diabetes	3
rinvGamma	3
rjeffreys	4
rmu	5
rmvmeta	6
runif_tau	7
runif_tau2	8

Index	10
--------------	-----------

BANMA-package	<i>The 'BANMA' package.</i>
---------------	-----------------------------

Description

Bayesian Estimation and Prediction for Network Meta-Analysis with Non-Informative Priors.

References

Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

data.edit	<i>Transforming arm-level data to contrast-based summary statistics</i>
-----------	---

Description

Transforming arm-level data to contrast-based summary statistics.

Usage

```
data.edit(study, trt, d, n)
```

Arguments

study	Study ID
trt	Numbered treatment (=1,2,...)
d	Number of events
n	Sample size

Value

Contrast-based summary statistics are generated.

- y: Contrast-based summary estimates.
- S: Vectored within-study covariance matrix.

Examples

```
data(diabetes)
attach(diabetes)

edat <- data.edit(study, trt, d, n)
```

diabetes

*Elliott and Mayer (2007)'s network meta-analysis data***Description**

- study: Study ID
- trt: Numbered treatment (1:Diuretic, 2:ACE inhibitor, 3:ARB, 4:Beta blocker, 5:Calcium-channel blocker, 6:Placebo)
- n: Sample size
- d: Number of events

Usage

```
data(diabetes)
```

Format

A arm-based dataset with 22 rows and 4 variables

References

Elliott, W. J., and Meyer, P. M. (2007). Incident diabetes in clinical trials of antihypertensive drugs: a network meta-analysis. *Lancet*. **369**: 201-207.

rinvGamma

*Posterior sampling of tau2 and tau for the multivariate random-effects model using the inverse-gamma prior distribution***Description**

Posterior sampling of tau2 and tau for the multivariate random-effects model using the inverse-gamma prior distribution.

Usage

```
rinvGamma(y, S, shape=0.001, scale=0.001, B=10000, n.grid=10^4)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y
B	Number of posterior samples to be generated.
shape	Shape parameter of the prior inverse-gamma distribution (default:0.001).
scale	Scale parameter of the prior inverse-gamma distribution (default:0.001).
n.grid	Number of grids for the numerical integration (default:10000).

Value

Posterior samples of tau2 and tau and their summaries.

- tau2: Posterior samples of tau2.
- tau: Posterior samples of tau.
- post.summary: Summary of the posterior samples of tau2 and tau.

References

Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaries
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

# Generating the posterior samples of tau2
R1 <- rinvgamma(y,S,B=20000)
R1$post.summary
```

rjeffreys

Posterior sampling of tau2 and tau for the multivariate random-effects model using the Jeffreys prior

Description

Posterior sampling of tau2 and tau for the multivariate random-effects model using the Jeffreys prior.

Usage

```
rjeffreys(y, S, B=10000, n.grid=10^4)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y
B	Number of posterior samples to be generated.
n.grid	Number of grids for the numerical integration (default:10000).

Value

Posterior samples of tau2 and tau and their summaries.

- tau2: Posterior samples of tau2.
- tau: Posterior samples of tau.
- post.summary: Summary of the posterior samples of tau2 and tau.

References

Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaryies
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

# Generating the posterior samples of tau2
R1 <- rjeffreys(y,S,B=20000)
R1$post.summary
```

rmu

Posterior sampling of mu and theta_new for the multivariate random-effects model

Description

Posterior sampling of mu and theta_new for the multivariate random-effects model.

Usage

```
rmu(y, S, rtau2)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y
rtau2	Posterior samples from the marginal posterior distribution of tau ² .

Value

Posterior samples of mu and theta_new and their summaries.

- mu: Posterior samples of mu.
- theta_new: Posterior samples of theta_new.
- mu_summary: Summary of the posterior samples of mu.
- theta.new_summary: Summary of the posterior samples of theta_new.

References

Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaries
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

# Generating the posterior samples of tau2
R1 <- runif_tau2(y,S,B=20000)
R1$post.summary

rtau2 <- R1$tau2

# Generating the posterior samples of mu and theta_new
# The number of samples are determined to be equal to that of rtau2.
R2 <- rmu(y,S,rtau2)
R2$mu_summary
R2$theta.new_summary
```

rmvmeta

The ordinary REML estimation of network meta-analysis

Description

The ordinary REML estimation of network meta-analysis.

Usage

```
rmvmeta(y, S)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y

Value

The REML estimates with 95% confidence intervals and ordinary t-approximation prediction intervals for network meta-analysis.

- Estimates: Restricted maximum likelihood (REML) estimates, their SE, and Wald-type 95% confidence intervals.
- Between-studies_SD: Between-studies SD estimate.
- 95%PI: 95% prediction intervals by the ordinary t-approximation.

References

- Cooper, H., Hedges, L. V., and Valentine, J. C. (2009). *The Handbook of Research Synthesis and Meta-Analysis*, 2nd edition. New York: Russell Sage Foundation.
- White, I. R. (2015). Network meta-analysis. *Stata Journal* **15**, 951-985.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaries
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

rmvmeta(y,S) # Results of the NMA analysis (log OR scale)
```

runif_tau	<i>Posterior sampling of tau2 and tau for the multivariate random-effects model using improper uniform prior for the scale of tau</i>
-----------	---

Description

Posterior sampling of tau2 and tau for the multivariate random-effects model for the scale of tau.

Usage

```
runif_tau(y, S, B=10000, n.grid=10^4)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y
B	Number of posterior samples to be generated.
n.grid	Number of grids for the numerical integration (default:10000).

Value

Posterior samples of tau2 and tau and their summaries.

- tau2: Posterior samples of tau2.
- tau: Posterior samples of tau.
- post.summary: Summary of the posterior samples of tau2 and tau.

References

- Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaries
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

# Generating the posterior samples of tau2
R1 <- runif_tau(y,S,B=20000)
R1$post.summary
```

runif_tau2	<i>Posterior sampling of tau2 and tau for the multivariate random-effects model using improper uniform prior for the scale of tau2</i>
------------	--

Description

Posterior sampling of tau2 and tau for the multivariate random-effects model for the scale of tau2.

Usage

```
runif_tau2(y, S, B=10000, n.grid=10^4)
```

Arguments

y	Contrast-based summary data of the outcome measure
S	Covariance estimates of y
B	Number of posterior samples to be generated.
n.grid	Number of grids for the numerical integration (default:10000).

Value

Posterior samples of tau2 and tau and their summaries.

- tau2: Posterior samples of tau2.
- tau: Posterior samples of tau.
- post.summary: Summary of the posterior samples of tau2 and tau.

References

Noma, H. (2022+). Bayesian estimation and prediction for network meta-analysis with non-informative priors. Forthcoming.

Examples

```
data(diabetes)
attach(diabetes)

# Transforming the arm-level data to the contrast-based summaries
edat <- data.edit(study,trt,d,n)

y <- edat$y
S <- edat$S

# Generating the posterior samples of tau2
R1 <- runif_tau2(y,S,B=20000)
R1$post.summary
```

Index

* **datasets**

diabetes, [3](#)

BANMA-package, [2](#)

data.edit, [2](#)

diabetes, [3](#)

rinvGamma, [3](#)

rjeffreys, [4](#)

rmu, [5](#)

rmvmeta, [6](#)

runif_tau, [7](#)

runif_tau2, [8](#)