

Hopf Module Algebras

Brandon Mather

UNT Master's

- (1939) Heinz Hopf works on homology of sphere groups
- (1969) Moss Sweedler writes seminal book “Hopf Algebras”
- (1986) Vladimir Drinfeld gives ICM address on quantum groups
- (1992) Susan Montgomery writes “Hopf Algebras and Their Actions on Rings”

Goal

To understand the actions of Hopf algebras on other algebras

Notation: $\mathbb{C}[v_1, \dots, v_n] = \mathbb{C} \langle v_1, \dots, v_n \mid v_j v_i - v_i v_j \rangle$

Quantum Polynomial Ring

Let $Q = (q_{ij})$ be an $n \times n$ matrix of roots of unity where

$$q_{ii} = 1 = q_{ji} q_{ij}.$$

A **quantum polynomial ring** is

$$\mathbb{C}_Q[v_1, \dots, v_n] = \mathbb{C} \langle v_1, \dots, v_n \mid v_j v_i - q_{ij} v_i v_j \rangle.$$

Example: $\mathbb{C}_{-1}[v_1, v_2] = \mathbb{C} \langle v_1, v_2 \mid v_1 v_2 + v_2 v_1 \rangle$

Motivation

- 1 When a grp G acts on a space V over \mathbb{C} linearly, the action can be extended to $V \otimes V$ by $g \in G$ acting as

$$g \otimes g = \Delta(g).$$

Then Δ defines a coproduct map

$$\Delta : \mathbb{C}G \rightarrow \mathbb{C}G \otimes \mathbb{C}G.$$

For arbitrary coproducts,

$$\Delta : A \rightarrow A \otimes A,$$

we call $g \in A$ **grouplike** if

$$\Delta(g) = g \otimes g.$$

Motivation

- 1 When a grp G acts on a space V over \mathbb{C} linearly, the action can be extended to $V \otimes V$ by $g \in G$ acting as

$$g \otimes g = \Delta(g).$$

Then Δ defines a coproduct map

$$\Delta : \mathbb{C}G \rightarrow \mathbb{C}G \otimes \mathbb{C}G.$$

For arbitrary coproducts,

$$\Delta : A \rightarrow A \otimes A,$$

we call $g \in A$ **grouplike** if

$$\Delta(g) = g \otimes g.$$

- 2 When a Lie alg \mathfrak{g} acts on a space V over \mathbb{C} , the action can be extended to $V \otimes V$ by $x \in \mathfrak{g}$ acting as

$$x \otimes 1 + 1 \otimes x = \Delta(x).$$

Then Δ defines a coproduct map

$$\Delta : \mathfrak{g} \rightarrow \mathfrak{g} \otimes \mathfrak{g}.$$

For arbitrary coproducts,

$$\Delta : A \rightarrow A \otimes A,$$

we call $x \in A$ **primitive** if

$$\Delta(x) = x \otimes 1 + 1 \otimes x.$$

Hopf algebras combine the notions of Algebras and Coalgebras that have coproducts. We are looking for actions of Hopf algebras. In particular on non-commutative algebras like Quantum Polynomial Rings.

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

with operations $\Delta : H_4 \rightarrow H_4 \otimes H_4, \varepsilon : H_4 \rightarrow \mathbb{C}, S : H_4 \rightarrow H_4$

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x$$

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

with operations $\Delta : H_4 \rightarrow H_4 \otimes H_4, \varepsilon : H_4 \rightarrow \mathbb{C}, S : H_4 \rightarrow H_4$

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x$$

$$\varepsilon(g) = 1, \quad \varepsilon(x) = 0$$

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

with operations $\Delta : H_4 \rightarrow H_4 \otimes H_4, \varepsilon : H_4 \rightarrow \mathbb{C}, S : H_4 \rightarrow H_4$

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x$$

$$\varepsilon(g) = 1, \quad \varepsilon(x) = 0$$

$$S(g) = g^{-1}, \quad S(x) = -x$$

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

with operations $\Delta : H_4 \rightarrow H_4 \otimes H_4, \varepsilon : H_4 \rightarrow \mathbb{C}, S : H_4 \rightarrow H_4$

$$\Delta(g) = g \otimes g,$$

$$\Delta(x) = x \otimes 1 + g \otimes x$$

$$\varepsilon(g) = 1,$$

$$\varepsilon(x) = 0$$

$$S(g) = g^{-1},$$

$$S(x) = -x$$

Group-like

$(1, g)$ – Primitive

Sweedler's Algebra

The unique 4-dim'l non-commutative, non-cocommutative Hopf alg given by M. Sweedler (1969):

$$H_4 = \langle g, x \mid g^2 = 1, x^2 = 0, gx = -xg \rangle$$

with operations $\Delta : H_4 \rightarrow H_4 \otimes H_4, \varepsilon : H_4 \rightarrow \mathbb{C}, S : H_4 \rightarrow H_4$

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x$$

$$\varepsilon(g) = 1, \quad \varepsilon(x) = 0$$

$$S(g) = g^{-1}, \quad S(x) = -x$$

Group-like

$(1, g)$ – Primitive

Let τ be the 'flip' over the tensor product, so $\tau(u \otimes v) = v \otimes u$.
Note that $\tau \circ \Delta(x) \neq \Delta(x)$, so H is non-cocommutative.

H_4 acts on $\mathbb{C}_{-1}[v_1, v_2]$ by

$$g \cdot v_1 = v_1, \quad g \cdot v_2 = -v_2, \quad x \cdot v_1 = 0, \quad x \cdot v_2 = v_1.$$

H_4 acts on $\mathbb{C}_{-1}[v_1, v_2]$ by

$$g \cdot v_1 = v_1, \quad g \cdot v_2 = -v_2, \quad x \cdot v_1 = 0, \quad x \cdot v_2 = v_1.$$

We can express this action on the generators as

$$g \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad x \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

Kac-Paljutkin Algebra

The unique 8-dim'l non-comm., non-cocomm. Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\left\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \right\rangle$$

with operations

Kac-Paljutkin Algebra

The unique 8-dim'l non-comm., non-cocomm. Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\left\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \right\rangle$$

with operations

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

Kac-Paljutkin Algebra

The unique 8-dim'l non-comm., non-cocomm. Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \rangle$$

with operations

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

Kac-Paljutkin Algebra

The unique 8-dim'l non-comm., non-cocomm. Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\left\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \right\rangle$$

with operations

$$\Delta(x) = x \otimes x, \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

$$S(x) = x, S(y) = y, S(z) = z.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[v_1, v_2]$ where $q^2 = -1$ via the representation

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[v_1, v_2]$ where $q^2 = -1$ via the representation

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_Q[v_1, v_2, v_3, v_4]$ for

$q_{12} = q_{34}^{-1}$, $q_{13} = q_{24}^{-1}$, $q_{14}^2 = 1$, $q_{23}^2 = -1$ via the rep

$$x \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[v_1, v_2]$ where $q^2 = -1$ via the representation

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_Q[v_1, v_2, v_3, v_4]$ for

$q_{12} = q_{34}^{-1}$, $q_{13} = q_{24}^{-1}$, $q_{14}^2 = 1$, $q_{23}^2 = -1$ via the rep

$$x \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_{-1}[v_1, v_2]$ via the rep

$$x \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Quantized Universal Enveloping Algebra

Described by P. Kulish and N. Reshetikhin in “Quantum linear problem...” (1983), leading Vladimir Drinfeld to quantum groups $\mathcal{U}_q(\mathfrak{sl}_2) =$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

Quantized Universal Enveloping Algebra

Described by P. Kulish and N. Reshetikhin in “Quantum linear problem...” (1983), leading Vladimir Drinfeld to quantum groups $\mathcal{U}_q(\mathfrak{sl}_2) =$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \quad \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \quad \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

Quantized Universal Enveloping Algebra

Described by P. Kulish and N. Reshetikhin in “Quantum linear problem...” (1983), leading Vladimir Drinfeld to quantum groups $\mathcal{U}_q(\mathfrak{sl}_2) =$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \varepsilon(K) = \varepsilon(K^{-1}) = 1,$$

Quantized Universal Enveloping Algebra

Described by P. Kulish and N. Reshetikhin in “Quantum linear problem...” (1983), leading Vladimir Drinfeld to quantum groups $\mathcal{U}_q(\mathfrak{sl}_2) =$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \varepsilon(K) = \varepsilon(K^{-1}) = 1,$$

$$S(E) = -K^{-1}E, S(F) = -FK, S(K) = K^{-1}, S(K^{-1}) = K.$$

Quantized Universal Enveloping Algebra

Described by P. Kulish and N. Reshetikhin in “Quantum linear problem...” (1983), leading Vladimir Drinfeld to quantum groups $\mathcal{U}_q(\mathfrak{sl}_2) =$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \quad \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \quad \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \quad \varepsilon(K) = \varepsilon(K^{-1}) = 1,$$

$$S(E) = -K^{-1}E, \quad S(F) = -FK, \quad S(K) = K^{-1}, \quad S(K^{-1}) = K.$$

Note: You can recover $\mathcal{U}(\mathfrak{sl}_2)$ by limiting $q \rightarrow 1$.

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[v_1, v_2]$ by the representation

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[v_1, v_2]$ by the representation

$$E \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$F \mapsto \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[v_1, v_2]$ by the representation

$$E \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$K \mapsto \begin{bmatrix} q & 0 \\ 0 & q^{-1} \end{bmatrix}$$

$$F \mapsto \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

$$K^{-1} \mapsto \begin{bmatrix} q^{-1} & 0 \\ 0 & q \end{bmatrix}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode $S : H \rightarrow H$ where the bialgebra operations are

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$1_H : \mathbb{C} \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ 1_H \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes 1_H & \\ \mathbb{C} \otimes H & & H & & H \otimes \mathbb{C} \\ \searrow = & & \nwarrow = & & \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$\Delta : H \rightarrow H \otimes H,$$

$$1_H : \mathbb{C} \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Coassociativity:

$$\begin{array}{ccc} H & \xrightarrow{\Delta} & H \otimes H \\ \Delta \downarrow & & \downarrow \Delta \otimes id \\ H \otimes H & \xrightarrow{id \otimes \Delta} & H \otimes H \otimes H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ 1_H \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes 1_H & \\ \mathbb{C} \otimes H & & H & & H \otimes \mathbb{C} \\ \searrow = & & & \swarrow = & \\ & & H & & \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$1_H : \mathbb{C} \rightarrow H,$$

$$\Delta : H \rightarrow H \otimes H,$$

$$\varepsilon : H \rightarrow \mathbb{C}$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Coassociativity:

$$\begin{array}{ccc} H & \xrightarrow{\Delta} & H \otimes H \\ \Delta \downarrow & & \downarrow \Delta \otimes id \\ H \otimes H & \xrightarrow{id \otimes \Delta} & H \otimes H \otimes H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ 1_H \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes 1_H & \\ \mathbb{C} \otimes H & & & & H \otimes \mathbb{C} \\ & \searrow = & H & \swarrow = & \end{array}$$

Counit:

$$\begin{array}{ccccc} & & H & & \\ \varepsilon \otimes id \nwarrow & & \downarrow \Delta & \nearrow id \otimes \varepsilon & \\ \mathbb{C} \otimes H & & H \otimes H & & H \otimes \mathbb{C} \\ & \nwarrow = & & \swarrow = & \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad \nabla \quad} & H & \xrightarrow{\quad \Delta \quad} & H \otimes H \\
 \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{\quad id \otimes \tau \otimes id \quad} & H \otimes H \otimes H \otimes H & &
 \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\nabla} & H & \xrightarrow{\Delta} & H \otimes H \\
 \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{id \otimes \tau \otimes id} & H \otimes H \otimes H \otimes H & &
 \end{array}$$

Unit and Counit compatibility:

$$\begin{array}{ccc}
 \mathbb{C} & \xrightarrow{1_H} & H \\
 1_H \otimes 1_H \searrow & & \downarrow \Delta \\
 & & H \otimes H
 \end{array}
 \qquad
 \begin{array}{ccc}
 H & \xrightarrow{\epsilon} & \mathbb{C} \\
 \nabla \uparrow & \nearrow \epsilon \otimes \epsilon & \\
 H \otimes H & &
 \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad \nabla \quad} & H & \xrightarrow{\quad \Delta \quad} & H \otimes H \\
 \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{\quad id \otimes \tau \otimes id \quad} & H \otimes H \otimes H \otimes H & &
 \end{array}$$

Unit and Counit compatibility:

$$\begin{array}{ccc}
 \mathbb{C} & \xrightarrow{1_H} & H \\
 1_H \otimes 1_H \searrow & & \downarrow \Delta \\
 & & H \otimes H
 \end{array}
 \qquad
 \begin{array}{ccc}
 H & \xrightarrow{\epsilon} & \mathbb{C} \\
 \nabla \uparrow & \nearrow \epsilon \otimes \epsilon & \\
 H \otimes H & &
 \end{array}$$

Antipode:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad id \otimes S \quad} & H \otimes H & & \\
 \Delta \uparrow & & \downarrow \nabla & & \\
 H & \xrightarrow{\quad \epsilon \quad} & \mathbb{C} & \xrightarrow{1_H} & H \\
 \Delta \downarrow & & & & \uparrow \nabla \\
 H \otimes H & \xrightarrow{\quad S \otimes id \quad} & H \otimes H & &
 \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$.

Then we say H **acts** on A by α if the following diagrams commute:

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$.

Then we say H **acts** on A by α if the following diagrams commute:

$$\begin{array}{ccc} H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\ \downarrow id \otimes \alpha & & \downarrow \alpha \\ H \otimes A & \xrightarrow{\alpha} & A \end{array} \quad \begin{array}{ccc} \mathbb{C} \otimes A & \xrightarrow{1_H \otimes id} & H \otimes A \\ & \searrow = & \downarrow \alpha \\ & & A \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$.

Then we say H **acts** on A by α if the following diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{1_H \otimes id} & H \otimes A \\
 & \searrow = & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called an H -**module algebra** if the following also commute:

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$.
Then we say H **acts** on A by α if the following diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 C \otimes A & \xrightarrow{1_H \otimes id} & H \otimes A \\
 & \searrow = & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called an **H -module algebra** if the following also commute:

$$\begin{array}{ccccc}
 H \otimes A \otimes A & \xrightarrow{id \otimes \nabla} & H \otimes A & \xrightarrow{\alpha} & A \xleftarrow{\nabla} A \otimes A \\
 \Delta \otimes id \otimes id \downarrow & & & & \uparrow \alpha \otimes \alpha \\
 H \otimes H \otimes A \otimes A & \xrightarrow{id \otimes \tau \otimes id} & H \otimes A \otimes H \otimes A & &
 \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$.
Then we say H **acts** on A by α if the following diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{1_H \otimes id} & H \otimes A \\
 & \searrow = & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called an **H -module algebra** if the following also commute:

$$\begin{array}{ccccc}
 H \otimes A \otimes A & \xrightarrow{id \otimes \nabla} & H \otimes A & \xrightarrow{\alpha} & A \xleftarrow{\nabla} A \otimes A \\
 \Delta \otimes id \otimes id \downarrow & & & & \uparrow \alpha \otimes \alpha \\
 H \otimes H \otimes A \otimes A & \xrightarrow{id \otimes \tau \otimes id} & H \otimes A \otimes H \otimes A & &
 \end{array}$$

$$\begin{array}{ccc}
 H & \xrightarrow{\varepsilon} & \mathbb{C} \xrightarrow{1_A} A \\
 id \otimes 1_A \downarrow & & \nearrow \alpha \\
 H \otimes A & &
 \end{array}$$

In words, H **acts on** A iff you can multiply in H and then act on A or act on A consecutively, $\forall h, h' \in H, \forall a \in A$

$$(hh')(a) = h(h'(a)), \quad 1_H(a) = a.$$

And A is an H -**module alg** iff H acts on A and $\forall h \in H, \forall a, a' \in A$

$$h(aa') = \sum h_i(a) \cdot h_j(a'), \quad h(1_A) = \varepsilon(h)1_A$$

where $\Delta(h) = \sum h_i \otimes h_j$.

Semidirect Product

Let G and G' be groups where G' acts on G by automorphisms, giving the semidirect product group, $G \rtimes G'$.

The action can be extended to the group algebras:

$$\mathbb{C}(G \rtimes G') = \mathbb{C}G \# \mathbb{C}G'$$

with product $g'g = g'(g)g'$.

Smash Product Algebra

If H is a Hopf algebra and A an H -module algebra, then $A \# H$ is defined as $A \otimes H$ as a vector space and with product

$$(a' \otimes h)(a \otimes h') = \sum_i a' h_{i1}(a) \otimes h_{i2} h'$$

where $a, a' \in A$, $h, h' \in H$ and $\Delta(h) = \sum_i h_{i1} \otimes h_{i2}$.

Smash Product Algebra

"Group-like" and "Lie-like"

For Hopf alg H , define

$$G(H) = \{h \in H \mid \Delta(h) = h \otimes h\} = \text{grouplike elements}$$

which forms a group, and define

$$P(H) = \{h \in H \mid \Delta(h) = h \otimes 1 + 1 \otimes h\} = \text{primitive elements}$$

which forms a Lie alg.

Cartier-Kostant-Milnor-Moore Theorem

Let H be a cocommutative Hopf algebra over \mathbb{C} , then as Hopf alg,

$$H \cong \mathcal{U}(P(H)) \# \mathbb{C}G(H)$$

Cor: Any cocomm, finite-dim'l Hopf alg over \mathbb{C} is iso to a grp alg.

- Which actions on algebras by Hopf algebras factor through a group action? If the action does not factor through a group action, it is said to have "quantum symmetry".

- Which actions on algebras by Hopf algebras factor through a group action? If the action does not factor through a group action, it is said to have "quantum symmetry".
- Which Hopf Algebras act on AS-regular algebras?

- Which actions on algebras by Hopf algebras factor through a group action? If the action does not factor through a group action, it is said to have "quantum symmetry".
- Which Hopf Algebras act on AS-regular algebras?
- When are the invariant subrings from Hopf actions AS-Gorenstein?

- Which actions on algebras by Hopf algebras factor through a group action? If the action does not factor through a group action, it is said to have "quantum symmetry".
- Which Hopf Algebras act on AS-regular algebras?
- When are the invariant subrings from Hopf actions AS-Gorenstein?
- If H is semisimple and finite-dimensional, and A is semiprime, is $A \# H$ semiprime?

- Which actions on algebras by Hopf algebras factor through a group action? If the action does not factor through a group action, it is said to have "quantum symmetry".
- Which Hopf Algebras act on AS-regular algebras?
- When are the invariant subrings from Hopf actions AS-Gorenstein?
- If H is semisimple and finite-dimensional, and A is semiprime, is $A \# H$ semiprime?
- If B is a Koszul algebra, are there nontrivial PBW deformations of $B \# \mathcal{U}_q(\mathfrak{sl}_2)$?