

Hopf Algebra Actions on Quantum Planes

Brandon Mather

Algebra Seminar November 2023

Kac-Paljutkin Algebra

The unique 8-dim'l non-commutative, non-cocommutative Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \rangle$$

with operations

Kac-Paljutkin Algebra

The unique 8-dim'l non-commutative, non-cocommutative Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \rangle$$

with operations

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

Kac-Paljutkin Algebra

The unique 8-dim'l non-commutative, non-cocommutative Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \rangle$$

with operations

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

Kac-Paljutkin Algebra

The unique 8-dim'l non-commutative, non-cocommutative Hopf alg given by G. Kac and V. Paljutkin in "Finite Ring Groups" (1966):

$$H_8 =$$

$$\langle x, y, z \mid x^2 = y^2 = 1, xy = yx, zx = yz, zy = xz, z^2 = \frac{1}{2}(1+x+y-xy) \rangle$$

with operations

$$\Delta(x) = x \otimes x, \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

$$S(x) = x, S(y) = y, S(z) = z.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[x, y]$ where $q^2 = -1$ by

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[x, y]$ where $q^2 = -1$ by

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_Q[x_1, x_2, x_3, x_4]$ for

$q_{12} = q_{34}^{-1}$, $q_{13} = q_{24}^{-1}$, $q_{14}^2 = 1$, $q_{23}^2 = -1$ by

$$x \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

Actions of Kac-Paljutkin Algebra

H_8 acts on $\mathbb{C}_q[x, y]$ where $q^2 = -1$ by

$$x \mapsto \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_Q[x_1, x_2, x_3, x_4]$ for

$q_{12} = q_{34}^{-1}$, $q_{13} = q_{24}^{-1}$, $q_{14}^2 = 1$, $q_{23}^2 = -1$ by

$$x \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

And on $\mathbb{C}_{-1}[u, v]$ by

$$x \mapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad y \mapsto \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \quad z \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

16-dim Hopf Algebra

The 16-dimensional semisimple Hopf algebras have been classified by Kashina. One such example is

$$H_{16} = \langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, zx = xyz, zy = yz \rangle$$

with operations:

16-dim Hopf Algebra

The 16-dimensional semisimple Hopf algebras have been classified by Kashina. One such example is

$$H_{16} = \langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, zx = xyz, zy = yz \rangle$$

with operations:

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x^2 + y \otimes 1 - y \otimes x^2)(z \otimes z),$$

16-dim Hopf Algebra

The 16-dimensional semisimple Hopf algebras have been classified by Kashina. One such example is

$$H_{16} = \langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, zx = xyz, zy = yz \rangle$$

with operations:

$$\Delta(x) = x \otimes x, \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x^2 + y \otimes 1 - y \otimes x^2)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

16-dim Hopf Algebra

The 16-dimensional semisimple Hopf algebras have been classified by Kashina. One such example is

$$H_{16} = \langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, zx = xyz, zy = yz \rangle$$

with operations:

$$\Delta(x) = x \otimes x, \Delta(y) = y \otimes y,$$

$$\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x^2 + y \otimes 1 - y \otimes x^2)(z \otimes z),$$

$$\varepsilon(x) = \varepsilon(y) = \varepsilon(z) = 1,$$

$$S(x) = x^3, S(y) = y, S(z) = \frac{1}{2}(1 + x^2 + y - x^2y)z.$$

Action of H_{16}

H_{16} acts on the algebra

$$\mathbb{C}[t, u, v, w]/(tw + wt, uw + wu)$$

by the representation

$$x \mapsto \begin{bmatrix} i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}, \quad y \mapsto \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix},$$

$$z \mapsto \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Quantized Universal Enveloping Algebra

Described by Piotr Kulish and Nicolai Reshetikhin in “Quantum linear problem for the sine-Gordon equation and highest weight representations” (1983), leading Vladimir Drinfeld to quantum groups

$$\mathcal{U}_q(\mathfrak{sl}_2) =$$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

Quantized Universal Enveloping Algebra

Described by Piotr Kulish and Nicolai Reshetikhin in “Quantum linear problem for the sine-Gordon equation and highest weight representations” (1983), leading Vladimir Drinfeld to quantum groups

$$\mathcal{U}_q(\mathfrak{sl}_2) =$$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

Quantized Universal Enveloping Algebra

Described by Piotr Kulish and Nicolai Reshetikhin in “Quantum linear problem for the sine-Gordon equation and highest weight representations” (1983), leading Vladimir Drinfeld to quantum groups

$$\mathcal{U}_q(\mathfrak{sl}_2) =$$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \varepsilon(K) = \varepsilon(K^{-1}) = 1,$$

Quantized Universal Enveloping Algebra

Described by Piotr Kulish and Nicolai Reshetikhin in “Quantum linear problem for the sine-Gordon equation and highest weight representations” (1983), leading Vladimir Drinfeld to quantum groups

$$\mathcal{U}_q(\mathfrak{sl}_2) =$$

$$\left\langle E, F, K, K^{-1} \mid EF - FE = (q - q^{-1})^{-1} (K - K^{-1}), KEK^{-1} = q^2 E, \right. \\ \left. KFK^{-1} = q^{-2} F, KK^{-1} = K^{-1}K = 1 \right\rangle$$

with operations:

$$\Delta(E) = E \otimes 1 + K \otimes E, \Delta(F) = F \otimes K^{-1} + 1 \otimes F,$$

$$\Delta(K) = K \otimes K, \Delta(K^{-1}) = K^{-1} \otimes K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \varepsilon(K) = \varepsilon(K^{-1}) = 1,$$

$$S(E) = -K^{-1}E, S(F) = -FK, S(K) = K^{-1}, S(K^{-1}) = K.$$

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[x, y]$ with $q^2 = -1$ by the representation

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[x, y]$ with $q^2 = -1$ by the representation

$$E \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad F \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

Actions of $\mathcal{U}_q(\mathfrak{sl}_2)$

$\mathcal{U}_q(\mathfrak{sl}_2)$ acts on $\mathbb{C}_q[x, y]$ with $q^2 = -1$ by the representation

$$E \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad F \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$K \mapsto \begin{bmatrix} q & 0 \\ 0 & q^{-1} \end{bmatrix} \quad K^{-1} \mapsto \begin{bmatrix} q^{-1} & 0 \\ 0 & q \end{bmatrix}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode $S : H \rightarrow H$ where the bialgebra operations are

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$\eta : \mathbb{C} \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ \eta \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes \eta & \\ \mathbb{C} \otimes H & & H & & H \otimes \mathbb{C} \\ \searrow = & & & \swarrow = & \\ & & H & & \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$\Delta : H \rightarrow H \otimes H,$$

$$\eta : \mathbb{C} \rightarrow H,$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Coassociativity:

$$\begin{array}{ccc} H & \xrightarrow{\Delta} & H \otimes H \\ \Delta \downarrow & & \downarrow \Delta \otimes id \\ H \otimes H & \xrightarrow{id \otimes \Delta} & H \otimes H \otimes H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ \eta \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes \eta & \\ \mathbb{C} \otimes H & & H & & H \otimes \mathbb{C} \\ \searrow = & & & \swarrow = & \\ & & H & & \end{array}$$

Hopf Algebra

A **Hopf algebra** is a bialgebra H over a field with an antipode

$S : H \rightarrow H$ where the bialgebra operations are

$$\nabla : H \otimes H \rightarrow H,$$

$$\Delta : H \rightarrow H \otimes H,$$

$$\eta : \mathbb{C} \rightarrow H,$$

$$\varepsilon : H \rightarrow \mathbb{C}$$

so that the following commute:

Associativity:

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{\nabla \otimes id} & H \otimes H \\ id \otimes \nabla \downarrow & & \downarrow \nabla \\ H \otimes H & \xrightarrow{\nabla} & H \end{array}$$

Coassociativity:

$$\begin{array}{ccc} H & \xrightarrow{\Delta} & H \otimes H \\ \Delta \downarrow & & \downarrow \Delta \otimes id \\ H \otimes H & \xrightarrow{id \otimes \Delta} & H \otimes H \otimes H \end{array}$$

Unit:

$$\begin{array}{ccccc} & & H \otimes H & & \\ \eta \otimes id \nearrow & & \downarrow \nabla & \nwarrow id \otimes \eta & \\ \mathbb{C} \otimes H & & H & & H \otimes \mathbb{C} \\ \searrow = & & \nwarrow = & & \end{array}$$

Counit:

$$\begin{array}{ccccc} & & H & & \\ = \nwarrow & & \downarrow \Delta & \nearrow = & \\ \mathbb{C} \otimes H & & H \otimes H & & H \otimes \mathbb{C} \\ \nwarrow \varepsilon \otimes id & & \nearrow id \otimes \varepsilon & & \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccc}
 H \otimes H & \xrightarrow{\quad \nabla \quad} & H \xrightarrow{\quad \Delta \quad} H \otimes H \\
 \Delta \otimes \Delta \downarrow & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{\quad id \otimes \tau \otimes id \quad} & H \otimes H \otimes H \otimes H
 \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad \nabla \quad} & H & \xrightarrow{\quad \Delta \quad} & H \otimes H \\
 \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{\quad id \otimes \tau \otimes id \quad} & H \otimes H \otimes H \otimes H & &
 \end{array}$$

Unit and Counit compatibility:

$$\begin{array}{ccc}
 \mathbb{C} & \xrightarrow{\eta} & H \\
 \eta \otimes \eta \searrow & & \downarrow \Delta \\
 & & H \otimes H
 \end{array}
 \qquad
 \begin{array}{ccc}
 H & \xrightarrow{\varepsilon} & \mathbb{C} \\
 \nabla \uparrow & & \nearrow \varepsilon \otimes \varepsilon \\
 H \otimes H & &
 \end{array}$$

Hopf Algebra Diagrams

Product and Coproduct compatibility:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad \nabla \quad} & H & \xrightarrow{\quad \Delta \quad} & H \otimes H \\
 \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\
 H \otimes H \otimes H \otimes H & \xrightarrow{\quad id \otimes \tau \otimes id \quad} & H \otimes H \otimes H \otimes H & &
 \end{array}$$

Unit and Counit compatibility:

$$\begin{array}{ccc}
 \mathbb{C} & \xrightarrow{\eta} & H \\
 \eta \otimes \eta \searrow & & \downarrow \Delta \\
 & & H \otimes H
 \end{array}
 \qquad
 \begin{array}{ccc}
 H & \xrightarrow{\epsilon} & \mathbb{C} \\
 \nabla \uparrow & & \nearrow \epsilon \otimes \epsilon \\
 H \otimes H & &
 \end{array}$$

Antipode:

$$\begin{array}{ccccc}
 H \otimes H & \xrightarrow{\quad id \otimes S \quad} & H \otimes H & & \\
 \Delta \uparrow & & & & \downarrow \nabla \\
 H & \xrightarrow{\quad \epsilon \quad} & \mathbb{C} & \xrightarrow{\quad \eta \quad} & H \\
 \Delta \downarrow & & & & \uparrow \nabla \\
 H \otimes H & \xrightarrow{\quad S \otimes id \quad} & H \otimes H & &
 \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$. Then we say H **acts** on A by α if the following the diagrams commute:

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$. Then we say H **acts** on A by α if the following the diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{\eta \otimes id} & H \otimes A \\
 \searrow = & & \downarrow \alpha \\
 & & A
 \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$. Then we say H **acts** on A by α if the following the diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{\eta \otimes id} & H \otimes A \\
 \searrow = & & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called a **module algebra** if the following also commute:

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$. Then we say H **acts** on A by α if the following the diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{\eta \otimes id} & H \otimes A \\
 \searrow = & & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called a **module algebra** if the following also commute:

$$\begin{array}{ccccc}
 H \otimes A \otimes A & \xrightarrow{\nabla} & H \otimes A & \xrightarrow{\alpha} & A & \xleftarrow{\nabla} & A \otimes A \\
 \Delta \otimes id \otimes id \downarrow & & & & & & \uparrow \alpha \otimes \alpha \\
 H \otimes H \otimes A \otimes A & \xrightarrow{id \otimes \tau \otimes id} & H \otimes A \otimes H \otimes A & & & &
 \end{array}$$

Hopf Algebra Actions

Let H be a Hopf alg and A an alg with a map $\alpha : H \otimes A \rightarrow A$. Then we say H **acts** on A by α if the following the diagrams commute:

$$\begin{array}{ccc}
 H \otimes H \otimes A & \xrightarrow{\nabla \otimes id} & H \otimes A \\
 \downarrow id \otimes \alpha & & \downarrow \alpha \\
 H \otimes A & \xrightarrow{\alpha} & A
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C} \otimes A & \xrightarrow{\eta \otimes id} & H \otimes A \\
 \searrow = & & \downarrow \alpha \\
 & & A
 \end{array}$$

A is called a **module algebra** if the following also commute:

$$\begin{array}{ccc}
 H \otimes A \otimes A & \xrightarrow{\nabla} & H \otimes A \xrightarrow{\alpha} A \xleftarrow{\nabla} A \otimes A \\
 \Delta \otimes id \otimes id \downarrow & & \uparrow \alpha \otimes \alpha \\
 H \otimes H \otimes A \otimes A & \xrightarrow{id \otimes \tau \otimes id} & H \otimes A \otimes H \otimes A
 \end{array}$$

$$\begin{array}{ccc}
 H & \xrightarrow{\varepsilon} & \mathbb{C} \xrightarrow{\eta} A \\
 \eta \downarrow & \nearrow \alpha & \\
 H \otimes A & &
 \end{array}$$