



Waves:

- So far in this class we have focused on Newton's laws of motion for systems that we can describe as a single object.
- Now we turn to the equally important concept of waves.
- Perhaps the most familiar wave to many people are water waves.
- However, there are many other important physical processes that can also be modeled as waves: Sounds, light, cell phone signals, ...
- At the most basic level, the laws of physics tell us that everything must behave as a wave
- Waves and their properties will be the focus for most of the remaining semester.



The damped harmonic oscillator :

- The harmonic oscillator is defined by the linear restoring force: $F = -kx$
- The characteristic motion of the variation in the oscillators position as a function of time is given by

$$x(t) = A \cos(\omega_0 t)$$

A = amplitude

$$\omega_0 = \sqrt{\frac{k}{m}} = \text{angular frequency}$$

- The oscillators **natural frequency** f_0 and period T can both be defined from the angular frequency


$$f_0 = \frac{1}{T} = \frac{\omega_0}{2\pi}$$

- The natural frequency f_0 is the frequency of the oscillator with no external forces acting on it.



- With no external forces, the energy of the oscillator is conserved, with the energy constantly being exchanged between kinetic and potential energy.

- A good example of a system that behaves as a harmonic oscillator is the simple-pendulum created by hanging a mass from a massless string with length L



- If the angle θ is small (<10deg), then the angle of the mass from the vertical is given by

$$\theta(t) = \theta_{\max} \cos(\omega_0 t)$$

- The frequency of the pendulum is independent of the mass:

$$\omega_0 = \sqrt{\frac{L}{g}}$$

Friction :

- Of course, in the real world, there is always friction.
- To a good approximation, the frictional force is proportional to the velocity of the oscillator.



$$F_{\text{net}} = -kx - bv$$

b = fiction constant

friction term



- What happens to the motion of the harmonic oscillator when there is friction present?
- If the friction term is small, then we might guess that the motion will still be harmonic, but the amplitude will decrease
- This is correct. As a function of time, the motion can be expressed as

$$x(t) = \left(A e^{-\frac{bt}{2m}} \right) \cos(\omega_{\text{damp}} t)$$

- The frequency of the oscillator has been modified by the friction force


$$\boxed{\omega_{\text{damp}} = \sqrt{\frac{k}{m} - \frac{b^2}{4m^2}}}$$

- If there is no friction, then the energy of the oscillator is constant

$$E = 1/2kA^2$$

- Replacing the amplitude with the one found above:

$$\boxed{E = \frac{1}{2}kA^2 e^{-\frac{bt}{m}}}$$



Physics for Life Scientists (PHYS-183), Paul D. Nation

- Although the energy is lost exponentially, the period of the oscillation remains the same:

$$T = \frac{2\pi}{\omega_{\text{damp}}}$$

Ex. A 0.2kg mass is attached to a spring with a spring constant of $k=40\text{N/m}$ and with a friction constant $b=0.02\text{kg/s}$. Assuming the spring is initially stretched by 0.1m, what is the amount it takes for the oscillator to lose 1/2 of its energy?

- The energy as a function of time is given by $E(t) = \frac{1}{2}kA^2 e^{-\frac{bt}{m}} = E(0)e^{-\frac{bt}{m}}$

- If the energy has been reduced to 1/2 of its initial value then we have: $\frac{1}{2} = e^{-\frac{bt}{m}}$

- Solving for the time: $t = -\frac{m}{b} \log\left(\frac{1}{2}\right)$

- Plugging in the numbers: $t = \frac{0.2}{0.02} \log(2) = 10 \log(2) = \boxed{6.9\text{s}}$



Driven harmonic oscillators:

- In order to keep the amplitude constant, we must add energy to oscillator from some external force
- A harmonic oscillator that has an external force and friction acting on it is called a **damped-driven harmonic oscillator** (“damped”=friction)

$$F_{\text{net}} = -kx - bv + F_{\text{ext}}$$

- If the external force is sinusoidal with an angular frequency ω_{ext} then, after a sufficiently long time, the oscillator reaches a **steady-state** where the motion remains periodic

$$x(t) = A(\omega_0, \omega_{\text{ext}}) \cos(\omega_{\text{ext}}t + \phi)$$

- The amplitude of the oscillation depends on both the natural and external frequencies.
- There is also now a phase angle ϕ , indicating that the external force and spring force may not oscillate the same in time.
- In steady-state, the amplitude of the oscillation is fixed. This means that the energy lost due to friction is balanced by the energy gained from the external force.




Physics for Life Scientists (PHYS-183), Paul D. Nation

- How do we control the amount of energy from the external force to the oscillator?
- The amplitude of the external force will be important, but the frequency ω_{ext} of the external force is also important
- For a fixed amplitude force, the amount of energy input into the oscillator depends on how close the external frequency ω_{ext} is to the oscillators natural frequency ω_0 .
- Why? Because the power (energy/time) is given by $P = \vec{F}_{\text{ext}} \cdot \vec{v}_{\text{osc}}$
- If ω_{ext} and ω_0 are not close, then $\vec{F}_{\text{ext}} \cdot \vec{v}_{\text{osc}} \rightarrow 0$
- Mathematically, the amplitude is given by

$$A = \frac{F_{\text{ext}}/m}{\sqrt{(\omega_0^2 - \omega_{\text{ext}}^2)^2 + \left(\frac{b\omega_{\text{ext}}}{m}\right)^2}}$$

- The maximum occurs when $\omega_{\text{ext}} = \omega_0$
- This condition is called **resonance**.



Physics for Life Scientists (PHYS-183), Paul D. Nation



- When an oscillator is driven on resonance, the amplitude of the motion becomes extremely large.
- Changing a channel on the radio or television is an example of changing resonance frequencies.
- Resonance is particularly important for buildings and bridges.
- If the building experiences a force near one of the resonance frequencies of the building, then the building can shake itself apart.



Bridge driven on resonance by the wind.

Physics for Life Scientists (PHYS-183), Paul D. Nation

Wave concepts:

- There are many kinds of waves.
- Mechanical waves are vibrations that travel through a material such as water and air (sound waves), waves along a string (musical instruments), or seismic waves (from earthquakes).
- A general property of all waves (except electromagnetic waves) is that they travel through material with a characteristic speed over long distances.
- Although a wave travels, the actual molecules and atoms of the material do not move, instead they vibrate about their equilibrium positions at a different speed.
- Molecules and atoms in the material do not travel in the same direction as the wave.
- Here, the motion of the rope is vertical, but the wave pulses made by the man are moving in the tangential direction.
- Such waves are called **transverse waves**.



Example of transverse waves

Physics for Life Scientists (PHYS-183), Paul D. Nation



- If we replace the rope with a spring, then the oscillations in the spring are in the same direction as the wave moves

- These kind of waves are called **longitudinal waves**.


- Some waves are transverse, some are longitudinal, and some are a combination of both.

- Waves on the ocean are both transverse and longitudinal, but sound waves in air are longitudinal only.

Longitudinal Wave



Example of longitudinal waves.



Simulation of sound waves from a violin string



Physics for Life Scientists (PHYS-183), Paul D. Nation

- Waves that can be described by sinusoidal functions are called **harmonic waves**.


- There are two general kinds of waves:

Standing waves are waves that have a constant position

Traveling waves are waves that move in a specific direction as a function of time.


Standing Waves:

- One way to generate standing waves is to drive a harmonic oscillator on resonance



A = wave amplitude

λ = wavelength



- Instead of wavelength λ we can also define the **wave number**:

$$k = 2\pi/\lambda$$



Physics for Life Scientists (PHYS-183), Paul D. Nation

- If we take a picture of the string, then the displacement as a function of position x is:

$$y(x) = A \sin(kx)$$

- As we move along the x -direction, every time $kx = \frac{2\pi x}{\lambda}$ increases by 2π then the amplitude is the same

Traveling Waves:

- Our standing wave on the string is actually a specific example of a traveling wave.
- Standing waves are actually waves that move so that their shape does not change.
- We can describe a general traveling wave by using an expression where the amplitude changes as both a function of space and time.

$$y(x, t) = A \sin(kx - \omega t) \quad (\omega = 2\pi f)$$

- Using a constant time (ex. $t=0$) then this equation is the same as that of the standing wave amplitude.
- Different values for t (ex. $t=1, 2, \dots$) just shift the argument of the sine function.



- If we fix the position x instead of time, then we see that the amplitude oscillates up and down with amplitude A and frequency ω

$$y(0, t) = A \sin(-\omega t)$$

- Each element of the string moves up and down only.


- Since the wave is traveling, it must have a velocity. This velocity can be found using the waves parameters ω and k

$$v = \frac{x}{t} = \lambda f = \frac{2\pi f}{2\pi/\lambda} = \frac{\omega}{k}$$

- The entire wave $y(x, t)$ moves along the $+x$ -direction with velocity $v = \omega/k$

$$y(x, t) = A \sin(kx - \omega t) \quad \leftarrow \text{moves in } +x\text{-direction}$$

$$y(x, t) = A \sin(kx + \omega t) \quad \leftarrow \text{moves in } -x\text{-direction}$$



Ex. A traveling wave on a string is described by the equation:

$$y(x, t) = 0.025 \sin(1.5x - 200t)$$

what is the waves amplitude, wavelength, frequency, period, and velocity if x and y are measured in meters, and t in sec?

Solution:

- The amplitude of the wave can be obtained directly: $A = 0.025$ m
- To get the wavelength we must convert from wave number: $\lambda = 2\pi/k = 2\pi/(1.5) = 4.2$ m
- Frequency must come from angular frequency: $f = \omega/2\pi = 200/2\pi = 31.8$ Hz
- The period is just the inverse of the frequency: $T = 1/f = 1/31.8 = 0.03$ s
- The velocity can be calculated from ω and k : $v = \omega/k = 200/1.5 = 133$ m/s



Physics for Life Scientists (PHYS-183), Paul D. Nation

What is moving in a wave?:

- We have seen that each point in a wave does not move in the same direction as the wave, but instead moves harmonically in the tangential direction

$$y(x, t) = A \sin(kx - \omega t)$$

- **Key idea:** Each point in the wave moves exactly like a harmonic oscillator.

- Therefore, the question is what exactly is moving when a wave moves?

Waves transfer energy from one place to another.

- For example, given a external force acting on one end of a string, a wave transfers the energy from this force to the other end of the string at the velocity $v = \omega/k$

- As a wave moves down a string, the amplitude at each location changes, causing a change in the kinetic energy of the harmonic oscillator at each point.

Maximum kinetic energy when $y = 0$

No kinetic energy when $y = \pm A$



Physics for Life Scientists (PHYS-183), Paul D. Nation