

Mortality age

A reproducible, sex-dependent, multi-cohort mortality-equivalent age experiment

Version 0.1 · 9 October 2026 · Longevity World Cup

Finding: NHANES can support several jointly measured mortality models, but these public cohorts do not establish a validated clock using all eight requested measurement domains. No respondent has the complete panel. The fitness panel failed a sparse five-year temporal calibration test. The complete-panel equation is published as an unvalidated integration experiment, with its assumptions exposed.

The development integration includes 12,645 people aged 18-49 and 475 deaths over available follow-up. Marginal temporal evaluation includes 8,182 people, 183 total deaths, and 78 deaths within five years. Missing measurements are integrated out during this evaluation; this is **not validation of a jointly observed eight-domain prediction**.

In marginal temporal evaluation, the full experiment has observed/expected five-year mortality 1.10 and unweighted Harrell C 0.691. The observed inflammation/kidney panel has observed/expected 1.01, slope 1.04 and C 0.852. Its broader age range prevents interpreting this as a head-to-head comparison.

The fitness panel has only 4 five-year deaths in its temporal test, with observed/expected 0.19. Its numeric panel age is withheld in the calculator. Other panel estimates and any full integration result remain experimental, not clinical validation.

The displayed age matches modeled five-year all-cause mortality to a historical sex-specific age reference. It measures an association in selected survey populations. It does not estimate pace of aging, life expectancy, treatment benefit, or years gained.

Calculator: longevityworldcup.com/mortality-age

Frozen model: [mortality-age-model.json](#)

1. Measurements and populations

Domain	Definition and canonical unit
Circulation	Mean nonzero readings 1-3 of resting clinic systolic and diastolic BP, mmHg. Both enter the joint equation. Not 24-hour blood pressure. [4]
Fitness	CVDESVO2, exercise-based treadmill estimate, mL/kg/min. Do not substitute CVDVOMAX, the non-exercise protocol-assignment estimate. Eligibility excluded many people with health conditions; ages 12-49 were tested. Adult calculator range: 18-49. [5-7]
Strength	Highest MGX trial from either hand with effort code 1 (maximal). Unit kg. Not MGDCGSZ, which adds best readings from both hands. [8-9]
Body composition	Measured waist cm / standing height cm. The same unit is required in numerator and denominator. [10]
Lipids	LBXAPB ApoB, mg/dL, from the morning fasting subsample. [11-12]
Metabolism	LBXGH HbA1c, NGSP percent. IFCC conversion: % = $0.09148 \times \text{mmol/mol} + 2.152$. [13]
Kidney	SSCYST cystatin C, mg/L, revised SSCARD_A stored-sample release spanning 1999-2004. Laboratory testing was in 2018-2020. Dedicated updated sample weights are used for its distributions and observed panels. [3]
Inflammation	CRP: early LBXCRP mg/dL $\times 10$; 2015 LBXHSCRP is already mg/L. Released below-detection fill values are retained. The 2015 assay is separate sensitivity data. [14-15]

Recorded RIAGENDR categories (female/male) are required. Every component includes a regularized sex-dependent slope; sex is also included in the baseline and age reference. These categories do not encode gender identity or hormone treatment. Differences that shrink toward zero are uncertain, not proof that sexes have identical effects.

Analytic rows are mortality-eligible, nonpregnant where identified, aged 18-79, with event status 0/1 and positive examination follow-up. Ages 80+ are excluded because of top-coding. Unknown pregnancy is not equivalent to a measured negative pregnancy test. The common core requires BP, waist/height and HbA1c. Complete-core selection and measurement exclusions may affect transport.

2. Data, overlap and frozen split

Public NHANES 1999-2016 is linked by SEQN to the 2019 public-use mortality files, using PERMTH_EXM and MORTSTAT. All causes are included. Public follow-up can be perturbed for confidentiality; mortality status is preserved. Restricted newer linkage was not accessed. [1-2]

Development cycles: 1999, 2001, 2005, 2007, 2011. Temporal evaluation: 2003, 2009, 2013. The 2015 cycle is a separate two-year assay/short-follow-up sensitivity. The plan was saved after availability counts and before fitting or prediction performance. A deterministic respondent hash separates internal 80% development fitting from 20% model selection.

Review corrected combined-cycle weight scaling and latent residual subsample weights after the initial temporal results were inspected. All fits and diagnostics were regenerated with unchanged predefined choices. This is re-evaluation on the same temporal split, not a new untouched validation sample.

Cycle	Core N / D	CRP+cys N / D	ApoB+CRP N / D	ApoB+grip N / D	CRP+cys+VO2 N / D
1999-00	3,944 / 976	3,502 / 878	0 / 0	0 / 0	1,137 / 49
2001-02	4,257 / 810	3,965 / 760	0 / 0	0 / 0	1,379 / 42
2003-04	4,006 / 726	3,762 / 682	0 / 0	0 / 0	1,325 / 25
2005-06	4,068 / 523	0 / 0	1,935 / 246	0 / 0	0 / 0
2007-08	4,848 / 606	0 / 0	2,346 / 308	0 / 0	0 / 0
2009-10	5,191 / 429	0 / 0	2,503 / 202	0 / 0	0 / 0
2011-12	4,578 / 302	0 / 0	0 / 0	2,045 / 143	0 / 0
2013-14	4,955 / 216	0 / 0	0 / 0	2,234 / 80	0 / 0
2015-16	4,716 / 130	0 / 0	2,085 / 56	0 / 0	0 / 0

N = observed complete rows for that panel; D = deaths over all available follow-up, not five years. Panels include the common core. These availability counts precede positive-weight exclusions. **All-eight complete cases: zero in every cycle.** Exercise fitness and grip were never collected together in these cycles.

Model	Development N / D	Temporal N / D / D≤5y
Common core	21,695 / 3217	14,152 / 1371 / 495
Core + CRP + cystatin	7,467 / 1638	3,762 / 682 / 156
Core + ApoB + CRP	4,281 / 554	2,503 / 202 / 79
Core + ApoB + grip	2,045 / 143	2,234 / 80 / 64
Core + CRP + cystatin + fitness	2,516 / 91	1,325 / 25 / 4

The age/sex reference uses the complete-core development population aged 18-79, not every eligible NHANES respondent. Modelled five-year mortality is therefore a historical complete-core reference, not a forecast of current national mortality.

3. Survival model and sex-dependent components

Follow-up t is in years; d is death status. Sex s is 0 for female and 1 for male. For observed inputs, the sex-specific Gompertz model is:

$$\begin{aligned}\eta(x, a, s) &= \alpha + \delta s + b_{\text{age}, s}(a - 45)/10 + \sum_j f_j(x_j, s) \\ h(t|x, a, s) &= \exp(\eta) \exp(\gamma_s t) \\ H(t|x, a, s) &= \exp(\eta) A(\gamma_s t), \text{ where } A(\gamma, t) = (\exp(\gamma t) - 1)/\gamma \\ P(\text{death by 5 years}) &= 1 - \exp[-H(5)]\end{aligned}$$

Log transformations are used for ApoB, CRP, cystatin C, grip and fitness to model positive measurements. Other measurements use their raw scale. Development-weighted 10th/50th/90th percentiles are the three restricted-cubic-spline knots. z is centered/scaled transformed input; q is centered/scaled spline basis. The component is:

$$f_j(x, s) = \beta_j z_j(x) + \theta_j q_j(x) + \psi_j z_j(x)(s - 0.5)$$

This is $f(\text{VO2max}, \text{sex})$, $f(\text{grip}, \text{sex})$, and equivalent sex-dependent functions for the other markers. Shared nonlinear curvature plus sex-specific slope deviations borrow information. Separate unrestricted male/female nonlinear fits would consume more information than the young-event counts support. No biomarker is capped to a preferred clinical value, and U/J shapes are not forced.

Three knots $k_1 < k_2 < k_3$ define q before centering/scaling: $[(u-k_1)_+^3 - (u-k_2)_+^3(k_3-k_1)/(k_3-k_2) + (u-k_3)_+^3(k_2-k_1)/(k_3-k_2)]/(k_3-k_1)^2$. Its tails are linear mathematically; the calculator refuses values beyond declared empirical support instead of extrapolating those tails.

The weighted negative mean log-likelihood uses ridge shrinkage. Relative penalties are 1 for main linear terms, 4 for nonlinear terms, 8 for sex deviations, 0.05 for sex, 0.02 for age slopes and zero for intercept. The internal grid is linear/log $\lambda = 0.0003, 0.001, 0.003$ and smooth $\lambda = 0.001, 0.003$. The smooth $\lambda = 0.003$ candidate wins by internal marginal survival loss and is refit on development only.

Candidate	Internal loss	Converged
Linear/log, $\lambda=0.0003$	0.217703	True
Linear/log, $\lambda=0.001$	0.216993	True
Linear/log, $\lambda=0.003$	0.216585	True
Smooth, $\lambda=0.001$	0.215718	True
Smooth, $\lambda=0.003$	0.215672	True

Numerical age slopes are constrained positive for reference inversion; follow-up slopes are bounded -0.1 to 0.2 per year. These are optimizer constraints, not biomarker clipping. Survey weights are normalized by sum in the likelihood. The fit includes full right-censored follow-up; its five-year output depends on the time form and proportional-effects assumption.

4. What combining cohorts assumes

Adding adjusted marginal hazard ratios would double count shared information and would not recover a joint model. Here each original participant contributes a survival likelihood marginalized over their unobserved measurements:

$$L_i = E_{X_{\text{missing}} | X_{\text{observed}}, \text{age}, \text{sex}, \text{core}} \{ h(t_i | X)^{d_i} \exp[-H(t_i | X)] \}$$

Conditional measurement means are ridge regressions on age, age², sex, age×sex, the four common-core measurements and their sex interactions. Residuals for the five occasionally measured inputs are approximated by sex-specific multivariate Gaussian distributions on standardized transformed scales. Observed values remain fixed during integration. The draws are quadrature, not fabricated observed participants.

Genuinely observed residual pair covariance is retained. A pair with fewer than 50 jointly measured records in a sex is assigned zero residual covariance in the primary model. Unobserved relationships cannot be inferred merely because datasets share the common core. The zero-residual-dependence choice and the transport of conditional distributions/outcome relationships across cycles and fitness exclusions are assumptions. [16-17]

Pair	Female N	Male N
apob + crp	1147	1280
apob + cystatin	0	0
apob + grip	547	632
apob + vo2	0	0
crp + cystatin	2158	2309
crp + grip	0	0
crp + vo2	1293	1437
cystatin + grip	0	0
cystatin + vo2	1193	1323
grip + vo2	0	0

Conditional means and residual covariances use dedicated ApoB fasting weights or revised cystatin weights for the involved subsample markers, otherwise MEC weights. Observed-panel survival uses the panel-specific weights. Per CDC guidance, early four-year weight numerators are twice the two-year numerators; a common normalization cancels. The full outcome model uses MEC weights; no published joint subsample weight covers all eight domains. Fitness noncompletion lacks an official fitness response weight. These limitations affect population calibration. [3,18,21]

The primary integral uses 32 deterministic antithetic Gaussian draws; 64-draw refitting checks numerical sensitivity. Residual covariance matrices are shrunk toward their diagonal only if needed for positive definiteness. Unknown pair correlations are changed to -0.3, +0.3 and +0.6, then refit. PSD shrinkage changes the attained correlations; all matrices and shrinkage are saved.

5. Temporal evaluation and its limits

Model	O/E (80% PSU range)	C	Slope	Brier 5y
Full integration (marginal)	1.10 (0.96 to 1.24)	0.691	0.74	0.00932
Common core	0.99 (0.94 to 1.05)	0.814	0.79	0.02640
Core + CRP + cystatin	1.01 (0.91 to 1.10)	0.852	1.04	0.02376
Core + ApoB + CRP	0.87 (0.76 to 0.96)	0.832	0.81	0.02371
Core + ApoB + grip	0.96 (0.82 to 1.09)	0.794	0.72	0.02284
Core + CRP + cystatin + fitness	0.19 (0.09 to 0.31)	0.612	Not estimable	0.00124

O/E = weighted Kaplan-Meier observed five-year death probability / weighted mean predicted probability; 1 is the target. Its 80% interval uses 128 survey-PSU bootstraps within strata with the model fixed. C is unweighted Harrell concordance over available censored follow-up, not five-year AUC. Slopes use a five-year truncated Cox recalibration on log cumulative risk when at least 30 five-year events are available. A missing slope is insufficient event information. Brier error uses inverse-censoring-probability weights with the censoring distribution estimated in the evaluation cohort. C/slope/Brier point estimates do not have design intervals in this initial report.


Same under-50 holdout	O/E	C	Brier 5y
Age/sex	1.45	0.649	0.009325
Observed common core	1.45	0.675	0.009311
Full marginal integration	1.10	0.691	0.009320

The full marginal Brier improvement over age/sex is 0.000005, small in absolute terms. These predictions integrate missing measurements. Even good marginal calibration would not prove that the unseen joint eight-domain equation is correct. There is no external cohort validation.

Full marginal test	N	Deaths ≤5y	O/E (80% PSU range)
Female	4102	26	0.90 (0.75 to 1.08)
Male	4080	52	1.21 (1.02 to 1.40)


The fitness panel substantially overpredicts the sparse test outcomes. With four five-year deaths it cannot supply reliable calibration or elaborate sex-specific fitness curves. The release decision to withhold that panel is a post-evaluation usability decision; the frozen model and all failure metrics remain available. The full model is not promoted to a validated clock by partial-panel results.

6.1 Sex-dependent curves and conditional uncertainty



Each curve is the log-hazard contribution relative to that sex's development median profile, holding other inputs fixed. Shading is the 10th-90th percentile of 32 survey-cluster refits with fixed knots, conditional distributions, selected specification and age reference. It omits nuisance-estimation and unobserved-dependence uncertainty. Limits are development-weighted 1st-99th input percentiles by sex, not clinical normal ranges.


6.2 Sex-dependent curves and conditional uncertainty



Each curve is the log-hazard contribution relative to that sex's development median profile, holding other inputs fixed. Shading is the 10th-90th percentile of 32 survey-cluster refits with fixed knots, conditional distributions, selected specification and age reference. It omits nuisance-estimation and unobserved-dependence uncertainty. Limits are development-weighted 1st-99th input percentiles by sex, not clinical normal ranges.

ApoB has an inverse adjusted association in this experiment. This is not evidence that raising ApoB improves health. Selection, illness, confounding and the integration assumptions can affect this association.

6.3 Sex-dependent curves and conditional uncertainty



Each curve is the log-hazard contribution relative to that sex's development median profile, holding other inputs fixed. Shading is the 10th-90th percentile of 32 survey-cluster refits with fixed knots, conditional distributions, selected specification and age reference. It omits nuisance-estimation and unobserved-dependence uncertainty. Limits are development-weighted 1st-99th input percentiles by sex, not clinical normal ranges.

The fitness contribution is weakly protective over the displayed support in this corrected fit. Sparse outcomes and selective exercise eligibility make a biological interpretation unjustified. Shrunk sex deviations do not establish equivalence between sexes.

7. Sensitivity, uncertainty and failed hypotheses

Refit / assumption	Female example age	Male example age
Primary 32-draw integration	40.20	37.41
Unknown residual correlation -0.3	40.08	37.46
Unknown residual correlation +0.3	39.91	37.16
Unknown residual correlation +0.6	39.69	37.07
64-draw integration	40.12	37.35
Exponential baseline + matched reference	36.87	34.82

Examples have chronological age 40 and development medians for each sex. They are artificial profiles, not two jointly observed people. The exponential sensitivity refits both outcome model and age reference with a constant baseline hazard. It demonstrates time-form dependence; it does not establish the proportional-effects assumption.

Exploratory sex-stratified Cox diagnostics using the observed core/blood spline bases found ranked-time associations in scaled Schoenfeld residuals. The smallest approximate p values are core: age_male, $p=4e-23$; blood: age_male, $p=8.6e-16$. This raises concern about constant covariate effects over follow-up. These penalized, survey-weighted diagnostics have no survey-design or multiple-testing correction and are not a formal test of the integrated likelihood.

One-at-a-time sweeps across sex, ages 20/30/40/49 and nine values per input produced 648 artificial profiles. The median tested dependence age span is 0.72 years; the maximum is 3.54 years. 0 primary profiles lie outside the reference-age support. This is not a simultaneous joint-tail stress test and does not exhaust unidentified relationships.

Observed-panel two-year landmark fits restart follow-up among people surviving past two years, subtract 24 months and advance age by two years. Baseline measurements remain two years old. This handles the time origin correctly but does not remove survivor selection or reverse causality. The separate integrated landmark artifact is exploratory and also changes conditional missing-input assumptions; it is not treated as a validation result.

The 32 training bootstraps are conditional on fixed nuisance distributions, knots, model selection and reference; their 80% age/curve ranges are not comprehensive prediction intervals. Structural sensitivity spans are separate, finite scenario comparisons, not confidence intervals. Regularization may conceal weakly identified components. One must not infer that each domain is independently measured biological aging.

The proposed all-domain clock hypothesis remains unestablished. Additional joint measurements, mortality events, sex-specific evaluation, measurement-method calibration and independent cohort validation are required before clinical or intervention interpretation. Accessible public NHANES overlaps cannot supply the absent exercise/grip joint information. HRS health access and UK Biobank would require separately authorized access; neither dataset was used. [19-20]

7.1 Later assays and longer follow-up

Two-year later-assay and ten-year temporal diagnostics were specified in transport-plan.md after primary results were inspected and before these endpoint results were calculated. Horizons were selected from administrative follow-up coverage. All models, measurement distributions and release decisions remain frozen. These are diagnostic sensitivities, not new confirmatory validation.

2015-2016: newer hs-CRP assay, two-year endpoint

Model	N	D≤2y	O/E (80% PSU range)	Slope	Brier 2y
Full integration (marginal)	2,604	8	1.12 (0.67 to 1.53)	Not estimable	0.003621
Common core	4,716	68	1.04 (0.83 to 1.26)	0.83	0.011383
Core + ApoB + CRP	2,085	31	0.88 (0.66 to 1.07)	0.83	0.009968

2003-2004: ten-year endpoint

Model	N	D≤10y	O/E (80% PSU range)	Slope	Brier 10y
Full integration (marginal)	2,381	44	0.78 (0.66 to 0.91)	0.71	0.015713
Common core	4,006	373	1.00 (0.94 to 1.06)	0.98	0.052571
Core + CRP + cystatin	3,762	349	1.00 (0.94 to 1.06)	1.02	0.049485
Core + CRP + cystatin + fitness	1,325	9	0.34 (0.24 to 0.44)	Not estimable	0.005043

The later cycle has at least 2.83 years of observed administrative follow-up, so it supports the two-year check and is not presented as five-year validation. Its population, calendar period and selection also differ; the comparison cannot isolate the assay effect. The 2003 cycle has at least 14.75 years of administrative follow-up. Its ten-year results reuse an already inspected temporal cohort.

Same-row full marginal comparison	Age/sex Brier	Core Brier	Full Brier
2015, 2 years	0.003620	0.003618	0.003621
2003, 10 years	0.015317	0.015388	0.015713

The full marginal Brier error is not lower than either benchmark in these additional checks. Only eight two-year deaths occur in the later under-50 cohort; the ten-year full marginal O/E is below one. These findings limit prediction reliability and do not establish unseen joint-panel validity. Fitness remains numerically withheld.

Full marginal sex coverage	Female N / D≤h	Male N / D≤h
2015, 2 years	1,328 / 2	1,276 / 6
2003, 10 years	1,132 / 15	1,249 / 29

Ranges use 128 conditional PSU bootstraps with fixed models. Sparse sex strata cannot establish calibration. Exact sex-specific calibration, concordance over available follow-up, follow-up counts and same-row benchmarks are in the saved artifacts. The new evaluator agrees with the original five-year evaluator and an analytical censoring example. Diagnostic plan SHA-256: 003551b5ad9b28f57538f7409698ae07521ceb4a2adda4bffd70c424ed3e110

8. From risk to age, and supported inputs

Reference risk is fit in the complete-core development population aged 18-79. The same five-year endpoint and sex are used on both sides. Matching cumulative hazard rather than rounded probabilities gives a stable closed-form age:

$$\text{Age}_{\text{eq}} = 45 + 10 \{ \log H_{\text{model}}(5) - \log A(y_{\text{ref},s}, 5) - \alpha_{\text{ref}} - \delta_{\text{ref},s} \} / b_{\text{ref},\text{age},s}$$

Reference parameter	Value
intercept	-6.75843283148
male	0.85644335816
age_female	0.920587062491
age_male	0.832440010619
gamma_0	0.120357788382
gamma_1	0.0921579396873

The reference is monotone because its age coefficients are positive. Inversion is allowed only over ages 18-79. The full fitness integration accepts chronological ages 18-49; nonfitness panels accept 18-79. Inputs outside sex-specific weighted 1st-99th development percentiles return unsupported, not a capped measurement or clipped age. These marginal ranges do not certify support for every joint combination.

The calculator withholds a sampling or dependence range if any included refit exceeds reference-age support, rather than silently dropping its tail.

Input	Female support	Male support	Unit
Systolic BP	88.000 to 153.333	95.000 to 156.248	mmHg
Diastolic BP	47.341 to 95.000	42.667 to 101.333	mmHg
Waist / height	0.402 to 0.847	0.397 to 0.813	ratio
HbA1c	4.400 to 8.600	4.400 to 9.400	%
ApoB	41.000 to 155.000	44.000 to 159.000	mg/dL
hs-CRP	0.100 to 32.713	0.100 to 25.700	mg/L
Cystatin C	0.470 to 1.080	0.542 to 1.148	mg/L
Single-hand grip	20.440 to 44.800	30.289 to 70.049	kg
Estimated VO2max	21.235 to 63.534	29.544 to 80.007	mL/kg/min

The calculator accepts an exercise-based estimate; wearable/non-exercise fitness methods return unsupported for the fitness/full calculation because cross-method calibration was not established. If a laboratory reports a value below a known reporting limit, the entered limit is divided by $\sqrt{2}$ for CRP/cystatin, matching the public-data fill convention. Unknown limits are not assigned invented values. Inputs and drafts are local to the browser; no health values are sent to a server.

9. Frozen equation coefficients

Component	Linear β	Spline θ	Sex slope ψ
Systolic BP	0.0731042696	0.140414199	0.0326650307
Diastolic BP	-0.0768089709	0.0731256378	-0.0149549228
Waist / height	-0.11159028	0.0912758211	-0.0888911961
HbA1c	0.168682458	0.0306600532	0.0161358542
ApoB	-0.136205129	-0.0700707488	-0.0464397107
hs-CRP	0.245665942	0.0281243438	0.00821889422
Cystatin C	0.136169588	0.0969965178	-0.0346365643
Single-hand grip	-0.0862748358	-0.0150869237	0.00541760257
Estimated VO2max	-0.041393371	0.0179198716	-0.0116652611

The component sex slope multiplies $z(x)(s-0.5)$, not raw x . Coefficients require the exact transformations, knots, centers/scales and baseline. A component taken from another panel cannot be swapped into this full equation while retaining its interpretation: each panel is fitted jointly and calibrated against the reference.

Full-model baseline	Value
intercept	-6.70775957728
male	0.70722925402
age_female	0.594817385033
age_male	0.480861817632
gamma_0	0.0853512354113
gamma_1	0.0669631258632

Input	Transformed knots	Center / scale
Systolic BP	101.333 / 115.333 / 132.667	116.484 / 13.1484
Diastolic BP	58 / 71.3333 / 85	71.3022 / 11.3087
Waist / height	0.445946 / 0.542036 / 0.679291	0.554361 / 0.0936493
HbA1c	4.8 / 5.2 / 5.7	5.3129 / 0.765764
ApoB (log)	4.12713 / 4.48864 / 4.81218	4.47225 / 0.278599
hs-CRP (log)	-1.20397 / 0.405465 / 2.11626	0.407454 / 1.31926
Cystatin C (log)	-0.525939 / -0.328504 / -0.12897	-0.325157 / 0.182517
Single-hand grip (log)	3.31054 / 3.68872 / 4.04263	3.67641 / 0.285085
Estimated VO2max (log)	3.39047 / 3.67704 / 3.96556	3.67681 / 0.235852

All digits, nonlinear centers/scales, panel coefficients, conditional covariance matrices, failed specifications and numerical fixtures are saved in the repository artifacts. The downloadable browser JSON contains the exact evaluated equation. The report's rounded tables are descriptive; calculations use the full stored precision.

10. Reproduction and provenance

Source: LongevityWorldCup.Research/MortalityAge. Run `prepare.py`, `train.py`, `diagnostics.py`, `ph-diagnostics.py`, `transport.py`, `export.py`, `report.py` in that order in the pinned environment. Run `test_model.py` and `verify-browser.mjs` for independent numerical and unit parity. Raw downloads, respondent data and intermediate outputs remain in ignored `.artifacts/mortality-age`.

Model version: **mortality-age-0.1-19e880e4f33f**

Frozen plan SHA-256: `fae4b2b157c8a1f222aff3d17b2487ff2d1a6e86f3a8676655439f263b6e89e6`

Browser model SHA-256: `f04d41939c4ea3f868a6cbc20a47580433ff9984fab2cd5917992dc953d08933`

64 original public files are recorded with URL, retrieval UTC, byte size, SHA-256, row count and columns in `artifacts/downloads.json`. Joins are one-to-one on SEQN with duplicate checks. SAS file signatures and mortality-file content are verified. Primary fit completion UTC: `2026-10-09T14:51:33.306171+00:00`. Additional diagnostic completion UTC: `2026-10-09T15:24:26.420998+00:00`. Monte Carlo seed: 91237.

Runtime / package	Version
Python	3.14.7
numpy	2.4.3
scipy	1.17.1
pandas	2.3.3
lifelines	0.30.3
matplotlib	3.11.2
reportlab	4.4.10

Verification includes analytic likelihood gradients against independent finite differences, exponential-limit calculations, observed-row/quadrature equivalence, right-censored risk sets, sex-specific age roundtrips, spline tails and Python/browser fixtures. Website checks cover unit conversions, sex-dependent functions, unsupported inputs, loading recovery, editing and restoration, mobile/desktop behavior, PDF bytes and exclusion from discovery. Code verification does not constitute scientific validation.

This report distinguishes actual temporal performance from scientific identification. The model has no complete-panel external validation, no causal adjustment or competing-risk interpretation, and no calibrated smartwatch conversion. Medication, smoking, disease, ethnicity and socioeconomic conditions are not separately included in the requested panel. Repeated measurements and longitudinal biological change were not modelled. Those limitations are material, not resolved by producing an age number.

11. Primary sources

- [1] CDC/NCHS public-use linked mortality description
<https://www.cdc.gov/nchs/data/datalinkage/public-use-linked-mortality-file-description.pdf>
- [2] CDC/NCHS mortality data dictionary
<https://www.cdc.gov/nchs/data/datalinkage/public-use-linked-mortality-files-data-dictionary.pdf>
- [3] NHANES SSCARD_A: revised cystatin/stored samples
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/1999/DataFiles/SSCARD_A.htm
- [4] NHANES BPX_G blood pressure
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2011/DataFiles/BPX_G.htm
- [5] NHANES CVX exercise fitness 1999-2000
<https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/1999/DataFiles/CVX.htm>
- [6] NHANES CVX_B exercise fitness 2001-2002
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2001/DataFiles/CVX_B.htm
- [7] NHANES CVX_C exercise fitness 2003-2004
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2003/DataFiles/CVX_C.htm
- [8] NHANES MGX_G grip
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2011/DataFiles/MGX_G.htm
- [9] NHANES MGX_H grip
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2013/DataFiles/MGX_H.htm
- [10] NHANES BMX_G anthropometry
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2011/DataFiles/BMX_G.htm
- [11] NHANES TRIGLY_D ApoB and morning weights
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2005/DataFiles/TRIGLY_D.htm
- [12] NHANES APOB_G
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2011/DataFiles/APOB_G.htm
- [13] NGSP/IFCC master equation
<https://ngsp.org/ifcc.asp>
- [14] NHANES CRP_F
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2009/DataFiles/CRP_F.htm
- [15] NHANES HSCRP_I
https://wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2015/DataFiles/HSCRP_I.htm
- [16] Raessler: Data Fusion, Identification Problems, Validity and Multiple Imputation
<https://www.ajs.or.at/index.php/ajs/article/view/vol33,%20no1&2%20-%209>
- [17] White and Royston: missing covariates in Cox models
<https://pmc.ncbi.nlm.nih.gov/articles/PMC2998703/>
- [18] NHANES analytic guidelines
<https://wwwn.cdc.gov/Nchs/Nhanes/analyticguidelines.aspx>
- [19] HRS sensitive health data access
<https://hrsdata.isr.umich.edu/data-products/sensitive-health>
- [20] UK Biobank access procedures
<https://www.ukbiobank.ac.uk/access-procedures/>
- [21] CDC NHANES combined-cycle and subsample weighting tutorial
<https://wwwn.cdc.gov/nchs/nhanes/tutorials/weighting.aspx>