

Totally nonnegative matrices, chain enumeration and zeros of polynomials


Petter Brändén

KTH Royal Institute of Technology

joint work with
Leonardo Saud Maia Leite

Einar fest, Reykjavik, June 18, 2025

Til hamingju Einar!





"The one contribution of mine that I hope will be remembered has consisted in pointing out that all sorts of problems of combinatorics can be viewed as problems of the location of the zeros of certain polynomials..."



Gian-Carlo Rota


Chain enumeration in posets

- ▶ A central theme in enumerative and algebraic combinatorics is counting chains in posets.



Chain enumeration in posets

- ▶ A central theme in enumerative and algebraic combinatorics is counting chains in posets.




- ▶ A **chain** in P is a totally ordered subset of P , i.e.,

$$x, y \in P \implies x \leq y \text{ or } y \leq x.$$

- ▶ For example $\{1, 6, 7\}$.

Chain enumeration in posets

- ▶ A central theme in enumerative and algebraic combinatorics is counting chains in posets.



- ▶ A chain in P is a totally ordered subset of P , i.e.,

$$x, y \in P \implies x \leq y \text{ or } y \leq x.$$


- ▶ For example $\{1, 6, 7\}$.

- ▶ The chain polynomial of P is

$$c_P(t) = \sum_{C \text{ chain}} t^{|C|}.$$

Chain enumeration in posets

- ▶ A central theme in enumerative and algebraic combinatorics is counting chains in posets.



- ▶ A chain in P is a totally ordered subset of P , i.e.,

$$x, y \in P \implies x \leq y \text{ or } y \leq x.$$

- ▶ For example $\{1, 6, 7\}$.

- ▶ The chain polynomial of P is

$$c_P(t) = \sum_{C \text{ chain}} t^{|C|}.$$

- ▶ Example. If $P = \{1 < 2 < \dots < n\}$, then $c_P(t) = (1 + t)^n$.

Chain enumeration in posets

- ▶ Example. If P is the Boolean algebra \mathbb{B}_n , then

$$c_P(t) = (1+t)^2 \sum_{k=0}^{n-1} (k+1)! \cdot S(n, k+1) t^k.$$

- ▶ Frobenius proved that $c_{\mathbb{B}_n}(t)$ is real-rooted.

Chain enumeration in posets


- ▶ Example. If P is the Boolean algebra \mathbb{B}_n , then

$$c_P(t) = (1+t)^2 \sum_{k=0}^{n-1} (k+1)! \cdot S(n, k+1) t^k.$$

- ▶ Frobenius proved that $c_{\mathbb{B}_n}(t)$ is real-rooted.
- ▶ Chain polynomials tend to be real-rooted:
- ▶ The Poset Conjecture (Neggers, 1978): P is a distributive lattice. Counterexamples by B. and Stembridge (2004, -06).
- ▶ $(3+1)$ -free posets (Stanley, 1998).
- ▶ Cohen-Macaulay simplicial complexes (Brenti-Welker, B., 2006).
- ▶ Shellable cubical complexes (Athanasiadis, 2021).
- ▶ Geometric lattices (Conjecture of Athanasiadis and Kalampogia-Evangelinou, 2023).

Rank uniform posets

- ▶ We consider finite, bounded and graded posets P , i.e.,
- ▶ P has a least element $\hat{0}$, and a largest element $\hat{1}$, and
- ▶ P admits a rank function $\rho : P \rightarrow \mathbb{N}$ such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ whenever y covers x .



Rank uniform posets

- ▶ We consider finite, bounded and graded posets P , i.e.,
- ▶ P has a least element $\hat{0}$, and a largest element $\hat{1}$, and
- ▶ P admits a rank function $\rho : P \rightarrow \mathbb{N}$ such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ whenever y covers x .
- ▶ P is rank uniform if whenever $\rho(y) = \rho(z)$, then

$$|\{x \leq y : \rho(x) = k\}| = |\{x \leq z : \rho(x) = k\}|,$$

for all $0 \leq k \leq \rho(y)$.

Rank uniform posets

- ▶ We consider finite, bounded and graded posets P , i.e.,
- ▶ P has a least element $\hat{0}$, and a largest element $\hat{1}$, and
- ▶ P admits a rank function $\rho : P \rightarrow \mathbb{N}$ such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ whenever y covers x .
- ▶ P is rank uniform if whenever $\rho(y) = \rho(z)$, then

$$|\{x \leq y : \rho(x) = k\}| = |\{x \leq z : \rho(x) = k\}|,$$

for all $0 \leq k \leq \rho(y)$.

- ▶ These posets generalize binomial- and Sheffer-posets.

Rank uniform posets

- ▶ We consider finite, bounded and graded posets P , i.e.,
- ▶ P has a least element $\hat{0}$, and a largest element $\hat{1}$, and
- ▶ P admits a rank function $\rho : P \rightarrow \mathbb{N}$ such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ whenever y covers x .
- ▶ P is rank uniform if whenever $\rho(y) = \rho(z)$, then

$$|\{x \leq y : \rho(x) = k\}| = |\{x \leq z : \rho(x) = k\}|,$$

for all $0 \leq k \leq \rho(y)$.

- ▶ These posets generalize binomial- and Sheffer-posets.
- ▶ If P is rank uniform define a matrix $R(P) = (r_{n,k}(P))_{n,k}$, where

$$r_{n,k}(P) = |\{x \leq y : \rho(x) = k\}|,$$

where y is an element of P of rank n .

Matrices from rank uniform posets

- Example. $r_{n,k}(\mathbb{B}_m) = \binom{n}{k}$, where \mathbb{B}_m is a Boolean algebra.

Matrices from rank uniform posets


- ▶ Example. $r_{n,k}(\mathbb{B}_m) = \binom{n}{k}$, where \mathbb{B}_m is a Boolean algebra.
- ▶ Example. $r_{n,k}(\mathbb{B}_m(q)) = \binom{n}{k}_q$, where $\mathbb{B}_m(q)$ is the lattice of all subspaces of an m -dimensional vector space over the finite field \mathbb{F}_q .

Matrices from rank uniform posets

- ▶ Example. $r_{n,k}(\mathbb{B}_m) = \binom{n}{k}$, where \mathbb{B}_m is a Boolean algebra.
- ▶ Example. $r_{n,k}(\mathbb{B}_m(q)) = \binom{n}{k}_q$, where $\mathbb{B}_m(q)$ is the lattice of all subspaces of an m -dimensional vector space over the finite field \mathbb{F}_q .
- ▶ Example. $r_{n,k}(\Pi'_m) = S(n+1, k+1)$, where Π'_m is the dual of the partition lattice.

Matrices from rank uniform posets

- ▶ Example. $r_{n,k}(\mathbb{B}_m) = \binom{n}{k}$, where \mathbb{B}_m is a Boolean algebra.
- ▶ Example. $r_{n,k}(\mathbb{B}_m(q)) = \binom{n}{k}_q$, where $\mathbb{B}_m(q)$ is the lattice of all subspaces of an m -dimensional vector space over the finite field \mathbb{F}_q .
- ▶ Example. $r_{n,k}(\Pi'_m) = S(n+1, k+1)$, where Π'_m is the dual of the partition lattice.
- ▶ Example. $r_{n,k}(Q) = 2^{n-k} \binom{n-1}{k-1}$, where Q is the face lattice of a hypercube.



What do these matrices have in common?



Totally nonnegative matrices

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry, algebraic combinatorics and physics.



Felix Gantmacher



Mark Krein



Samuel Karlin

TN-matrices and TN-posets

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry and algebraic combinatorics.
- ▶ We say that a rank uniform poset P is TN if $R(P)$ is TN.

TN-matrices and TN-posets

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry and algebraic combinatorics.
- ▶ We say that a rank uniform poset P is TN if $R(P)$ is TN.
- ▶ Examples: \mathbb{B}_m , $\mathbb{B}_m(q)$, Π'_m , the face lattice of a hypercube.

TN-matrices and TN-posets

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry and algebraic combinatorics.
- ▶ We say that a rank uniform poset P is TN if $R(P)$ is TN.
- ▶ **Examples:** \mathbb{B}_m , $\mathbb{B}_m(q)$, Π'_m , the face lattice of a hypercube.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.

TN-matrices and TN-posets

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry and algebraic combinatorics.
- ▶ We say that a rank uniform poset P is TN if $R(P)$ is TN.
- ▶ **Examples:** \mathbb{B}_m , $\mathbb{B}_m(q)$, Π'_m , the face lattice of a hypercube.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ Suppose P is a bounded and graded poset of rank r , and that $S \subseteq [r - 1]$. Then

$$P_S = \{x \in P : \rho(x) \in S \cup \{0, r\}\}$$

is the **rank selected subposet** induced by S .

TN-matrices and TN-posets

- ▶ A matrix with real entries is **totally nonnegative (TN)** if all of its minors are nonnegative.
- ▶ Important in linear algebra, statistics, analysis, geometry and algebraic combinatorics.
- ▶ We say that a rank uniform poset P is TN if $R(P)$ is TN.
- ▶ **Examples:** \mathbb{B}_m , $\mathbb{B}_m(q)$, Π'_m , the face lattice of a hypercube.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ Suppose P is a bounded and graded poset of rank r , and that $S \subseteq [r - 1]$. Then

$$P_S = \{x \in P : \rho(x) \in S \cup \{0, r\}\}$$

is the **rank selected subposet** induced by S .

- ▶ If P is TN, then so is P_S .

TN-matrices and TN-posets

- If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ **Theorem** (B., Saud).
 - ▶ A simplicial complex is TN if and only if its h -vector is nonnegative.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ **Theorem** (B., Saud).
 - ▶ A **simplicial complex** is TN if and only if its h -vector is nonnegative.
 - ▶ A **cubical complex** is TN if and only if its cubical h -vector is nonnegative.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ **Theorem** (B., Saud).
 - ▶ A simplicial complex is TN if and only if its h -vector is nonnegative.
 - ▶ A cubical complex is TN if and only if its cubical h -vector is nonnegative.
 - ▶ Shellable q -complexes are TN.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ **Theorem** (B., Saud).
 - ▶ A simplicial complex is TN if and only if its h -vector is nonnegative.
 - ▶ A cubical complex is TN if and only if its cubical h -vector is nonnegative.
 - ▶ Shellable q -complexes are TN.
 - ▶ Dual Dowling lattices are TN.

TN-matrices and TN-posets

- ▶ If P is graded but fails to have a largest element $\hat{1}$, then P is said to be TN if $\hat{P} = P \cup \{\hat{1}\}$ is TN.
- ▶ **Theorem** (B., Saud). The chain polynomial of any TN-poset is real-rooted.
- ▶ **Theorem** (B., Saud).
 - ▶ A simplicial complex is TN if and only if its h -vector is nonnegative.
 - ▶ A cubical complex is TN if and only if its cubical h -vector is nonnegative.
 - ▶ Shellable q -complexes are TN.
 - ▶ Dual Dowling lattices are TN.
- ▶ **Corollary**. Rank selected subposets of the above posets have real-rooted chain polynomials.

The shape of characteristic polynomials

- ▶ Let $\mathcal{H} = \{H_1, \dots, H_m\}$ be a **hyperplane arrangement** (list of hyperplanes) in \mathbb{F}_q^n .

The shape of characteristic polynomials

- ▶ Let $\mathcal{H} = \{H_1, \dots, H_m\}$ be a **hyperplane arrangement** (list of hyperplanes) in \mathbb{F}_q^n .
- ▶ Recall that the **characteristic polynomial** of \mathcal{H} is

$$\chi_{\mathcal{H}}(t) = \sum_{A \subseteq [m]} (-1)^{|A|} t^{\dim(\cap_{i \in A} H_i)}.$$

The shape of characteristic polynomials

- ▶ Let $\mathcal{H} = \{H_1, \dots, H_m\}$ be a **hyperplane arrangement** (list of hyperplanes) in \mathbb{F}_q^n .
- ▶ Recall that the **characteristic polynomial** of \mathcal{H} is

$$\chi_{\mathcal{H}}(t) = \sum_{A \subseteq [m]} (-1)^{|A|} t^{\dim(\cap_{i \in A} H_i)}.$$

- ▶ Can we describe all linear inequalities satisfied by all characteristic polynomials of hyperplane arrangements in \mathbb{F}_q^n ?

The shape of characteristic polynomials

- ▶ Let $\mathcal{H} = \{H_1, \dots, H_m\}$ be a **hyperplane arrangement** (list of hyperplanes) in \mathbb{F}_q^n .
- ▶ Recall that the **characteristic polynomial** of \mathcal{H} is

$$\chi_{\mathcal{H}}(t) = \sum_{A \subseteq [m]} (-1)^{|A|} t^{\dim(\cap_{i \in A} H_i)}.$$

- ▶ Can we describe all linear inequalities satisfied by all characteristic polynomials of hyperplane arrangements in \mathbb{F}_q^n ?
- ▶ **Theorem** (B., Saud). The convex hull of all characteristic polynomials of hyperplane arrangements in \mathbb{F}_q^n is equal to the simplex with vertices

$$t^{n-k}(t-1)(t-q)\cdots(t-q^{k-1}), \quad 0 \leq k \leq n.$$

The shape of chromatic polynomials

- ▶ **Problem.** Describe the convex hull of all chromatic polynomials of graphs on $V = \{1, 2, \dots, n\}$.

Congratulations Einar!

