


Symmetries in the sandpile model and the shuffle conjecture

Mark Dukes


School of Mathematics and Statistics, University College Dublin, Ireland

EinarFest @ NORCOM 2025


1. Chip-firing on $K_{m,n}$ and sandpile dynamics





Toppling Rule:
vertex v_i unstable \Rightarrow
$$\begin{cases} x_i \rightarrow x_i - d_i \\ x_j \rightarrow x_j + e_{ij} \quad \forall j \neq i \end{cases}$$



1. Chip-firing on $K_{m,n}$ and sandpile dynamics




Toppling Rule:
vertex v_i unstable \Rightarrow
$$\begin{cases} x_i \rightarrow x_i - d_i \\ x_j \rightarrow x_j + e_{ij} \quad \forall j \neq i \end{cases}$$





recurrent
configuration

1. Chip-firing on $K_{m,n}$ and sandpile dynamics




Toppling Rule:
vertex v_i unstable \Rightarrow

$$\begin{cases} x_i \rightarrow x_i - d_i \\ x_j \rightarrow x_j + e_{ij} \quad \forall j \neq i \end{cases}$$




recurrent
configuration





non-recurrent
configuration

1. Chip-firing on $K_{m,n}$ and sandpile dynamics




Toppling Rule:
vertex v_i unstable \Rightarrow

$$\begin{cases} x_i \rightarrow x_i - d_i \\ x_j \rightarrow x_j + e_{ij} \quad \forall j \neq i \end{cases}$$




recurrent
configuration




non-recurrent
configuration

$\text{rec}(K_{m,n}) :=$
set of all (weakly-decreasing)
recurrent configurations


2. Characterizing recurrent configurations



2. Characterizing recurrent configurations




2. Characterizing recurrent configurations



Theorem (D and Le Borgne 2013)

$c \in \text{rec}(K_{m,n})$ iff $f(c)$ is a $m \times n$ parallelogram polyomino.

2. Characterizing recurrent configurations



Theorem (D and Le Borgne 2013)

$c \in \text{rec}(K_{m,n})$ iff $f(c)$ is a $m \times n$ parallelogram polyomino.

Corollary

If $c \in \text{rec}(K_{m,n})$ then $\sum c_i = \text{area}(f(c)) + (m + n - 3)$.

3. The parabounce of a polyomino

Fix a canonical toppling on recurrent configurations. Initially add 1 to all vertices (\equiv toppling the sink)

1. Topple all unstable right vertices.
2. Topple all unstable left vertices.
3. If any vertices are unstable then go to 1.

3. The parabounce of a polyomino

Fix a canonical toppling on recurrent configurations. Initially add 1 to all vertices (\equiv toppling the sink)

1. Topple all unstable right vertices.
2. Topple all unstable left vertices.
3. If any vertices are unstable then go to 1.


This canonical toppling can be identified within the corresponding parallelogram polyomino:

3. The parabounce of a polyomino

Fix a canonical toppling on recurrent configurations. Initially add 1 to all vertices (\equiv toppling the sink)

1. Topple all unstable right vertices.
2. Topple all unstable left vertices.
3. If any vertices are unstable then go to 1.

This canonical toppling can be identified within the corresponding parallelogram polyomino:




$$\begin{aligned}\text{parabounce}(P) &= 1 + 1 + 1 + 1 \\ &\quad + 2 + 2 + 2 \\ &= 10 \\ \text{area}(P) &= 8.\end{aligned}$$

3. The parabounce of a polyomino

Fix a canonical toppling on recurrent configurations. Initially add 1 to all vertices (\equiv toppling the sink)

1. Topple all unstable right vertices.
2. Topple all unstable left vertices.
3. If any vertices are unstable then go to 1.

This canonical toppling can be identified within the corresponding parallelogram polyomino:



$$\begin{aligned}
 \text{parabounce}(P) &= 1 + 1 + 1 + 1 \\
 &\quad + 2 + 2 + 2 \\
 &= 10 \\
 \text{area}(P) &= 8.
 \end{aligned}$$

Define

$$\text{Nara}_{m,n}(q, t) := \sum_{P \in \text{Para}_{m,n}} q^{\text{area}(P)} t^{\text{parabounce}(P)}.$$

$\text{Nara}_{m,n}(1, 1)$ gives the Narayana numbers hence our naming ‘ q, t -Narayana polynomials’.

4. Unexpected symmetries

For example, $\text{Nara}_{2,2}(q, t) = (qt)^4(1 + q + t)$,

4. Unexpected symmetries

For example, $\text{Nara}_{2,2}(q, t) = (qt)^4(1 + q + t)$, and

$$\begin{aligned} \text{Nara}_{3,3}(q, t) = (qt)^6 \big(& 1 + t + 2t^2 + t^3 + t^4 \\ & + q + 2qt + 2qt^2 + qt^3 \\ & + 2q^2 + 2q^2t + q^2t^2 \\ & + q^3 + q^3t \\ & + q^4 \big). \end{aligned}$$

4. Unexpected symmetries

For example, $\text{Nara}_{2,2}(q, t) = (qt)^4(1 + q + t)$, and

$$\begin{aligned} \text{Nara}_{3,3}(q, t) = (qt)^6 \big(& 1 + t + 2t^2 + t^3 + t^4 \\ & + q + 2qt + 2qt^2 + qt^3 \\ & + 2q^2 + 2q^2t + q^2t^2 \\ & + q^3 + q^3t \\ & + q^4 \big). \end{aligned}$$

Conjecture (D and Le Borgne 2013)

$\text{Nara}_{m,n}(q, t)$ is symmetric in both m, n and q, t :


$$\text{Nara}_{m,n}(q, t) = \text{Nara}_{m,n}(t, q)$$

$$\text{Nara}_{m,n}(q, t) = \text{Nara}_{n,m}(q, t).$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$$t = (3, 1, 4, 4, 1)$$



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$


$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned} F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\ &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots \end{aligned}$$


5. Parking functions

A sequence (t_1, \dots, t_n) of non-negative integers is an **n -parking function** if there exists a permutation π such that $t_{\pi(i)} \leq i$ for all $1 \leq i \leq n$.

The interpretation is that t_i is the preferred parking spot of car i . If their spot is taken then the car parks in the next available ($> t_i$) spot. If all cars starting with car 1 and ending in car n can park without 'going around' then t is a parking function.



$1 \ 2 \ 3 \ 4 \ 5$
 $t = (3, 1, 4, 4, 1)$
 is a 5-parking function



$$\text{area}(t) = 2$$

$$\text{dinv}(t) = 4$$

$$\text{word}(t) = (4, 5, 3, 1, 2)$$

$$\begin{aligned}
 F_{5, \text{idcs}}(t) &= \sum_{i_1 \leq i_2 < i_3 < i_4 \leq i_5} z_{i_1} z_{i_2} z_{i_3} z_{i_4} z_{i_5} \\
 &= z_1^2 z_2 z_3^2 + z_1^2 z_2 z_3 z_4 + z_1^2 z_2 z_3 z_5 + \dots
 \end{aligned}$$

6. The shuffle conjecture: setting

Diagonal harmonic polynomials are the solutions to a system of PDEs:

$$DH_n := \left\{ f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] : \sum_{i=1}^n \partial_{x_i}^a \partial_{y_i}^b f = 0 \text{ for all } a + b > 0 \right\}.$$

6. The shuffle conjecture: setting

Diagonal harmonic polynomials are the solutions to a system of PDEs:

$$DH_n := \left\{ f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] : \sum_{i=1}^n \partial_{x_i}^a \partial_{y_i}^b f = 0 \text{ for all } a + b > 0 \right\}.$$

E.g.

$$DH_2 = \{c_1 \cdot 1 + c_2 \cdot (x_1 - x_2) + c_3 \cdot (y_1 - y_2) : c_i \in \mathbb{C}\}.$$

6. The shuffle conjecture: setting

Diagonal harmonic polynomials are the solutions to a system of PDEs:

$$DH_n := \left\{ f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] : \sum_{i=1}^n \partial_{x_i}^a \partial_{y_i}^b f = 0 \text{ for all } a + b > 0 \right\}.$$

E.g.

$$DH_2 = \{c_1 \cdot 1 + c_2 \cdot (x_1 - x_2) + c_3 \cdot (y_1 - y_2) : c_i \in \mathbb{C}\}.$$

Set $DH_n^{c,d}$ to be those $f \in DH_n$ whose total x degree is c and whose total y degree is d .

6. The shuffle conjecture: setting

Diagonal harmonic polynomials are the solutions to a system of PDEs:

$$DH_n := \left\{ f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] : \sum_{i=1}^n \partial_{x_i}^a \partial_{y_i}^b f = 0 \text{ for all } a + b > 0 \right\}.$$

E.g.

$$DH_2 = \{c_1 \cdot 1 + c_2 \cdot (x_1 - x_2) + c_3 \cdot (y_1 - y_2) : c_i \in \mathbb{C}\}.$$

Set $DH_n^{c,d}$ to be those $f \in DH_n$ whose total x degree is c and whose total y degree is d . The *bigraded Frobenius characteristic* of DH_n is the power series in q, t and $Z = \{z_1, z_2, \dots\}$:

$$DH_n(Z; q, t) = \sum_{c,d \geq 0} t^c q^d \sum_{\lambda \vdash n} s_\lambda \text{mult}(\chi^\lambda, \text{char } DH_n^{c,d}) = \nabla e_n,$$

where s_λ is a Schur function in variables Z , χ^λ is the character of $DH_n^{c,d}$.

6. The shuffle conjecture: setting

Diagonal harmonic polynomials are the solutions to a system of PDEs:

$$DH_n := \left\{ f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] : \sum_{i=1}^n \partial_{x_i}^a \partial_{y_i}^b f = 0 \text{ for all } a + b > 0 \right\}.$$

E.g.

$$DH_2 = \{c_1 \cdot 1 + c_2 \cdot (x_1 - x_2) + c_3 \cdot (y_1 - y_2) : c_i \in \mathbb{C}\}.$$

Set $DH_n^{c,d}$ to be those $f \in DH_n$ whose total x degree is c and whose total y degree is d . The *bigraded Frobenius characteristic* of DH_n is the power series in q, t and $Z = \{z_1, z_2, \dots\}$:

$$DH_n(Z; q, t) = \sum_{c,d \geq 0} t^c q^d \sum_{\lambda \vdash n} s_\lambda \text{mult}(\chi^\lambda, \text{char } DH_n^{c,d}) = \nabla e_n,$$

where s_λ is a Schur function in variables Z , χ^λ is the character of $DH_n^{c,d}$.

E.g.

$$\begin{aligned} DH_2(Z, q, t) &= \nabla e_2 = s_2 + (q + t)s_{11} \\ DH_3(Z, q, t) &= \nabla e_3 = s_3 + (q^2 + qt + t^2 + q + t)s_{21} \\ &\quad + (q^3 + q^2t + qt^2 + t^3 + qt)s_{111}. \end{aligned}$$

7. The shuffle conjecture: statement + special cases

The shuffle conjecture (and now theorem!) is a combinatorial interpretation for this power series:

Theorem (Haglund et al. 2003 (publ. 2005))

$$\nabla e_n = \sum_{\pi \in PF_n} t^{\text{area}(\pi)} q^{\text{dinv}(\pi)} F_{n, \text{ides}(\pi)},$$

where PF_n is the set of parking functions of order n , and $\text{area}(\pi)$ and $\text{dinv}(\pi)$ are two statistics on parking functions. $F_{n, \text{ides}(\pi)}$ is a fundamental quasi-symmetric function, and each such expression can be written in terms of Schur functions.

7. The shuffle conjecture: statement + special cases

The shuffle conjecture (and now theorem!) is a combinatorial interpretation for this power series:

Theorem (Haglund et al. 2003 (publ. 2005))

$$\nabla e_n = \sum_{\pi \in PF_n} t^{\text{area}(\pi)} q^{\text{dinv}(\pi)} F_{n, \text{ides}(\pi)},$$

where PF_n is the set of parking functions of order n , and $\text{area}(\pi)$ and $\text{dinv}(\pi)$ are two statistics on parking functions. $F_{n, \text{ides}(\pi)}$ is a fundamental quasi-symmetric function, and each such expression can be written in terms of Schur functions.

Theorem (Haglund 2005)

$$\langle \nabla e_{m+n-2}, h_{m-1} h_{n-1} \rangle = \sum_{\pi \in \text{Park}_{m-1, n-1}} t^{\text{area}(\pi)} q^{\text{dinv}(\pi)} =: \text{Park}_{m-1, n-1}(q, t),$$

where $\text{Park}_{m-1, n-1}$ is the set of all parking functions π of order $m+n-2$ whose reading word $\sigma(\pi)$ is a shuffle of the sequences $(1, \dots, m-1)$ and $(m, \dots, m+n-2)$.

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

The same recursion holds for the function

$$(qt)^{m+n-1} \text{Park}_{n-1, m-1}^{(r, s-1)}$$

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

The same recursion holds for the function

$$(qt)^{m+n-1} \text{Park}_{n-1, m-1}^{(r, s-1)}$$

and this provides the following connection

Theorem (Aval, D'adderio, D, Hicks, Le Borgne 2014)

$$\text{Nara}_{m,n}(q, t) = (qt)^{m+n-1} \langle \nabla e_{m+n-2}, h_{m-1} h_{n-1} \rangle.$$

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

The same recursion holds for the function

$$(qt)^{m+n-1} \text{Park}_{n-1, m-1}^{(r, s-1)}$$

and this provides the following connection

Theorem (Aval, D'adderio, D, Hicks, Le Borgne 2014)

$$\text{Nara}_{m,n}(q, t) = (qt)^{m+n-1} \langle \nabla e_{m+n-2}, h_{m-1} h_{n-1} \rangle.$$

Symmetry in q and t of the above expression is a property of the ∇ operator.

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

The same recursion holds for the function

$$(qt)^{m+n-1} \text{Park}_{n-1, m-1}^{(r, s-1)}$$

and this provides the following connection

Theorem (Aval, D'adderio, D, Hicks, Le Borgne 2014)

$$\text{Nara}_{m,n}(q, t) = (qt)^{m+n-1} \langle \nabla e_{m+n-2}, h_{m-1} h_{n-1} \rangle.$$

Symmetry in q and t of the above expression is a property of the ∇ operator. Symmetry in m and n is easily seen to hold due to the form of the RHS in the theorem.

8. Symmetry established

By conditioning on the first 'bounce' in a parallelogram polyomino, we find the following recursion holds for $\text{Nara}_{m,n}(q, t)$:

$$\text{Nara}_{m,n}^{(r,s)}(q, t) = t^{m+n-1} q^r \sum_{h=1}^{n-r} \sum_{k=0}^{m-s-1} q^s \binom{s+r-1}{s}_q \binom{s+h-1}{h}_q \text{Nara}_{n-r, m-s}^{(h,k)}(q, t).$$

The same recursion holds for the function

$$(qt)^{m+n-1} \text{Park}_{n-1, m-1}^{(r, s-1)}$$

and this provides the following connection

Theorem (Aval, D'adderio, D, Hicks, Le Borgne 2014)

$$\text{Nara}_{m,n}(q, t) = (qt)^{m+n-1} \langle \nabla e_{m+n-2}, h_{m-1} h_{n-1} \rangle.$$


Symmetry in q and t of the above expression is a property of the ∇ operator. Symmetry in m and n is easily seen to hold due to the form of the RHS in the theorem.

Corollary


$$\text{Nara}_{m,n}(q, t) = \text{Nara}_{m,n}(t, q) \text{ and } \text{Nara}_{m,n}(q, t) = \text{Nara}_{n,m}(q, t).$$

9. Chip-firing on the complete split graph

Complete split graph $S_{m,n}$:




\xrightarrow{g}




A representation of the recurrent states $\text{rec}(S_{m,n})$ comes in the form of a new type of polyomino that we call *sawtooth polyominoes*.

9. Chip-firing on the complete split graph

Complete split graph $S_{m,n}$:



g
 \rightarrow




A representation of the recurrent states $\text{rec}(S_{m,n})$ comes in the form of a new type of polyomino that we call *sawtooth polyominoes*.

Theorem (Derycke, D, and Le Borgne 2024)

$c \in \text{rec}(S_{m,n})$ iff $g(c) \in \text{Sawtooth}_{m,n}$.

10. Another surprise




$$\text{area}(P) = 12$$

$$\begin{aligned}\text{parabounce}(P) &= 1 \cdot (2 + 1) + 2 \cdot (2 + 1) \\ &\quad + 3 \cdot (0 + 1) + 4 \cdot (1 + 1) \\ &= 20.\end{aligned}$$

Let

$$F_{n,d}^{ITC}(q, t) := \sum_{P \in \text{Sawtooth}_{m,n}} q^{\text{area}(P)} t^{\text{itcbounce}(P)}.$$

10. Another surprise



$$\text{area}(P) = 12$$

$$\begin{aligned} \text{parabounce}(P) &= 1 \cdot (2 + 1) + 2 \cdot (2 + 1) \\ &\quad + 3 \cdot (0 + 1) + 4 \cdot (1 + 1) \\ &= 20. \end{aligned}$$

Let

$$F_{n,d}^{ITC}(q, t) := \sum_{P \in \text{Sawtooth}_{m,n}} q^{\text{area}(P)} t^{\text{itcbounce}(P)}.$$

Theorem (Derycke, D, and Le Borgne 2024)

$$\begin{aligned} F_{n,d}^{ITC}(q, t) &= \sum_{k=1}^n \sum_{\substack{(b_1, \dots, b_{k+1}) \models_k^* d \\ (a_1, \dots, a_k) \models_k n}} \prod_{i=1}^{k+1} q^{\binom{a_i}{2}} \binom{a_i + b_i}{b_i}_q \binom{a_i + b_i + a_{i-1} - 1}{a_{i-1} - 1}_q t^{(i-1)(a_i + b_i)} \\ &= \langle \nabla e_{n+d}, e_n h_d \rangle. \end{aligned}$$

Symmetry in q and t follows.

11. A more general result

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be a pair of integer compositions with $n = |\lambda| + |\mu|$.

11. A more general result

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be a pair of integer compositions with $n = |\lambda| + |\mu|$.

Let $G_{\lambda, \mu}$ be the graph that consists of cliques $K_{\lambda_1}, K_{\lambda_2}, \dots$, and independent sets $I_{\mu_1}, I_{\mu_2}, \dots$: every pair of vertices that are members of different sets (be they cliques/independent sets) are connected by an edge.

11. A more general result

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be a pair of integer compositions with $n = |\lambda| + |\mu|$.

Let $G_{\lambda, \mu}$ be the graph that consists of cliques $K_{\lambda_1}, K_{\lambda_2}, \dots$, and independent sets $I_{\mu_1}, I_{\mu_2}, \dots$: every pair of vertices that are members of different sets (be they cliques/independent sets) are connected by an edge.

Theorem (D'Adderio, D, Iraci, Lazar, Le Borgne, Vanden Wyngaerd 2024)

For every pair of compositions λ, μ such that $n = |\lambda| + |\mu|$,

$$\langle \nabla e_n, e_\lambda h_\mu \rangle = \sum_{c \in \text{rec}(G_{\lambda, \mu})} q^{\text{level}(c)} t^{\text{delaybounce}(c)}.$$

Symmetry in q and t follows from this form. (Each of the coefficients of ∇e_n when written as a linear combination of Schur functions is a symmetric function.)

11. A more general result

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be a pair of integer compositions with $n = |\lambda| + |\mu|$.

Let $G_{\lambda, \mu}$ be the graph that consists of cliques $K_{\lambda_1}, K_{\lambda_2}, \dots$, and independent sets $I_{\mu_1}, I_{\mu_2}, \dots$: every pair of vertices that are members of different sets (be they cliques/independent sets) are connected by an edge.

Theorem (D'Adderio, D, Iraci, Lazar, Le Borgne, Vanden Wyngaerd 2024)

For every pair of compositions λ, μ such that $n = |\lambda| + |\mu|$,

$$\langle \nabla e_n, e_\lambda h_\mu \rangle = \sum_{c \in \text{rec}(G_{\lambda, \mu})} q^{\text{level}(c)} t^{\text{delaybounce}(c)}.$$

Symmetry in q and t follows from this form. (Each of the coefficients of ∇e_n when written as a linear combination of Schur functions is a symmetric function.)

Special cases of the above theorem include the results for the complete bipartite graph and the complete split graph.

12. Comments

- ▶ These instances of symmetry, that are related to the shuffle conjecture, suggest something more general is afoot. An interesting question to consider is whether there is a parameterized graph whose bivariate q, t -polynomial (in the sense of what we have discussed) corresponds to other instances of the inner product of ∇e_n with some other symmetric functions.

12. Comments

- ▶ These instances of symmetry, that are related to the shuffle conjecture, suggest something more general is afoot. An interesting question to consider is whether there is a parameterized graph whose bivariate q, t -polynomial (in the sense of what we have discussed) corresponds to other instances of the inner product of ∇e_n with some other symmetric functions.
- ▶ Somewhat absent from our discussion is the choice of toppling regimes, and how bivariate polynomials related to them seem to be equal such as $F_{n,k}^{ITC}(q, t) = F_n^{CTI}(q, t)$. This is another instance of unexplained symmetry.

12. Comments

- ▶ These instances of symmetry, that are related to the shuffle conjecture, suggest something more general is afoot. An interesting question to consider is whether there is a parameterized graph whose bivariate q, t -polynomial (in the sense of what we have discussed) corresponds to other instances of the inner product of ∇e_n with some other symmetric functions.
- ▶ Somewhat absent from our discussion is the choice of toppling regimes, and how bivariate polynomials related to them seem to be equal such as $F_{n,k}^{ITC}(q, t) = F_n^{CTI}(q, t)$. This is another instance of unexplained symmetry.
- ▶ Symmetries that we are able to explain are thanks to known results related to the theory of symmetric functions. Absent from this is a combinatorial proof of the symmetries.

12. Comments

- ▶ These instances of symmetry, that are related to the shuffle conjecture, suggest something more general is afoot. An interesting question to consider is whether there is a parameterized graph whose bivariate q, t -polynomial (in the sense of what we have discussed) corresponds to other instances of the inner product of ∇e_n with some other symmetric functions.
- ▶ Somewhat absent from our discussion is the choice of toppling regimes, and how bivariate polynomials related to them seem to be equal such as $F_{n,k}^{ITC}(q, t) = F_n^{CTI}(q, t)$. This is another instance of unexplained symmetry.
- ▶ Symmetries that we are able to explain are thanks to known results related to the theory of symmetric functions. Absent from this is a combinatorial proof of the symmetries.

Thanks for your attention!