

Regular digraphs from finite groups

Andrea Švob (asvob@math.uniri.hr)

Faculty of Mathematics, University of Rijeka, Croatia

Joint work with A. E. Brouwer, D. Crnković, M. Zubović Žutolija and T. Zrinski

15th Nordic Combinatorial Conference (NORCOM 2025)
Reykjavik, Iceland, June, 2025



This work has been supported by Croatian Science Foundation under the project HRZZ-UIP-2020-02-5713.

18th June 2025

The outline of the talk:

- ① The method
- ② Examples of dsrgs
- ③ Orbit matrices of directed strongly regular graphs

A t -(v, k, λ) **design** is a finite incidence structure $\mathcal{D} = (\mathcal{P}, \mathcal{B}, \mathcal{I})$ satisfying the following requirements:

- 1 $|\mathcal{P}| = v$,
- 2 every element of \mathcal{B} is incident with exactly k elements of \mathcal{P} ,
- 3 every t elements of \mathcal{P} are incident with exactly λ elements of \mathcal{B} .

Every element of \mathcal{P} is incident with exactly r elements of \mathcal{B} .

The number of blocks is denoted by b .

If $b = v$ (or equivalently $k = r$) then the design is called **symmetric**.

- A $2-(v, k, \lambda)$ design is called a block design.
- If \mathcal{D} is a t -design, then it is also a s -design, for $1 \leq s \leq t - 1$.
- An **incidence matrix** of a design \mathcal{D} is a matrix $A = [a_{ij}]$ where $a_{ij} = 1$ if j th point is incident with the i th block and $a_{ij} = 0$ otherwise.




Figure: $2-(7, 3, 1)$ design

- D. Crnković, V. Mikulić Crnković, A. Švob, On some transitive combinatorial structures constructed from the unitary group $U(3, 3)$, J. Statist. Plann. Inference 144 (2014), 19–40.

Theorem

Let G be a finite permutation group acting transitively on the sets Ω_1 and Ω_2 of size m and n , respectively. Let $\alpha \in \Omega_1$ and $\Delta_2 = \bigcup_{i=1}^s \delta_i G_\alpha$, where $\delta_1, \dots, \delta_s \in \Omega_2$ are representatives of distinct G_α -orbits. If $\Delta_2 \neq \Omega_2$ and

$$\mathcal{B} = \{\Delta_2 g : g \in G\},$$

then $\mathcal{D}(G, \alpha, \delta_1, \dots, \delta_s) = (\Omega_2, \mathcal{B})$ is a $1 - (n, |\Delta_2|, \frac{|G_\alpha|}{|G_{\Delta_2}|} \sum_{i=1}^s |\alpha G_{\delta_i}|)$ design with $\frac{m \cdot |G_\alpha|}{|G_{\Delta_2}|}$ blocks. The group $H \cong G / \bigcap_{x \in \Omega_2} G_x$ acts as an automorphism group on (Ω_2, \mathcal{B}) , transitively on points and blocks of the design. If $\Delta_2 = \Omega_2$ then the set \mathcal{B} consists of one block, and $\mathcal{D}(G, \alpha, \delta_1, \dots, \delta_s)$ is a design with parameters $1 - (n, n, 1)$.

- D. Crnković, V. Mikulić Crnković, A. Švob, New 3-designs and 2-designs having $U(3,3)$ as an automorphism group, Discrete Math. 340 (2017), 2507–2515.
- D. Crnković, A. Švob, New symmetric 2-(176,50,14) designs, Discrete Math. 344 (2021), 112623, 3 pages.
- D. Crnković, N. Mostarac, A. Švob, Distance-regular graphs and new block designs obtained from the Mathieu groups, Appl. Algebra Engrg. Comm. Comput. 35 (2024), 177–194.
- ...

If a group G acts transitively on Ω , $\alpha \in \Omega$, and Δ is an orbit of G_α , then $\Delta' = \{\alpha g \mid g \in G, \alpha g^{-1} \in \Delta\}$ is also an orbit of G_α . Δ' is called the orbit of G_α paired with Δ . It is obvious that $\Delta'' = \Delta$ and $|\Delta'| = |\Delta|$. If $\Delta' = \Delta$, then Δ is said to be self-paired.

Corollary

If $\Omega_1 = \Omega_2$ and Δ_2 is a union of self-paired and mutually paired orbits of G_α , then the design $\mathcal{D}(G, \alpha, \delta_1, \dots, \delta_s)$ is a symmetric self-dual design and the incidence matrix of that design is the adjacency matrix of a $|\Delta_2|$ -regular graph.

The construction described in Theorem gives us all simple designs on which the group G acts transitively on the points and blocks, *i.e.* if G acts transitively on the points and blocks of a simple 1-design \mathcal{D} , then \mathcal{D} can be obtained as described in Theorem.

Note that the construction from Theorem gives us 1-designs, and **the incidence matrices of some of these 1-designs may be the adjacency matrices of directed strongly regular graphs.**

Since the construction given in Theorem gives all designs having G as an automorphism group acting transitively on points and blocks, it gives us also **all directed strongly regular graphs admitting a transitive action of the set of vertices.**

Clearly, the adjacency matrix of a directed strongly regular graph with parameters (n, k, t, λ, μ) is the incidence matrix of a $1-(n, k, k)$ design. In that way, the neighbourhoods of a directed strongly regular graph correspond to the blocks of a design, where the neighbourhood of a vertex x is the set of all vertices y such that there is an arc $x \rightarrow y$.

dsrgs

A directed strongly regular graph with parameters (v, k, t, λ, μ) is a directed graph on v vertices without loops such that

- (i) every vertex has in-degree and out-degree k ,
- (ii) every vertex x has t out-neighbours that are also in-neighbours of x ,
- (iii) the number of directed paths of length 2 from a vertex x to another vertex y is λ if there is an edge from x to y , and is μ if there is no edge from x to y .

- A. E. Brouwer, D. Crnković, A. Švob, A construction of directed strongly regular graphs with parameters $(63, 11, 8, 1, 2)$, Discrete Math. 347 (2024), 114146, 3 pages.

Theorem

Up to isomorphism, there are exactly two directed strongly regular graphs with parameters $(63, 11, 8, 1, 2)$ on which the linear group $PSL(2, 8)$ acts transitively. These directed strongly regular graphs have $PSL(2, 8) : Z_3$ as the full automorphism group.

The linear group $PSL(2, 8)$ is the simple group of order 504 and up to conjugation it has exactly one subgroup of order 8, which is isomorphic to the elementary abelian group E_8 .

By taking $G = PSL(2, 8)$ and $G_\alpha = E_8$, we constructed two non-isomorphic directed strongly regular graphs with parameters $(63, 11, 8, 1, 2)$, both having $PSL(2, 8) : Z_3$ as the full automorphism group.

Description:

Let Γ be a distance-regular graph with intersection array $\{q, q-2, 1; 1, q-2, q\}$, an antipodal $(q-1)$ -cover of the complete graph K_{q+1} , with $q^2 - 1$ vertices and $\lambda = \mu = 1$.


Remark

The automorphism group G of Γ is the semilinear group $P\Sigma L(2, q)$, acting transitively and edge-transitively on Γ . The orbits of the stabilizer in $L_2(q)$ of the vertex p (which is elementary abelian of order q) are the $q-1$ singletons in the antipodal class of p together with the $q-1$ sets of size q that are the conics with equations $Q(p)Q(x) + c(x, p)^2 = 0$ for $c \in \mathbb{F}_q \setminus \{0\}$ minus the point $C \cap p^\perp$.

In the special case $q = 8$, with $G = L_2(8) : 3$, the six antipodes of a point p fall into two orbits of size 3.

Adding to Γ (viewed as a directed graph by viewing an undirected edge xy as a pair of directed edges xy and yx) arrows from p to the three antipodes in one orbit, and adding all images of these arrows under G , produces a directed strongly regular graph with parameters $(63, 11, 8, 1, 2)$.

Let Δ_1 and Δ_2 be the two dsrgs obtained for the two choices of orbit, with adjacency matrices A_1 and A_2 , respectively. Then Δ_2 is not isomorphic to Δ_1 , but Δ_2 is isomorphic to the dsrg with adjacency matrix A_1^T .

$\text{dsrg}(63, 11, 8, 1, 2)$
 Δ :


$$v = 63$$

- A. E. Brouwer, D. Crnković, A. Švob, M. Zubović Žutolija, On some directed strongly regular graphs constructed from linear groups, preprint, 2025.

We classify directed strongly regular graphs admitting a transitive action of the linear groups $L_2(q)$, $q \leq 32$, $L_3(q)$, $q \leq 7$, and $L_4(2)$, for which the rank of the permutation representation is at most 20 (i.e. the number of orbits of the stabilizer of a vertex is at most 20).

We have looked at other small simple groups, of order at most 10^6 .

Several of the graphs found have a so far unknown parameter set.

Some examples generalize to an infinite family.

Main steps:

- The method of construction applied for obtaining dsrgs.
- Checking for each of the constructed digraphs if its parameters can be obtained from a construction described by A. E. Brouwer and S. A. Hobart database of dsrgs.
- Checking the isomorphism with the known digraphs.
- Finding the constructions for parametrically new dsrgs.

(126,40,20,14,12), (126,45,25,16,16), (140,19,7,6,2), (165,60,36,23,21), (240,50,32,4,12),
 (280,39,15,14,4), (280,123,63,54,54), (288,70,28,20,16), (288,77,35,22,20), (336,89,59,22,24),
 (420,167,68,67,66), (465,156,56,51,53), (560,117,39,26,24), (560,234,156,100,96),
 (750,224,84,83,60), (930,313,111,106,105), (1360,159,39,38,16), (3192,794,206,196,198)

Table: New dsrg parameter sets

A geometric construction

Let q be a prime power. Construct a directed graph as follows. Take as vertices the ordered pairs (s, t) of distinct points in a projective plane $PG(2, q)$ (desarguesian or not). Let L_{st} be the line on s and t . Make an arrow $(s, t) \rightarrow (u, v)$ when either $(L_{st} = L_{uv} \text{ and } u = t)$ or $(L_{st} \neq L_{uv} \text{ and } u = L_{st} \cap L_{uv})$.

The graph thus obtained is a directed strongly regular graph with parameters $n = q(q+1)(q^2+q+1)$, $k = q(q^2+q+1)$, $t = q^2+1$, $\lambda = q^2$, $\mu = q^2+1$. The graph is invariant under $P\Gamma L(3, q)$.

A partition $\Pi = \{C_0, C_1, \dots, C_{t-1}\}$ of the vertex set of a graph Γ is called *equitable* (or *regular*) if for each pair of (possibly equal) indices $i, j \in \{0, 1, \dots, t-1\}$ there exists a nonnegative integer b_{ij} so that every vertex $x \in C_i$ is adjacent with b_{ij} vertices in C_j , independent of the selection of x .

The $t \times t$ matrix $B = [b_{ij}]$ is called a *quotient matrix* of Γ with respect to the equitable partition Π .

The action of an automorphism group G of a dsrg induces equitable partition of its vertex set. The corresponding quotient matrix is called an orbit matrix of the dsrg with respect to the action of G .

Theorem [D. Crnković, T. Zrinski, AŠ, 202?]

Let Γ be a $\text{dsrg}(v, k, t, \lambda, \mu)$ and let G be an automorphism group of Γ . Further, let O_1, O_2, \dots, O_b be the G -orbits on vertex set of Γ , and let $|O_i| = n_i$, $i = 1, \dots, b$, be the corresponding orbit lengths. If $R = [r_{ij}]$ is the row orbit matrix of Γ with respect to the action of G , then the following hold

$$0 \leq r_{ij} \leq n_j, \text{ for } 1 \leq i, j \leq b,$$

$$0 \leq r_{ii} \leq n_i - 1, \text{ for } 1 \leq i \leq b,$$

$$\sum_{j=1}^b r_{ij} = k, \text{ for } 1 \leq i \leq b,$$

$$\sum_{s=1}^b r_{is} r_{sj} = \delta_{ij}(t - \mu) + r_{ij}\lambda + (n_j - r_{ij})\mu, \text{ for } 1 \leq i, j \leq b,$$

where δ_{ij} is the Kronecker delta.

Definition

Let n_i , $i = 1, \dots, b$, be positive integers such that $\sum_{i=1}^b n_i = v$. A $(b \times b)$ -matrix $R = [r_{ij}]$, where r_{ij} , $1 \leq i, j \leq b$, are non-negative integers satisfying conditions given by Theorem, is a **row orbit matrix associated with a $\text{dsrg}(v, k, t, \lambda, \mu)$ and the distribution of orbit lengths (n_1, n_2, \dots, n_b)** .

Remark

Orbit matrices from Definition may or may not correspond to a directed strongly regular graph with parameters (v, k, t, λ, μ) . Those matrices can be used for constructing directed strongly regular graphs with a presumed automorphism group in a similar way as orbit matrices of 2-designs are used for a construction of 2-designs and orbit matrices of strongly regular graphs are used for a construction of strongly regular graphs

Thank you very much for your attention!

