

Naturally labelled posets and a hierarchy related to interval orders

Sergey Kitaev

University of Strathclyde

Joint work with David Bevan and Gi-Sang Cheon

Einar Fest

The Party After My PhD Defense (January 2003)



The Party After My PhD Defense (January 2003)



The Party After My PhD Defense (January 2003)



The Party After My first student's PhD Defense (2019)



Meet the research team!



David Bevan



Gi-Sang Cheon



SK

European Journal of Combinatorics 126 (2025) 104117



Contents lists available at [ScienceDirect](#)

European Journal of Combinatorics

journal homepage: <https://www.sciencedirect.com/journal/european-journal-of-combinatorics>



On naturally labelled posets and permutations avoiding 12-34

David Bevan ^a, Gi-Sang Cheon ^b, Sergey Kitaev ^a

^a Department of Mathematics and Statistics, University of Strathclyde, Glasgow, Scotland, United Kingdom

^b Department of Mathematics, Sungkyunkwan University, Suwon, South Korea



Naturally labelled posets

A **partially ordered set (poset)** of cardinality n is **labelled** if its elements are in $[n] := \{1, 2, \dots, n\}$.

Naturally labelled poset

A poset (P, \prec) is **naturally labelled** (or **natural** for brievity) if $i \prec j$ implies $i < j$ (in the usual order).


Naturally labelled posets

A **partially ordered set (poset)** of cardinality n is **labelled** if its elements are in $[n] := \{1, 2, \dots, n\}$.

Naturally labelled poset


A poset (P, \prec) is **naturally labelled** (or **natural** for brevity) if $i \prec j$ implies $i < j$ (in the usual order).

Non-example of a naturally labelled poset



$(3 \prec 1 \text{ but } 3 \not\prec 1)$


Example of a naturally labelled poset



Naturally labelled posets

Natural posets on $[n]$ are **in bijection** with **lower triangular binary matrices** with **ones on the main diagonal** that contain no $(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix})$ submatrix whose **upper right entry** (shown in bold) is on the **main diagonal**.

Example




\leftrightarrow

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

Naturally labelled posets

Natural posets on $[n]$ are in bijection with lower triangular binary matrices with ones on the main diagonal that contain no $(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix})$ submatrix whose upper right entry (shown in bold) is on the main diagonal.

Example



\leftrightarrow


$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

These are counted by A006455 in OEIS. Connected natural posets are counted by A323502 in OEIS.

Brightwell, Prömel and Steger ("The average number of linear extensions of a partial order. *J. Comb. Theory Ser. A*, 73(2):193–206, 1996.") prove that the number of natural posets on $[n]$ is asymptotically $C_{\text{even}} n 2^{n^2/4}$ for even n and $C_{\text{odd}} n 2^{n^2/4}$ for odd n , where the constants differ by less than 10^{-4} and both are approximately equal to 12.7636.

3-free naturally labelled posets

Non-example of a 3-free natural poset




(elements 1, 3, 5 form a chain of length 3)

Natural posets on $[n]$ of **height at most 2** are in bijection with **lower triangular binary matrices** with **ones on the main diagonal** that contain no $\begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}$ whose **upper right entry** is on the **main diagonal**.

3-free naturally labelled posets

Non-example of a 3-free natural poset



(elements 1, 3, 5 form a chain of length 3)

Natural posets on $[n]$ of **height at most 2** are in bijection with **lower triangular binary matrices** with **ones on the main diagonal** that contain no $\begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}$ whose **upper right entry** is on the **main diagonal**.

Stanley graphs

A labelled graph on $[n]$ is a **Stanley graph** iff no vertex v has **both** left neighbours ($u < v$) and right neighbours ($u > v$). Clearly, every Stanley graph is **bipartite**. Assigning the **minimal elements** of a 3-free natural poset to one part of a bipartite graph and the remaining elements to the other part gives a **bijection** with Stanley graphs.

3-free naturally labelled posets

Stanley graphs are counted by A135922 in [OEIS](#). These numbers grow like $7.371968\dots \cdot 2^{n^2/4}$.

The **generating function** for the number of **3-free natural posets** with k minimal elements is

$$P_k(t) = \frac{t^k}{(1-t)(1-3t)\cdots(1-(2^k-1)t)}, \quad (k=0,1,\dots).$$

Hence, the **generating function** for all **3-free natural posets** is given by

$$\sum_{k \geq 0} P_k(t) = \sum_{k \geq 0} \frac{t^k}{\prod_{i=0}^k (1-(2^i-1)t)}.$$

Alternatively, the **generating function** is $F(t, 1)$, where

$$F(t, y) = 1 + t((y-1)F(t, y) + F(t, 2y)).$$

3-free naturally labelled posets

Stanley graphs are counted by A135922 in [OEIS](#). These numbers grow like $7.371968\dots \cdot 2^{n^2/4}$.

The **generating function** for the number of **3-free natural posets** with k minimal elements is

$$P_k(t) = \frac{t^k}{(1-t)(1-3t)\cdots(1-(2^k-1)t)}, \quad (k=0,1,\dots).$$

Hence, the **generating function** for all **3-free natural posets** is given by

$$\sum_{k \geq 0} P_k(t) = \sum_{k \geq 0} \frac{t^k}{\prod_{i=0}^k (1-(2^i-1)t)}.$$

Alternatively, the **generating function** is $F(t, 1)$, where

$$F(t, y) = 1 + t((y-1)F(t, y) + F(t, 2y)).$$

(For connected such posets the **generating function** is $G(1, 0)$, where $G(u, v) = 1 + t((v-1)G(u, v) + G(2u, u+v))$; A323843 in [OEIS](#).)

An influential paper



[Home](#) [Preferences](#) [Free Tools](#) [Help](#) [Contact Us](#) [Terms of Use](#) [Blog](#)

University of Strathclyde



[Previous](#) [Up](#) [Next](#)

Select alternative format

Publications results for "Author=(Kitaev, S*)"

MR2652101 (2011h:05022)

Bousquet-Mélou, Mireille (F-BORD-LB); Claesson, Anders (ICE-RU-IM); Dukes, Mark (ICE-UICE-SI); Kitaev, Sergey (ICE-RU-IM)

(2 + 2)-free posets, ascent sequences and pattern avoiding permutations. (English summary)

J. Combin. Theory Ser. A 117 (2010), no. 7, 884–909.

05A19 (05A05 06A07)

[Review PDF](#) | [Clipboard](#) | [Journal](#) | [Article](#) | [Make Link](#)

Citations

From References: 92

From Reviews: 21

[Find it @Strathclyde](#)



Mireille Bousquet-Mélou



Anders Claesson




Mark Dukes



SK

A hierarchy of combinatorial objects related to interval orders



2+2-free naturally labelled posets

2+2-free poset


A poset is **2+2-free** if it does not contain an **induced** subposet that is isomorphic to 2+2, the **union of two disjoint 2-element chains**.

2+2-free naturally labelled posets

2+2-free poset


A poset is **2+2-free** if it does not contain an **induced** subposet that is isomorphic to 2+2, the **union of two disjoint 2-element chains**.

Non-example of a 2+2-free natural poset



(the elements 3, 5 and 2, 4 form 2+2)

Example of a 2+2-free natural poset



2+2-free naturally labelled posets

Brute-force generation yields the following enumeration of 2+2-free natural posets that is **not** in **OEIS**: 1, 2, 7, 37, 272, 2637, 32469,

Open problem

Find a **recurrence** / **functional equation** / **generating function** / **formula** for 2+2-free naturally labelled posets.

2+2-free naturally labelled posets


Brute-force generation yields the following enumeration of 2+2-free natural posets that is **not** in [OEIS](#): 1, 2, 7, 37, 272, 2637, 32469,

Open problem

Find a **recurrence** / **functional equation** / **generating function** / **formula** for 2+2-free naturally labelled posets.

This problem is **hard**, but imposing **extra restrictions** may give interesting results. [Claesson](#) and [Linusson](#) (“ $n!$ matchings, $n!$ posets. *Proc. Amer. Math. Soc.*, **139**(2):435–449, 2011”)) prove that there are exactly $n!$ **factorial posets**, i.e. naturally labelled posets s.t. $i < j \prec k \Rightarrow i \prec k$ (these posets are proved to be **2+2-free**).

Non-example of a factorial poset



Naturally labelled, but $1 < 2 \prec 3$ while $1 \not\prec 3$.

$\{3,2+2\}$ -free naturally labelled posets

Brute-force generation yields the following for the enumeration of **$\{3,2+2\}$ -free natural posets** on $[n]$:

$$1, 2, 6, 23, 107, 585, 3669, 25932, \dots$$

This matches A113226 in [OEIS](#) enumerating permutations avoiding the [vincular pattern](#) 12-34 ($= \underline{12} \underline{34}$); a permutation $\pi_1 \pi_2 \cdots \pi_n$ avoids the pattern if there is no $i < j - 1$ such that $\pi_i < \pi_{i+1} < \pi_j < \pi_{j+1}$.

$\{3,2+2\}$ -free naturally labelled posets

Brute-force generation yields the following for the enumeration of $\{3,2+2\}$ -free natural posets on $[n]$:

$$1, 2, 6, 23, 107, 585, 3669, 25932, \dots$$


This matches A113226 in [OEIS](#) enumerating permutations avoiding the **vincular pattern** 12-34 ($= \underline{12} \underline{34}$); a permutation $\pi_1 \pi_2 \cdots \pi_n$ avoids the pattern if there is no $i < j - 1$ such that $\pi_i < \pi_{i+1} < \pi_j < \pi_{j+1}$.

We proved the connection **bijectively** via the **decomposition** of **unlabelled** 2+2-free posets, and using a bijection between the set of $\{3,2+2\}$ -free n -element natural posets and certain decorated binary words of length n .

Theorem

$\{3,2+2\}$ -free naturally labelled posets are in bijection with 12-34-avoiding permutations.

The place of naturally labelled posets in the hierarchy



Thank you for your attention!

Any questions?