

Monochromatic structures in two-edge-colored ordered complete graphs


János Barát

Alfréd Rényi Institute of Mathematics, Budapest, Hungary
University of Pannonia, Veszprém, Hungary

16.06.2025

joint work with András Gyárfás and Géza Tóth

Co-authors



Preliminary results

Preliminary results

Every 2-colored complete graph has a monochromatic spanning tree.

Preliminary results

Every 2-colored complete graph has a monochromatic spanning tree.

Every 2-colored complete graph K_{3n-1} contains a monochromatic matching M_n and this is not true for K_{3n-2} .

Preliminary geometric results

Preliminary geometric results

Károlyi, Pach and Tóth generalized both results to **geometric graphs**, that are graphs drawn in the plane with straight-line segments as edges.

Preliminary geometric results

Károlyi, Pach and Tóth generalized both results to **geometric graphs**, that are graphs drawn in the plane with straight-line segments as edges.

Theorem (KPT)

Every 2-colored complete geometric graph has a monochromatic plane spanning tree.

Theorem (KPT)

Every 2-colored complete geometric graph K_{3n-1} contains a monochromatic plane matching M_n .

Here a **plane** subgraph is one, whose edges in the embedding do not have common internal points.

Our starting point

Our starting point

Conjecture: Every 2-colored complete simple drawing has a monochromatic plane spanning tree.

Our starting point

Conjecture: Every 2-colored complete simple drawing has a monochromatic plane spanning tree.

OPEN


GOAL: Show it for the twisted drawing (Harborth, Mengersen '92)

Our starting point

Conjecture: Every 2-colored complete simple drawing has a monochromatic plane spanning tree.

OPEN

GOAL: Show it for the twisted drawing (Harborth, Mengersen '92)



Ordered graphs

Ordered graphs

An **ordered graph** G is a simple graph with $V(G) = [m] = \{1, 2, \dots, m\}$.

We also use $[i, j] = \{i, i+1, \dots, j\}$

The vertex set is considered with the natural ordering and the edges are denoted by (i, j) , where we always assume $i < j$.

The length of (i, j) is $j-i$.

Independent edges


Independent edges

Independent edges in ordered graphs can be classified as follows


Independent edges

Independent edges in ordered graphs can be classified as follows

- Edges (a, b) and (c, d) are **crossing** if either $a < c < b < d$ or $c < a < d < b$.
- Edges (a, b) and (c, d) are **nested** if either $a < c < d < b$ or $c < a < b < d$.
- Edges (a, b) and (c, d) are **separated** if either $a < b < c < d$ or $c < d < a < b$.



an ordered graph



6 types of questions

6 types of questions

	length of the	longest path	
	lower bound	upper bound	
<i>nested</i>	$n/4$	$n/2 + 1$	
<i>crossing</i>	$n/4$	$n/2$	
<i>separated</i>	\sqrt{n}	$\sqrt{n} + 2$	
<i>non-nested</i>	$n/4$	$2n/3$	→ twisted drawing
<i>non-crossing</i>	$\lfloor \frac{n+1}{2} \rfloor$	$\lfloor \frac{n+1}{2} \rfloor$	→ convex drawing
<i>non-separated</i>	$n/4$	$n/2$	

Monochromatic spanning trees

Monochromatic spanning trees

Theorem (JB, AGy, GT)

In every 2-coloring of the ordered complete graph, there exists

- (i) a monochromatic non-crossing spanning tree.
- (ii) a monochromatic non-nested spanning tree.
- (iii) a monochromatic non-separated spanning tree.

Monochromatic spanning trees

Theorem (JB, AGy, GT)

In every 2-coloring of the ordered complete graph, there exists

- (i) a monochromatic non-crossing spanning tree.
- (ii) a monochromatic non-nested spanning tree.
- (iii) a monochromatic non-separated spanning tree.

(i) Bialostocki and Dierker conjectured. Proved by KPT.

Monochromatic spanning trees

Theorem (JB, AGy, GT)

In every 2-coloring of the ordered complete graph, there exists

- (i) a monochromatic non-crossing spanning tree.
- (ii) a monochromatic non-nested spanning tree.
- (iii) a monochromatic non-separated spanning tree.

- (i) Bialostocki and Dierker conjectured. Proved by KPT.
- (ii) connection to twisted drawings

Non-crossing spanning tree

Non-crossing spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(i) a monochromatic non-crossing spanning tree.

Non-crossing spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(i) a monochromatic non-crossing spanning tree.

If all edges $(i, i + 1)$, $i = 1, \dots, n - 1$ have the same color, then we are done, they form the desired spanning tree.

Non-crossing spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(i) a monochromatic non-crossing spanning tree.

If all edges $(i, i + 1)$, $i = 1, \dots, n - 1$ have the same color, then we are done, they form the desired spanning tree.

Otherwise there is an i , $1 < i < n$ such that $(i - 1, i)$ and $(i, i + 1)$ have different colors.

Non-crossing spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(i) a monochromatic non-crossing spanning tree.

If all edges $(i, i+1)$, $i = 1, \dots, n-1$ have the same color, then we are done, they form the desired spanning tree.

Otherwise there is an i , $1 < i < n$ such that $(i-1, i)$ and $(i, i+1)$ have different colors.

Delete vertex i . Use induction.

Non-nested spanning tree

Non-nested spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(ii) a monochromatic non-nested spanning tree.

Non-nested spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (ii) a monochromatic non-nested spanning tree.

Let c be any 2-coloring of the edges. $(1, 2)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done.

Non-nested spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (ii) a monochromatic non-nested spanning tree.

Let c be any 2-coloring of the edges. $(1, 2)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done.

Otherwise, let s be the smallest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s-1]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Non-nested spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(ii) a monochromatic non-nested spanning tree.

Let c be any 2-coloring of the edges. $(1, 2)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done.

Otherwise, let s be the smallest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s-1]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Suppose first that we find a blue spanning tree B without nested edges. Delete the edges in B induced by $[2, s-1]$. The resulting graph can also be found in the original coloring c . Now add the blue edges $(1, 2), (1, 3), \dots, (1, s-1)$.

Non-nested spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(ii) a monochromatic non-nested spanning tree.

Let c be any 2-coloring of the edges. $(1, 2)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done.

Otherwise, let s be the smallest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s-1]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Suppose first that we find a blue spanning tree B without nested edges. Delete the edges in B induced by $[2, s-1]$. The resulting graph can also be found in the original coloring c . Now add the blue edges $(1, 2), (1, 3), \dots, (1, s-1)$.

Suppose now that we found a red spanning tree R on $[2, n]$. It cannot contain any edges induced by $[2, s-1]$ since they are blue. So, R can also be found in the original coloring c . Simply add edge $(1, s)$.

Non-separated spanning tree

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (iii) a monochromatic non-separated spanning tree.

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (iii) a monochromatic non-separated spanning tree.

Let c be any 2-coloring of the edges. $(1, n)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done,

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (iii) a monochromatic non-separated spanning tree.

Let c be any 2-coloring of the edges. $(1, n)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done,

Otherwise, let s be the largest number such that $(1, s)$ is red.

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists (iii) a monochromatic non-separated spanning tree.

Let c be any 2-coloring of the edges. $(1, n)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done,

Otherwise, let s be the largest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s+1, n]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(iii) a monochromatic non-separated spanning tree.

Let c be any 2-coloring of the edges. $(1, n)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done,

Otherwise, let s be the largest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s+1, n]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Suppose first that we find a blue spanning tree B without separated edges. Delete the edges of B induced by $[s+1, n]$. The resulting graph can also be found in the original coloring c . Now add the blue edges $(1, s+1), \dots, (1, n)$.

Non-separated spanning tree

In every 2-coloring of the ordered complete graph on $[n]$, there exists
(iii) a monochromatic non-separated spanning tree.

Let c be any 2-coloring of the edges. $(1, n)$ is blue. If $(1, i)$ is blue for every i , $2 \leq i \leq n$, then we are done,

Otherwise, let s be the largest number such that $(1, s)$ is red. We now change the coloring c to \tilde{c} as follows: we recolor each edge induced by $[s+1, n]$ blue, and keep c otherwise. Consider the coloring \tilde{c} on $[2, n]$ and apply the induction hypothesis.

Suppose first that we find a blue spanning tree B without separated edges. Delete the edges of B induced by $[s+1, n]$. The resulting graph can also be found in the original coloring c . Now add the blue edges $(1, s+1), \dots, (1, n)$.

Suppose now that we found a red spanning tree R on $[2, n]$. It cannot contain any edges induced by $[s+1, n]$ since they are blue. So, R can also be found in the original coloring c . Simply add edge $(1, s)$.

Monochromatic spanning subgraphs

Monochromatic spanning subgraphs

Theorem (JB, AGy, GT)

In every 2-coloring of the ordered complete graph, there exists

- (i) a monochromatic non-crossing spanning tree.
- (ii) a monochromatic non-nested spanning tree.
- (iii) a monochromatic non-separated spanning tree.

Monochromatic spanning subgraphs

Theorem (JB, AGy, GT)

In every 2-coloring of the ordered complete graph, there exists

- (i) a monochromatic non-crossing spanning tree.
- (ii) a monochromatic non-nested spanning tree.
- (iii) a monochromatic non-separated spanning tree.

Proposition (JB, AGy, GT)

- (i) There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.
- (ii) There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Non-crossing monochromatic subgraphs

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i + 1)$ blue, for $1 \leq i \leq n - 1$, all other edges red.

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i + 1)$ blue, for $1 \leq i \leq n - 1$, all other edges red.
Let H be a non-crossing subgraph with red edges.

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i+1)$ blue, for $1 \leq i \leq n-1$, all other edges red.

Let H be a non-crossing subgraph with red edges.

Consider the convex drawing of K_n , where the blue edges are on the outer cycle. Now H becomes a planar subgraph of the convex drawing.

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i+1)$ blue, for $1 \leq i \leq n-1$, all other edges red.

Let H be a non-crossing subgraph with red edges.

Consider the convex drawing of K_n , where the blue edges are on the outer cycle. Now H becomes a planar subgraph of the convex drawing.

Let us add all edges of the outer cycle ($n-1$ blue and possibly one red) to H to get H' .

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i+1)$ blue, for $1 \leq i \leq n-1$, all other edges red.

Let H be a non-crossing subgraph with red edges.

Consider the convex drawing of K_n , where the blue edges are on the outer cycle. Now H becomes a planar subgraph of the convex drawing.

Let us add all edges of the outer cycle ($n-1$ blue and possibly one red) to H to get H' .

It is still an outerplanar graph. Therefore, H' has at most $2n-3$ edges, $n-1$ of them are blue.

Non-crossing monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-crossing monochromatic subgraph with n edges.

Color all edges $(i, i+1)$ blue, for $1 \leq i \leq n-1$, all other edges red.

Let H be a non-crossing subgraph with red edges.

Consider the convex drawing of K_n , where the blue edges are on the outer cycle. Now H becomes a planar subgraph of the convex drawing.

Let us add all edges of the outer cycle ($n-1$ blue and possibly one red) to H to get H' .

It is still an outerplanar graph. Therefore, H' has at most $2n-3$ edges, $n-1$ of them are blue.

Thus H can have at most $n-2$ edges.

Non-nested monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Non-nested monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Color edge (i, j) blue if $i + j$ is even and red, if $i + j$ is odd.

Non-nested monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Color edge (i, j) blue if $i + j$ is even and red, if $i + j$ is odd.

For the red edges, the value of $i + j$ can have at most $n - 1$ different values.

Non-nested monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Color edge (i, j) blue if $i + j$ is even and red, if $i + j$ is odd.

For the red edges, the value of $i + j$ can have at most $n - 1$ different values.

So, among n red edges, there are two with $i + j = i' + j'$, therefore, these edges are nested.

Non-nested monochromatic subgraphs

There is a 2-coloring of the ordered complete graph on $[n]$, which does not contain a non-nested monochromatic subgraph with n edges.

Color edge (i, j) blue if $i + j$ is even and red, if $i + j$ is odd.

For the red edges, the value of $i + j$ can have at most $n - 1$ different values.

So, among n red edges, there are two with $i + j = i' + j'$, therefore, these edges are nested.

The argument is the same for the blue edges.

Non-separated subgraph

Non-separated subgraph

Proposition (JB, AGy, GT)

In any 2-coloring of the ordered complete graph on $[n]$, there is a non-separated monochromatic subgraph of $\lfloor n^2/8 \rfloor$ edges.

Non-separated subgraph

Proposition (JB, AGy, GT)

In any 2-coloring of the ordered complete graph on $[n]$, there is a non-separated monochromatic subgraph of $\lfloor n^2/8 \rfloor$ edges.

Consider all edges (i, j) with $i \leq \lfloor n/2 \rfloor < j$.

Non-separated subgraph

Proposition (JB, AGy, GT)

In any 2-coloring of the ordered complete graph on $[n]$, there is a non-separated monochromatic subgraph of $\lfloor n^2/8 \rfloor$ edges.

Consider all edges (i, j) with $i \leq \lfloor n/2 \rfloor < j$.

$$\# \text{ edges} = \frac{n^2}{4}$$

Non-separated subgraph

Proposition (JB, AGy, GT)

In any 2-coloring of the ordered complete graph on $[n]$, there is a non-separated monochromatic subgraph of $\lfloor n^2/8 \rfloor$ edges.

Consider all edges (i, j) with $i \leq \lfloor n/2 \rfloor < j$.

edges = $\frac{n^2}{4}$

no separated pairs

Non-separated subgraph

Proposition (JB, AGy, GT)

In any 2-coloring of the ordered complete graph on $[n]$, there is a non-separated monochromatic subgraph of $\lfloor n^2/8 \rfloor$ edges.

Consider all edges (i, j) with $i \leq \lfloor n/2 \rfloor < j$.

edges = $\frac{n^2}{4}$

no separated pairs

majority color

Non-nested matchings

Non-nested matchings

Every 2-colored complete graph K_{3n-1} contains a monochromatic matching M_n

Non-nested matchings

Every 2-colored complete graph K_{3n-1} contains a monochromatic matching M_n

Conjecture

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-nested matching M_n

Non-nested matchings

Every 2-colored complete graph K_{3n-1} contains a monochromatic matching M_n

Conjecture

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-nested matching M_n

Theorem (JB, AGy, GT)

If an ordered complete graph on $[3n-1]$ contains either

- (i) a red K_{2n-1} or
- (ii) a blue $K_{n-1,2n}$ as a subgraph, then there is a monochromatic non-nested M_n .

Non-crossing matchings

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.
here is a simple proof

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.
here is a simple proof
Consider the Hamiltonian path.

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.
here is a simple proof
Consider the Hamiltonian path.
Either it is monochromatic or

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.
here is a simple proof
Consider the Hamiltonian path.
Either it is monochromatic or
there are two consecutive edges of different colors.

Non-crossing matchings

Every 2-colored ordered complete graph on $[3n-1]$ contains a monochromatic non-crossing M_n .

This follows from a more general theorem by Károlyi, Pach and Tóth.
here is a simple proof
Consider the Hamiltonian path.
Either it is monochromatic or
there are two consecutive edges of different colors.
use induction.

Non-separated matchings

Non-separated matchings

Theorem (JB, AGy, GT)

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-separated M_n .

Non-separated matchings

Theorem (JB, AGy, GT)

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-separated M_n .

Consider the complete bipartite graph on $A = [1, n]$ and $B = [2n, 3n-1]$.

Non-separated matchings

Theorem (JB, AGy, GT)

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-separated M_n .

Consider the complete bipartite graph on $A = [1, n]$ and $B = [2n, 3n-1]$.
either this is monochromatic

Non-separated matchings

Theorem (JB, AGy, GT)

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-separated M_n .

Consider the complete bipartite graph on $A = [1, n]$ and $B = [2n, 3n-1]$.
either this is monochromatic
or we find a 2-colored V and use induction

Non-separated matchings

Theorem (JB, AGy, GT)

Every 2-colored ordered complete graph on $[3n - 1]$ contains a monochromatic non-separated M_n .

Consider the complete bipartite graph on $A = [1, n]$ and $B = [2n, 3n-1]$.
either this is monochromatic
or we find a 2-colored V and use induction
check the new edge and the non-separated property

Open questions

Open questions

Problem 1

What is the minimum number m such that every 2-colored ordered complete graph on $[m]$ contains a monochromatic non-nested M_n ?

Open questions

Problem 1

What is the minimum number m such that every 2-colored ordered complete graph on $[m]$ contains a monochromatic non-nested M_n ?

Problem 2

Show that every 2-colored ordered complete graph on $[11]$ contains a monochromatic non-nested M_4 .

Results on cycles

Results on cycles

	length of the	longest cycle	
	lower bound	upper bound	
<i>nested</i>	3	3	
<i>crossing</i>	$\log n$	$n/2$	
<i>separated</i>	3	3	
<i>non-nested</i>	$\log n$	$2n/3$	→ twisted drawing
<i>non-crossing</i>	$\sqrt{n/2}$	\sqrt{n}	→ convex drawing
<i>non-separated</i>	$n/8$	$n/2$	