

# Product representation of perfect cubes

Péter Pál Pach

MTA–HUN-REN RI “Momentum” Arithmetic Combinatorics Research Group  
HUN-REN Rényi Institute of Mathematics  
and  
Budapest University of Technology and Economics

18 June 2025

This research was supported by the Lendület (Momentum) Grant of the Hungarian Academy of Sciences and the Hungarian Scientific Research Funds (Grant Nr. OTKA NKFIH K146387).

Joint work with Zsigmond György Fleiner, Márk Hunor Juhász,  
Blanka Kövér and Csaba Sándor.

## Definition

$F_{k,d}(n)$ : maximal size of a set  $A \subseteq [n]$  such that

$$a_1 a_2 \dots a_k = x^d, \quad a_1 < a_2 < \dots < a_k$$

has no solution with  $a_1, a_2, \dots, a_k \in A$  and integer  $x$ .

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$



# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

- $F_{2,2}(n) = \left(\frac{6}{\pi^2} + o(1)\right) n$
- $\frac{n^{3/4}}{(\log n)^{3/2}} \ll F_{4,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$
- $\frac{n^{\frac{2\ell}{4\ell-1}}}{(\log n)^{\frac{4\ell}{4\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n) \ll \frac{n^{3/4}}{(\log n)^{3/2}}$
- $\frac{n^{\frac{2\ell+1}{4\ell+1}}}{(\log n)^{\frac{4\ell+2}{4\ell+1}}} \ll F_{4\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n$$

# Product representation of perfect squares

Erdős, Sárközy, T. Sós, 1995

$$\frac{n^{2/3}}{(\log n)^{4/3}} \ll F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{7/9} \log n$$

Győri, 1997

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{2/3} \log n$$

P., 2015

$$F_{6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll cn^{2/3} \log n / \log \log n$$

# Product representation of perfect squares

P., 2015

$$\frac{n^{3/5}}{(\log n)^{6/5}} \ll F_{8,2}(n) - \pi(n)$$

P., Vizer, 2023

- $\frac{n^{3/5}}{(\log n)^{6/5}} \ll F_{10,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right)$
- $\frac{n^{6/11}}{(\log n)^{12/11}} \ll F_{22,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right)$
- $\frac{n^{\frac{3\ell}{6\ell-2}}}{(\log n)^{\frac{3\ell}{3\ell-1}}} \ll F_{4\ell,2}(n) - \pi(n)$
- $\frac{n^{\frac{3\ell}{6\ell-1}}}{(\log n)^{\frac{6\ell}{6\ell-1}}} \ll F_{8\ell+2,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right)$
- $\frac{n^{\frac{6\ell-1}{12\ell-4}}}{(\log n)^{\frac{6\ell-1}{6\ell-2}}} \ll F_{8\ell+6,2}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right)$

# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{\varepsilon \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{e \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$$F_{5,2}(n) = (1 - o(1))n \text{ or } F_{5,2}(n) < (1 - c)n$$

# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{\varepsilon \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{\varepsilon \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$



# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{e \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

# Odd and even cases

## Generalized multiplicative Sidon sets

For  $k = 2K$ :

$$a_1 a_2 \dots a_K = b_1 b_2 \dots b_K \implies a_1 a_2 \dots a_K b_1 b_2 \dots b_K = x^2$$

Erdős, Sárközy, T. Sós, 1995

- $\frac{n}{(\log n)^{1+\varepsilon}} \ll n - F_{3,2}(n) \ll n(\log n)^{\frac{\varepsilon \log 2}{2} - 1 + \varepsilon}$
- $\liminf_{n \rightarrow \infty} \frac{F_{2\ell+1,2}(n)}{n} \geq \log 2 = 0.69 \dots$
- $\frac{n}{(\log n)^2} \ll n - F_{2\ell+1,2}(n)$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

## Tao, 2025

For  $k \geq 4$  we have  $F_{k,2}(n) \leq (1 - c + o(1))n$ .

## Granville, Soundararajan, 2001

For  $\ell \geq 2$  we have  $(1 - c_0 + o(1))n \leq F_{2\ell+1,2}(n)$ , where  $c_0 = 1 - \log(1 + \sqrt{e}) + 2 \int_1^{\sqrt{e}} \frac{\log t}{t+1} dt \approx 0.1715$  is the Hall-Montgomery constant.

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

## Tao, 2025

For  $k \geq 4$  we have  $F_{k,2}(n) \leq (1 - c + o(1))n$ .

## Granville, Soundararajan, 2001

For  $\ell \geq 2$  we have  $(1 - c_0 + o(1))n \leq F_{2\ell+1,2}(n)$ , where  $c_0 = 1 - \log(1 + \sqrt{e}) + 2 \int_1^{\sqrt{e}} \frac{\log t}{t+1} dt \approx 0.1715$  is the Hall-Montgomery constant.

## Question

$F_{5,2}(n) = (1 - o(1))n$  or  $F_{5,2}(n) < (1 - c)n$

## Tao, 2025

For  $k \geq 4$  we have  $F_{k,2}(n) \leq (1 - c + o(1))n$ .

## Granville, Soundararajan, 2001

For  $\ell \geq 2$  we have  $(1 - c_0 + o(1))n \leq F_{2\ell+1,2}(n)$ , where  $c_0 = 1 - \log(1 + \sqrt{e}) + 2 \int_1^{\sqrt{e}} \frac{\log t}{t+1} dt \approx 0.1715$  is the Hall-Montgomery constant.

## Conjecture (Verstraëte, 2006)

Let  $f \in \mathbb{Z}[x]$  and let  $k$  be a positive integer. Then, for some constant  $\rho = \rho(k, f)$  depending only on  $k$  and  $f$ , the maximal size of a set  $A \subseteq [n]$  such that no product of  $k$  distinct elements of  $A$  is in the value set of  $f$  is either  $(\rho + o(1))n$  or  $(\rho + o(1))\pi(n)$  as  $n \rightarrow \infty$ .

# Product representation of perfect powers

$F_{k,d}(n)$ : maximal size of a set  $A \subseteq [n]$  such that

$$a_1 a_2 \dots a_k = x^d, \quad a_1 < a_2 < \dots < a_k$$

has no solution with  $a_1, a_2, \dots, a_k \in A$  and integer  $x$ .

$f_{k,d}(n)$ : maximal size of a set  $A \subseteq [n]$  such that

$$a_1 a_2 \dots a_k = x^d$$

has no solution with  $a_1, a_2, \dots, a_k \in A$  and integer  $x$  except trivial solutions where the multiset  $\{a_1, a_2, \dots, a_k\}$  can be partitioned into  $d$ -tuples where each  $d$ -tuple consists of  $d$  copies of the same number. (E.g.  $a^3 b^3 c^3 = x^3$  is a trivial solution, but  $a^5 b^2 c^2 = x^3$  is not.) Note that  $f_{k,d}(n) \leq F_{k,d}(n)$ .

# Product representation of perfect powers

$F_{k,d}(n)$ : maximal size of a set  $A \subseteq [n]$  such that

$$a_1 a_2 \dots a_k = x^d, \quad a_1 < a_2 < \dots < a_k$$

has no solution with  $a_1, a_2, \dots, a_k \in A$  and integer  $x$ .

$f_{k,d}(n)$ : maximal size of a set  $A \subseteq [n]$  such that

$$a_1 a_2 \dots a_k = x^d$$

has no solution with  $a_1, a_2, \dots, a_k \in A$  and integer  $x$  except trivial solutions where the multiset  $\{a_1, a_2, \dots, a_k\}$  can be partitioned into  $d$ -tuples where each  $d$ -tuple consists of  $d$  copies of the same number. (E.g.  $a^3 b^3 c^3 = x^3$  is a trivial solution, but  $a^5 b^2 c^2 = x^3$  is not.) Note that  $f_{k,d}(n) \leq F_{k,d}(n)$ .



# Product representation of perfect cubes

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

# Product representation of perfect cubes

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

# Product representation of perfect cubes

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

# Product representation of perfect cubes

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

# Product representation of perfect cubes

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{5/6}$

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$



## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left( \pi(n) + \pi\left(\frac{n}{2}\right) \right) \ll n^{5/6}$

## Our results

- $c_1 n^{2/3} < n - F_{2,3}(n) \leq n - f_{2,3}(n) < c_2 n^{2/3}$
- $f_{3,3}(n) = (c_{3,3} + o(1))n$ , where  $0.6224 \leq c_{3,3} \leq 0.6420$
- $F_{3,3}(n) = (C_{3,3} + o(1))n$ , where  $0.6919 \leq C_{3,3} \leq 0.7136$
- $\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{\varepsilon \log 3}{2\sqrt{3}} - \varepsilon}}$
- For every  $d \geq 2$ ,  $d \nmid k$ , we have  $\frac{n}{(\log n)^d} \ll n - F_{k,d}(n)$ .
- $c_1 \frac{n^{3/4}}{(\log n)^{3/2}} < f_{6,3}(n) - \pi(n) < c_2 \frac{n^{3/4}}{(\log n)^{3/2}}$
- $F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$
- $\frac{n^{2/3}}{(\log n)^{4/3}} \ll f_{9,3}(n) - \pi(n) \ll n^{2/3} \log n$
- $\frac{n^{5/6}}{(\log n)^{5/3}} < F_{9,3}(n) - \left(\pi(n) + \pi\left(\frac{n}{2}\right)\right) \ll n^{5/6}$

| $k$ and $d$       | growth rate of $F_{k,d}(n)$           |
|-------------------|---------------------------------------|
| $d \mid k, d < k$ | $o(n)$                                |
| $k < d$           | $(1 - o(1))n$                         |
| $k = d$           | $(c_d + o(1))n$ with $c_d \in (0, 1)$ |
| $k = d + 1$       | $(1 - o(1))n$                         |
| $k \geq d + 2$    | $< (1 - c)n$ with $c > 0$             |

**Table:** Summary of results of Tao and FJKPS.

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Lower bound:  $G : C_4$ -free graph on  $\{\text{primes} \leq \sqrt{n}\}$

$$A = \{uv : uv \text{ is an edge of } G\} \cup \{\text{primes} \in (\sqrt{n}, n]\}$$

$$a_1 a_2 = b_1 b_2 \ (a_1, a_2, b_1, b_2) \implies \{a_1, a_2\} = \{b_1, b_2\}$$

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Lower bound:  $G : C_4$ -free graph on  $\{\text{primes} \leq \sqrt{n}\}$

$$A = \{uv : uv \text{ is an edge of } G\} \cup \{\text{primes} \in (\sqrt{n}, n]\}$$

$$a_1 a_2 = b_1 b_2 \ (a_1, a_2, b_1, b_2) \implies \{a_1, a_2\} = \{b_1, b_2\}$$

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Lower bound:  $G : C_4$ -free graph on  $\{\text{primes} \leq \sqrt{n}\}$

$$A = \{uv : uv \text{ is an edge of } G\} \cup \{\text{primes} \in (\sqrt{n}, n]\}$$

$$a_1 a_2 = b_1 b_2 \ (a_1, a_2, b_1, b_2) \implies \{a_1, a_2\} = \{b_1, b_2\}$$

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Lower bound:  $G : C_4$ -free graph on  $\{\text{primes} \leq \sqrt{n}\}$

$$A = \{uv : uv \text{ is an edge of } G\} \cup \{\text{primes} \in (\sqrt{n}, n]\}$$

$$a_1 a_2 = b_1 b_2 \ (a_1, a_2, b_1, b_2) \implies \{a_1, a_2\} = \{b_1, b_2\}$$

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Upper bound: write each  $a \in A$  as  $a = u_a v_a$ , where  $u_a \leq v_a$  and  $v_a$  is either a prime or  $v_a \leq n^{2/3}$ .

Observe that the graph  $G$  with vertex set  $[n^{2/3}] \cup \{\text{primes} \leq n\}$  and edge set  $E = \{u_a v_a : a \in A\}$  is  $C_4$ -free.



$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Upper bound: write each  $a \in A$  as  $a = u_a v_a$ , where  $u_a \leq v_a$  and  $v_a$  is either a prime or  $v_a \leq n^{2/3}$ .

Observe that the graph  $G$  with vertex set  $[n^{2/3}] \cup \{\text{primes} \leq n\}$  and edge set  $E = \{u_a v_a : a \in A\}$  is  $C_4$ -free.

$$F_{4,2}(n) \leq G_2(n) =$$

size of the largest multiplicative Sidon subset of  $[n]$

$$\text{Erdős (1934): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + c_2 n^{3/4}$$

$$\text{Erdős (1969): } \pi(n) + \frac{c_1 n^{3/4}}{(\log n)^{3/2}} \leq G_2(n) \leq \pi(n) + \frac{c_2 n^{3/4}}{(\log n)^{3/2}}$$

Upper bound: write each  $a \in A$  as  $a = u_a v_a$ , where  $u_a \leq v_a$  and  $v_a$  is either a prime or  $v_a \leq n^{2/3}$ .

Observe that the graph  $G$  with vertex set  $[n^{2/3}] \cup \{\text{primes} \leq n\}$  and edge set  $E = \{u_a v_a : a \in A\}$  is  $C_4$ -free.

## Theorem

$$F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$$

Construction:

$$A = \left\{ m : m = pq, \frac{n}{\log n} < m \leq n, p, q \text{ primes}, p < \frac{q}{\log n} \right\}.$$

Upper bound: by a result of Erdős, there exist distinct  $a_1, a_2, \dots, a_6 \in A$  such that

$$a_1 a_2 = a_3 a_4 = a_5 a_6,$$

however, then  $a_1 a_2 a_3 a_4 a_5 a_6$  is a perfect cube.

The result refutes the conjecture of Verstraëte.

## Theorem

$$F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$$

Construction:

$$A = \left\{ m : m = pq, \frac{n}{\log n} < m \leq n, p, q \text{ primes}, p < \frac{q}{\log n} \right\}.$$

Upper bound: by a result of Erdős, there exist distinct  $a_1, a_2, \dots, a_6 \in A$  such that

$$a_1 a_2 = a_3 a_4 = a_5 a_6,$$

however, then  $a_1 a_2 a_3 a_4 a_5 a_6$  is a perfect cube.

The result refutes the conjecture of Verstraëte.

## Theorem

$$F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$$

Construction:

$$A = \left\{ m : m = pq, \frac{n}{\log n} < m \leq n, p, q \text{ primes}, p < \frac{q}{\log n} \right\}.$$

Upper bound: by a result of Erdős, there exist distinct  $a_1, a_2, \dots, a_6 \in A$  such that

$$a_1 a_2 = a_3 a_4 = a_5 a_6,$$

however, then  $a_1 a_2 a_3 a_4 a_5 a_6$  is a perfect cube.

The result refutes the conjecture of Verstraëte.

## Theorem

$$F_{6,3}(n) = (1 + o(1)) \frac{n \log \log n}{\log n}$$

Construction:

$$A = \left\{ m : m = pq, \frac{n}{\log n} < m \leq n, p, q \text{ primes}, p < \frac{q}{\log n} \right\}.$$

Upper bound: by a result of Erdős, there exist distinct  $a_1, a_2, \dots, a_6 \in A$  such that

$$a_1 a_2 = a_3 a_4 = a_5 a_6,$$

however, then  $a_1 a_2 a_3 a_4 a_5 a_6$  is a perfect cube.

The result refutes the conjecture of Verstraëte.

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$\Omega(d) \leq \frac{1}{3} \log \log n$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$\Omega(d) \leq \frac{1}{3} \log \log n$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .



## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$\Omega(d) \leq \frac{1}{3} \log \log n$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$$\Omega(d) \leq \frac{1}{3} \log \log n$$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$\Omega(d) \leq \frac{1}{3} \log \log n$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (lower bound):

$A \subseteq \{1, 2, \dots, n\}$ :  $a_1 a_2 a_3 a_4 \neq x^3$  if  $a_i \in A$ ,  $a_1 < a_2 < a_3 < a_4$

$D := \{d_1, \dots, d_t\}$ , the set of those  $d$  such that  $d \leq n^{1/3}$  and

$\Omega(d) \leq \frac{1}{3} \log \log n$

By calculation,

$$t = |D| > \frac{n^{1/3}}{(\log n)^{1 + \frac{1}{3} \log \frac{1}{3} - \frac{1}{3} + \frac{\varepsilon}{3}}}.$$

$H$ : 3-uniform hypergraph on  $\{P_1, \dots, P_t\}$  such that  $\{P_i, P_j, P_k\}$  is an edge in  $H$  if and only if  $d_i d_j d_k \in A$ .

$M$ : the set of those  $m \in [n]$  such that  $m \notin A$  and  $m = d_i d_j d_k$  for some  $1 \leq i < j < k \leq t$ , then  $|A| \leq n - |M|$ .

$h(m) := \#$  of triples  $(d_i, d_j, d_k)$  such that  $m = d_i d_j d_k$ ,  
 $1 \leq i < j < k \leq t$ .

If  $m = p_1^{k_1} p_2^{k_2} \cdots p_r^{k_r} \in M$ , then

$$\Omega(m) = \Omega(d_i) + \Omega(d_j) + \Omega(d_k) \leq \log \log n,$$

hence

$$h(m) \leq \prod_{i=1}^r \binom{k_i + 2}{2} \leq \prod_{i=1}^r 3^{k_i} \leq 3^{\log \log n} = (\log n)^{\log 3}.$$

If  $H$  contains a  $K_4^3$ , then for some  $d_{i_1} < d_{i_2} < d_{i_3} < d_{i_4}$  and

$$a_1 = d_{i_1} d_{i_2} d_{i_3}, \quad a_2 = d_{i_1} d_{i_2} d_{i_4}, \quad a_3 = d_{i_1} d_{i_3} d_{i_4}, \quad a_4 = d_{i_2} d_{i_3} d_{i_4}$$

we have  $a_1 a_2 a_3 a_4 = (d_{i_1} d_{i_2} d_{i_3} d_{i_4})^3$ .

Therefore,  $H$  does not contain any  $K_4^3$ .

$h(m) := \#$  of triples  $(d_i, d_j, d_k)$  such that  $m = d_i d_j d_k$ ,

$1 \leq i < j < k \leq t$ .

If  $m = p_1^{k_1} p_2^{k_2} \cdots p_r^{k_r} \in M$ , then

$$\Omega(m) = \Omega(d_i) + \Omega(d_j) + \Omega(d_k) \leq \log \log n,$$

hence

$$h(m) \leq \prod_{i=1}^r \binom{k_i + 2}{2} \leq \prod_{i=1}^r 3^{k_i} \leq 3^{\log \log n} = (\log n)^{\log 3}.$$

If  $H$  contains a  $K_4^3$ , then for some  $d_{i_1} < d_{i_2} < d_{i_3} < d_{i_4}$  and

$$a_1 = d_{i_1} d_{i_2} d_{i_3}, \quad a_2 = d_{i_1} d_{i_2} d_{i_4}, \quad a_3 = d_{i_1} d_{i_3} d_{i_4}, \quad a_4 = d_{i_2} d_{i_3} d_{i_4}$$

we have  $a_1 a_2 a_3 a_4 = (d_{i_1} d_{i_2} d_{i_3} d_{i_4})^3$ .

Therefore,  $H$  does not contain any  $K_4^3$ .

$h(m) := \#$  of triples  $(d_i, d_j, d_k)$  such that  $m = d_i d_j d_k$ ,

$1 \leq i < j < k \leq t$ .

If  $m = p_1^{k_1} p_2^{k_2} \cdots p_r^{k_r} \in M$ , then

$$\Omega(m) = \Omega(d_i) + \Omega(d_j) + \Omega(d_k) \leq \log \log n,$$

hence

$$h(m) \leq \prod_{i=1}^r \binom{k_i + 2}{2} \leq \prod_{i=1}^r 3^{k_i} \leq 3^{\log \log n} = (\log n)^{\log 3}.$$

If  $H$  contains a  $K_4^3$ , then for some  $d_{i_1} < d_{i_2} < d_{i_3} < d_{i_4}$  and

$$a_1 = d_{i_1} d_{i_2} d_{i_3}, \quad a_2 = d_{i_1} d_{i_2} d_{i_4}, \quad a_3 = d_{i_1} d_{i_3} d_{i_4}, \quad a_4 = d_{i_2} d_{i_3} d_{i_4}$$

we have  $a_1 a_2 a_3 a_4 = (d_{i_1} d_{i_2} d_{i_3} d_{i_4})^3$ .

Therefore,  $H$  does not contain any  $K_4^3$ .

$h(m) := \#$  of triples  $(d_i, d_j, d_k)$  such that  $m = d_i d_j d_k$ ,

$1 \leq i < j < k \leq t$ .

If  $m = p_1^{k_1} p_2^{k_2} \cdots p_r^{k_r} \in M$ , then

$$\Omega(m) = \Omega(d_i) + \Omega(d_j) + \Omega(d_k) \leq \log \log n,$$

hence

$$h(m) \leq \prod_{i=1}^r \binom{k_i + 2}{2} \leq \prod_{i=1}^r 3^{k_i} \leq 3^{\log \log n} = (\log n)^{\log 3}.$$

If  $H$  contains a  $K_4^3$ , then for some  $d_{i_1} < d_{i_2} < d_{i_3} < d_{i_4}$  and

$$a_1 = d_{i_1} d_{i_2} d_{i_3}, \quad a_2 = d_{i_1} d_{i_2} d_{i_4}, \quad a_3 = d_{i_1} d_{i_3} d_{i_4}, \quad a_4 = d_{i_2} d_{i_3} d_{i_4}$$

we have  $a_1 a_2 a_3 a_4 = (d_{i_1} d_{i_2} d_{i_3} d_{i_4})^3$ .

Therefore,  $H$  does not contain any  $K_4^3$ .



By a result of de Caen  $\exists \delta > 0$  such that there at least  $\delta t^3$  triples  $(i, j, k)$  such that  $\{P_i, P_j, P_k\}$  is not an edge in  $H$ .

Let  $h = \max_{m \in M} h(m) \leq (\log n)^{\log 3}$ .

Therefore,

$$|M| \geq \frac{\delta t^3}{h} \gg \frac{n}{(\log n)^{3+\log \frac{1}{3}-1+\varepsilon} \cdot (\log n)^{\log 3}} = \frac{n}{(\log n)^{2+\varepsilon}},$$

which completes the proof of the lower bound.

By a result of de Caen  $\exists \delta > 0$  such that there at least  $\delta t^3$  triples  $(i, j, k)$  such that  $\{P_i, P_j, P_k\}$  is not an edge in  $H$ .

Let  $h = \max_{m \in M} h(m) \leq (\log n)^{\log 3}$ .

Therefore,

$$|M| \geq \frac{\delta t^3}{h} \gg \frac{n}{(\log n)^{3 + \log \frac{1}{3} - 1 + \epsilon} \cdot (\log n)^{\log 3}} = \frac{n}{(\log n)^{2 + \epsilon}},$$

which completes the proof of the lower bound.

By a result of de Caen  $\exists \delta > 0$  such that there at least  $\delta t^3$  triples  $(i, j, k)$  such that  $\{P_i, P_j, P_k\}$  is not an edge in  $H$ .

Let  $h = \max_{m \in M} h(m) \leq (\log n)^{\log 3}$ .

Therefore,

$$|M| \geq \frac{\delta t^3}{h} \gg \frac{n}{(\log n)^{3 + \log \frac{1}{3} - 1 + \varepsilon} \cdot (\log n)^{\log 3}} = \frac{n}{(\log n)^{2 + \varepsilon}},$$

which completes the proof of the lower bound.

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (upper bound):

Construction:  $A$  is the set of the integers  $a$  such that

- (i)  $\frac{n}{\log n} \leq a \leq n$ ,
- (ii)  $d^2 \mid a$  implies  $d \leq \log n$ , and
- (iii)  $a$  cannot be written in the form  $a = uvw$  with integers  $u, v, w$  such that  $\frac{\sqrt[3]{n}}{(\log n)^{16}} \leq u, v, w \leq \sqrt[3]{n}(\log n)^{16}$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (upper bound):

Construction:  $A$  is the set of the integers  $a$  such that

- (i)  $\frac{n}{\log n} \leq a \leq n$ ,
- (ii)  $d^2 \mid a$  implies  $d \leq \log n$ , and
- (iii)  $a$  cannot be written in the form  $a = uvw$  with integers  $u, v, w$  such that  $\frac{\sqrt[3]{n}}{(\log n)^{16}} \leq u, v, w \leq \sqrt[3]{n}(\log n)^{16}$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (upper bound):

Construction:  $A$  is the set of the integers  $a$  such that

- (i)  $\frac{n}{\log n} \leq a \leq n$ ,
- (ii)  $d^2 \mid a$  implies  $d \leq \log n$ , and
- (iii)  $a$  cannot be written in the form  $a = uvw$  with integers  $u, v, w$  such that  $\frac{\sqrt[3]{n}}{(\log n)^{16}} \leq u, v, w \leq \sqrt[3]{n}(\log n)^{16}$ .

## Theorem

$$\frac{n}{(\log n)^{2+\varepsilon}} < n - F_{4,3}(n) \leq n - f_{4,3}(n) < \frac{n}{(\log n)^{1 - \frac{e \log 3}{2\sqrt{3}} - \varepsilon}}$$

Proof (upper bound):

Construction:  $A$  is the set of the integers  $a$  such that

- (i)  $\frac{n}{\log n} \leq a \leq n$ ,
- (ii)  $d^2 \mid a$  implies  $d \leq \log n$ , and
- (iii)  $a$  cannot be written in the form  $a = uvw$  with integers  $u, v, w$  such that  $\frac{\sqrt[3]{n}}{(\log n)^{16}} \leq u, v, w \leq \sqrt[3]{n}(\log n)^{16}$ .

Let  $1 < k < d$ . Is it true that

$$n^{k/d} \ll n - F_{k,d}(n) \leq n - f_{k,d}(n) \ll n^{k/d}?$$

Is it true that there exists a constant  $c$  such that

$$f_{2,3}(n) = n - (c + o(1))n^{2/3}?$$

Is it true that for any  $d \geq 4$  and  $k > d$ ,  $d \mid k$ , there exist constants  $c_{k,d} > 0$  and  $C_{k,d} \in \mathbb{Z}^+$  such that

$$F_{k,d}(n) = (c_{k,d} + o(1))\pi_{C_{k,d}}(n)?$$

Thank you for your attention!



Let  $1 < k < d$ . Is it true that

$$n^{k/d} \ll n - F_{k,d}(n) \leq n - f_{k,d}(n) \ll n^{k/d}?$$

Is it true that there exists a constant  $c$  such that

$$f_{2,3}(n) = n - (c + o(1))n^{2/3}?$$

Is it true that for any  $d \geq 4$  and  $k > d$ ,  $d \mid k$ , there exist constants  $c_{k,d} > 0$  and  $C_{k,d} \in \mathbb{Z}^+$  such that

$$F_{k,d}(n) = (c_{k,d} + o(1))\pi_{C_{k,d}}(n)?$$

Thank you for your attention!

Let  $1 < k < d$ . Is it true that

$$n^{k/d} \ll n - F_{k,d}(n) \leq n - f_{k,d}(n) \ll n^{k/d}?$$

Is it true that there exists a constant  $c$  such that

$$f_{2,3}(n) = n - (c + o(1))n^{2/3}?$$

Is it true that for any  $d \geq 4$  and  $k > d$ ,  $d \mid k$ , there exist constants  $c_{k,d} > 0$  and  $C_{k,d} \in \mathbb{Z}^+$  such that

$$F_{k,d}(n) = (c_{k,d} + o(1))\pi_{C_{k,d}}(n)?$$

Thank you for your attention!

Let  $1 < k < d$ . Is it true that

$$n^{k/d} \ll n - F_{k,d}(n) \leq n - f_{k,d}(n) \ll n^{k/d}?$$

Is it true that there exists a constant  $c$  such that

$$f_{2,3}(n) = n - (c + o(1))n^{2/3}?$$

Is it true that for any  $d \geq 4$  and  $k > d$ ,  $d \mid k$ , there exist constants  $c_{k,d} > 0$  and  $C_{k,d} \in \mathbb{Z}^+$  such that

$$F_{k,d}(n) = (c_{k,d} + o(1))\pi_{C_{k,d}}(n)?$$

Thank you for your attention!

Let  $1 < k < d$ . Is it true that

$$n^{k/d} \ll n - F_{k,d}(n) \leq n - f_{k,d}(n) \ll n^{k/d}?$$

Is it true that there exists a constant  $c$  such that

$$f_{2,3}(n) = n - (c + o(1))n^{2/3}?$$

Is it true that for any  $d \geq 4$  and  $k > d$ ,  $d \mid k$ , there exist constants  $c_{k,d} > 0$  and  $C_{k,d} \in \mathbb{Z}^+$  such that

$$F_{k,d}(n) = (c_{k,d} + o(1))\pi_{C_{k,d}}(n)?$$

Thank you for your attention!