Nordic Laboratories

PATIENT: XXXXXXXXXXXXXXXXXXX			TEST REF: TST-NL-XXXXX
TEST NUMBER: T-NL-XXXXXX		COLLECTED: 2025-XX-XX	PRACTITIONER: XXXXXXXXXXXXXXXXXXX
GENDER: XXXXX			XXXXXXXXXXXXXXXXXX
AGE: XX			

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report


Microbiome Summary Score

Your gut microbiome score evaluates all the relevant metrics for your age range into a single number, giving you a clear picture of how your microbiome stacks up. The score ranges from 0 to 100. A score closer to 100 suggests your microbiome is in excellent shape compared to the reference population. Conversely, a score closer to 0 indicates more areas for improvement.

We calculate this score by looking at various factors in your gut/vaginal microbiome, such as the balance of beneficial bacteria versus opportunistic pathogens. We then compare these factors to a wide range of individuals. Our dataset includes publicly available reference samples and proprietary data from Tiny Health customers.

To determine your percentile, we compare your Microbiome Summary Score to others in the same age range to help you understand what is typical and where your sample falls. For example, if you are in the 41st percentile, 41% of comparable samples have a lower Microbiome Summary Score and 59% have a higher score.

90 Microbiome summary score



Sample at a glance

1 Needs support

0 Needs improvement

Bacterial Vaginosis-Associated Bacteria

- Gardnerella vaginalis 6.915%

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Results Summary

	Metrics	Clinical indication examples
Beneficial microbes	For Your Overall Vaginal Health Common Lactobacillus Species	• Oral probiotic support • Vaginal suppository support • Hygiene suggestions (e.g., product use, no douching) • Alternatives to antimicrobials
Disruptive microbes	Bacterial Vaginosis-Associated Bacteria Aerobic Bacteria Candida Yeasts Sexually Transmitted Microbes Ureaplasma Group B Strep	• Dietary, supplement, and nutritional support • Close monitoring post-antibiotics to track recovery • Vaginal oral probiotic support • Vaginal suppository support • Products (e.g., Vitamin C tablets, boric acid suppositories, tea tree oil suppositories) • Review symptoms/conditions • Consider additional confirmatory testing (e.g., BV, culture, STI)

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

Nordic Laboratories

PATIENT: XXXXXXXXXXXXXXXXXXX			TEST REF: TST-NL-XXXXX
TEST NUMBER: T-NL-XXXXXX	COLLECTED: 2025-XX-XX		PRACTITIONER: XXXXXXXXXXXXXXXXXXXX
GENDER: XXXXX			XXXXXXXXXXXXXXXXXX
AGE: XX			

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Detailed Vaginal Health Test Results

Legend: (p) Phylum (f) Family (g) Genus (sp) Species

Beneficial microbes

For Your Overall Vaginal Health

Lactobacillus (g)	90.430%	90% / 60%

Common Lactobacillus Species

Lactobacillus crispatus (sp)	90.303%
Lactobacillus gasseri (sp)	0.000%
Lactobacillus iners (sp)	0.000%
Lactobacillus jensenii (sp)	0.000%
Lactobacillus paragasseri (sp)	0.000%
Lactobacillus delbrueckii (sp)	0.000%
Lactobacillus mulieris (sp)	0.000%
Lacticaseibacillus paracasei (sp)	0.000%
Lactobacillus acidophilus (sp)	0.000%
Lacticaseibacillus rhamnosus (sp)	0.000%
Lactobacillus johnsonii (sp)	0.000%
Lactobacillus helveticus (sp)	0.231%

Disruptive microbes

Bacterial Vaginosis-Associated Bacteria

Gardnerella vaginalis (sp)	6.915%	6%
Fannyhessea vaginae (sp)	0.000%	0.005%
28L (Megasphaera) (g)	0.000%	0.005%
UBA629 sp005465875 (sp)	0.000%	0.005%
Prevotella (g)	0.144%	4%

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

Nordic Laboratories

PATIENT: XXXXXXXXXXXXXXXXXXX
TEST NUMBER: T-NL-XXXXXX
GENDER: XXXXX
AGE: XX
COLLECTED: 2025-XX-XX
TEST REF: TST-NL-XXXXX
PRACTITIONER:
XXXXXXXXXXXXXXXXXXX
XXXXXXXXXXXXXXXXX

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Disruptive microbes

Aerobic Bacteria		
Total aerobic bacteria	1.108%	4%
Candida Yeasts		
Candida albicans (sp)	0.000%	0.005%
Candida dubliniensis (sp)	0.000%	0.005%
Candida parapsilosis (sp)	0.000%	0.005%
Candida tropicalis (sp)	0.000%	0.005%
Sexually Transmitted Microbes		
Trichomonas vaginalis (sp)	0.000%	0.005%
Chlamydia trachomatis (sp)	0.000%	0.005%
Neisseria gonorrhoeae (sp)	0.000%	0.005%
Mycoplasmoides genitalium (sp)	0.000%	0.005%
Treponema pallidum (sp)	0.000%	0.005%
Ureaplasma		
Ureaplasma urealyticum (sp)	0.000%	0.005%
Ureaplasma parvum (sp)	0.000%	0.3%, 0.005%
Group B Strep		
Streptococcus agalactiae (GBS) (sp)	0.000%	0.005%

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Microbiome Breakdown

- **Protective:** These support microbiome health and overall health.
- **Disruptive:** These microbes can cause issues and some are linked with symptoms and disease states. This should make up a minority of the microbiome, kept in check by protective and variable microbes. Any Disruptive microbes out of range will be flagged in the "Detailed Vaginal Health Test Results" section above.
- **Variable:** These can be protective or disruptive depending on their levels and what other microbes are around them. Lower levels are usually ok, and higher levels can cause issues.
- **Unknown:** These are lesser known microbes. Not enough is known to classify them as protective, disruptive or variable at this time.

Your breakdown



Protective 91% Variable 0% Disruptive 7% Unknown 1%

Top 20 species

Metric	Result	Description
Lactobacillus crispatus	90.303%	*L. crispatus* is your vagina's best friend. It is an expert producer of lactic acid, which keeps the vaginal pH low and ensures a protective environment. This beneficial Lactobacillus is very good at: • Preventing the growth of harmful microbes that cause vaginal infections. • Protecting you against HIV, herpes simplex virus type 2, and human papillomavirus. • Reducing the risk of preterm delivery. Sometimes though, *L. crispatus* can become too cozy and cause cytolytic vaginosis.
Bifidobacterium vaginale_E	3.309%	*G. vaginalis* are often found in healthy women in low amounts. When their levels go up, it's worth keeping an eye on. One of their main skills is the production of biofilm, a sticky layer of infectious bacteria that serves as a fort for other harmful buddies such as *Fannyhessea* and *Prevotella*. So when *G. vaginalis* are found along with other harmful bacteria, chances are you may have bacterial vaginosis (BV). In the context of BV, *G. vaginalis* have been associated with pregnancy complications and increased risk of sexually transmitted infections.

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Bifidobacterium vaginale_C	3.253%	*G. vaginalis* are often found in healthy women in low amounts. When their levels go up, it's worth keeping an eye on. One of their main skills is the production of biofilm, a sticky layer of infectious bacteria that serves as a fort for other harmful buddies such as *Fannyhessea* and *Prevotella*. So when *G. vaginalis* are found along with other harmful bacteria, chances are you may have bacterial vaginosis (BV). In the context of BV, *G. vaginalis* have been associated with pregnancy complications and increased risk of sexually transmitted infections.
Lactobacillus amylovorus	0.493%	*L. amylovorus* has been found in healthy vaginal microbiomes. Even if it's not very common, it is probably associated with good vaginal health. That's because all lactobacilli produce lactic acid, to a greater or lesser extent. This lowers the vaginal pH, creating an acidic environment that is not favorable for disruptive microbes that cause infection.
Limosilactobacillus vaginalis_A	0.344%	*L. vaginalis* has been found in healthy vaginal microbiomes. Laboratory experiments show that this beneficial bacterium has antifungal properties that may help keep *Candida* yeasts at bay. But *L. vaginalis* likes to tease immune cells, which may promote inflammation. So it may be that this species is not as protective as others.
Bifidobacterium vaginale_G	0.274%	*G. vaginalis* are often found in healthy women in low amounts. When their levels go up, it's worth keeping an eye on. One of their main skills is the production of biofilm, a sticky layer of infectious bacteria that serves as a fort for other harmful buddies such as *Fannyhessea* and *Prevotella*. So when *G. vaginalis* are found along with other harmful bacteria, chances are you may have bacterial vaginosis (BV). In the context of BV, *G. vaginalis* have been associated with pregnancy complications and increased risk of sexually transmitted infections.
Lactobacillus helveticus	0.231%	*L. helveticus* has been found in healthy vaginal microbiomes. This beneficial bacterium can inhibit the growth of disruptive bacteria that may cause urinary tract infection or bacterial vaginosis, such as *Escherichia coli*, *Gardnerella vaginalis*, and *Prevotella bivia*. It may also protect against yeast infection.
Finegoldia magna_H	0.198%	*Finegoldia* species are normal inhabitants of the skin and mucous membranes. On occasion, they can cause trouble. Not much is known about their role in the vagina. They are more often found in women whose vaginal community is not dominated by *Lactobacillus*.
Corynebacterium minutissimum_A	0.175%	
Streptococcus anginosus	0.165%	*S. anginosus* is a common member of the mouth, vagina, urinary tract, and gut microbiome. But under certain conditions it can cause infection. It can, for example, cause aerobic vaginitis (AV). Women with AV have higher abundance of *S. anginosus* than healthy women or women with bacterial vaginosis. It has also been associated with urinary tract infections and urinary incontinence, or poor bladder control. *S. anginosus* might be resistant to some common antibiotics, such as erythromycin and clindamycin.

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

Corynebacterium tuscaniense	0.159%	
Corynebacterium coyleae	0.129%	
Phocaeicola vulgatus	0.119%	We don't know much about the role of *P. vulgatus* in the vagina. This bacterium was previously known as *Bacteroides vulgatus*. And we know that *Bacteroides* species, in low numbers, can be found in healthy vaginal microbiomes. However, they are more frequently detected in women with low numbers of lactobacilli. So they are often detected in women with bacterial vaginosis
Peptoniphilus_A harei_A	0.116%	
Pseudoglutamicibacter albus_A	0.086%	
Bifidobacterium vaginale	0.079%	*G. vaginalis* are often found in healthy women in low amounts. When their levels go up, it's worth keeping an eye on. One of their main skills is the production of biofilm, a sticky layer of infectious bacteria that serves as a fort for other harmful buddies such as *Fannyhessea* and *Prevotella*. So when *G. vaginalis* are found along with other harmful bacteria, chances are you may have bacterial vaginosis (BV). In the context of BV, *G. vaginalis* have been associated with pregnancy complications and increased risk of sexually transmitted infections.
Corynebacterium riegelii	0.076%	
Helcococcus sueciensis	0.063%	
Lactobacillus kitasatonis	0.056%	*L. kitasatonis* has been found in healthy vaginal microbiomes. Even if it's not very common, it is probably associated with good vaginal health. That's because all lactobacilli produce lactic acid, to a greater or lesser extent. This lowers the vaginal pH, creating an acidic environment that is not favorable for disruptive microbes that cause infection.
Fastidiosipila sanguinis	0.053%	

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

All species

Corynebacterium kefirresidentii	0.053%	Bifidobacterium breve	0.050%
Corynebacterium sp001767255	0.046%	Bifidobacterium bifidum	0.046%
Bifidobacterium sp002884815	0.046%	Corynebacterium phoceense	0.040%

Our high confidence detection limit is 0.01% abundance. Very low abundance organisms can be important members of the microbiome but also very difficult to accurately detect. Detections below 0.01% abundance have an increased risk of being false positives. However, in our testing, we have observed that most false positive identifications are closely related to a detected species.

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: [17 Weeks Vaginal] Tiny Health Sample Report

References

[1] A. Uzan-Yulzari et al., "Neonatal antibiotic exposure impairs child growth during the first six years of life by perturbing intestinal microbial colonization," Nat Commun, vol. 12, no. 1, p. 443, 2021, doi: 10.1038/s41467-020-20495-4.
[2] K. E. Fujimura et al., "Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation," Nat Med, vol. 22, no. 10, pp. 1187–1191, 2016, doi: 10.1038/nm.4176.
[3] M. B. Azad et al., "Infant gut microbiota and food sensitization: associations in the first year of life," Clin Exp Allergy, vol. 45, no. 3, pp. 632–643, 2015, doi: 10.1111/cea.12487.
[4] H. M. Tun et al., "Roles of Birth Mode and Infant Gut Microbiota in Intergenerational Transmission of Overweight and Obesity From Mother to Offspring," Jama Pediatr, vol. 172, no. 4, p. 368, 2018, doi: 10.1001/jamapediatrics.2017.5535.
[5] T. Vatanen et al., "The human gut microbiome in early-onset type 1 diabetes from the TEDDY study," Nature, vol. 562, no. 7728, pp. 589–594, 2018, doi: 10.1038/s41586-018-0620-2.
[6] C. J. Stewart et al., "Temporal development of the gut microbiome in early childhood from the TEDDY study," Nature, vol. 562, no. 7728, pp. 583–588, 2018, doi: 10.1038/s41586-018-0617-x.
[7] S. Tamburini, N. Shen, H. C. Wu, and J. C. Clemente, "The microbiome in early life: implications for health outcomes," Nat Med, vol. 22, no. 7, pp. 713–722, 2016, doi: 10.1038/nm.4142.
[8] M. J. Ege, "The Hygiene Hypothesis in the Age of the Microbiome," Ann Am Thorac Soc, vol. 14, no. Supplement_5, pp. S348–S353, 2017, doi: 10.1513/annalsats.201702-139aw.
[9] B. M. Henrick et al., "Bifidobacteria-mediated immune system imprinting early in life," Cell, 2021, doi: 10.1016/j.cell.2021.05.030.
[10] A. Marcobal et al., "Bacteroides in the Infant Gut Consume Milk Oligosaccharides via Mucus-Utilization Pathways," Cell Host Microbe, vol. 10, no. 5, pp. 507–514, 2011, doi: 10.1016/j.chom.2011.10.007.
[11] T. Wilmanski et al., "Gut microbiome pattern reflects healthy ageing and predicts survival in humans," Nat Metabolism, vol. 3, no. 2, pp. 274–286, 2021, doi: 10.1038/s42255-021-00348-0.
[12] M. A. Jackson et al., "Signatures of early frailty in the gut microbiota," Genome Med, vol. 8, no. 1, p. 8, 2016, doi: 10.1186/s13073-016-0262-7.[13] J. M. Pickard, M. Y. Zeng, R. Caruso, and G. Núñez, "Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease," Immunol Rev, vol. 279, no. 1, pp. 70–89, 2017, doi: 10.1111/imr.12567.
[13] J. M. Pickard, M. Y. Zeng, R. Caruso, and G. Núñez, "Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease," Immunol Rev, vol. 279, no. 1, pp. 70–89, 2017, doi: 10.1111/imr.12567.
[14] O. Manor et al., "Health and disease markers correlate with gut microbiome composition across thousands of people," Nat Commun, vol. 11, no. 1, p. 5206, 2020, doi: 10.1038/s41467-020-18871-1.
[15] M. Scudellari, "Cleaning up the hygiene hypothesis," Proc National Acad Sci, vol. 114, no. 7, pp. 1433–1436, 2017, doi: 10.1073/pnas.1700688114.[16] R. Debray, R. A. Herbert, A. L. Jaffe, A. Crits-Christoph, M. E. Power, and B. Koskella, "Priority effects in microbiome assembly," Nat Rev Microbiol, vol. 20, no. 2, pp. 109–121, 2022, doi: 10.1038/s41579-021-00604-w.
[16] R. Debray, R. A. Herbert, A. L. Jaffe, A. Crits-Christoph, M. E. Power, and B. Koskella, "Priority effects in microbiome assembly," Nat Rev Microbiol, vol. 20, no. 2, pp. 109–121, 2022, doi: 10.1038/s41579-021-00604-w.
[17] C. Huttenhower et al., "Structure, function and diversity of the healthy human microbiome," Nature, vol. 486, no. 7402, pp. 207–214, 2012, doi: 10.1038/nature11234.
[18] P. I. Costea et al., "Enterotypes in the landscape of gut microbial community composition," Nat Microbiol, vol. 3, no. 1, pp. 8–16, 2018, doi: 10.1038/s41564-017-0072-8.
[19] W. E. Anthony et al., "Acute and persistent effects of commonly used antibiotics on the gut microbiome and resistome in healthy adults," Cell Reports, vol. 39, no. 2, p. 110649, 2022, doi: 10.1016/j.celrep.2022.110649.
[20] A. Palleja et al., "Recovery of gut microbiota of healthy adults following antibiotic exposure," Nat Microbiol, vol. 3, no. 11, pp. 1255–1265, 2018, doi: 10.1038/s41564-018-0257-9.
[21] A. Bell et al., "Elucidation of a sialic acid metabolism pathway in mucus-foraging Ruminococcus gnavus unravels mechanisms of bacterial adaptation to the gut," Nat Microbiol, vol. 4, no. 12, pp. 2393–2404, 2019, doi: 10.1038/s41564-019-0590-7.
[22] M. T. Henke, D. J. Kenny, C. D. Cassilly, H. Vlamakis, R. J. Xavier, and J. Clardy, "Ruminococcus gnavus, a member of the human gut microbiome associated with Crohn's disease, produces an inflammatory polysaccharide," Proc National Acad Sci, vol. 116, no. 26, pp. 12672–12677, 2019, doi: 10.1073/pnas.1904099116.
[23] W. Li et al., "Ecological and network analyses identify four microbial species with potential significance for the diagnosis/treatment of ulcerative colitis (UC)," Bmc Microbiol, vol. 21, no. 1, p. 138, 2021, doi: 10.1186/s12866-021-02201-6.
[24] F. Blachier, M. Beaumont, and E. Kim, "Cysteine-derived hydrogen sulfide and gut health," Curr Opin Clin Nutr, vol. 22, no. 1, pp. 68–75, 2019, doi: 10.1097/mco.0000000000000526.
[25] S. C. Peck, K. Denger, A. Burrichter, S. M. Irwin, E. P. Balskus, and D. Schleheck, "A glycyl radical enzyme enables hydrogen sulfide production by the human intestinal bacterium Bilophila wadsworthia," Proc National Acad Sci, vol. 116, no. 8, pp. 3171–3176, 2019, doi: 10.1073/pnas.1815661116.
[26] H. M. HAMER, D. JONKERS, K. VENEMA, S. VANHOUTVIN, F. J. TROOST, and R. -J. BRUMMER, "Review article: the role of butyrate on colonic function," Aliment Pharm Therap, vol. 27, no. 2, pp. 104–119, 2008, doi: 10.1111/j.1365-2036.2007.03562.x.
[27] Y. Furusawa et al., "Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells," Nature, vol. 504, no. 7480, pp. 446–450, 2013, doi: 10.1038/nature12721.
[28] H. Sokol et al., "Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients," Proc National Acad Sci, vol. 105, no. 43, pp. 16731–16736, 2008, doi: 10.1073/pnas.0804812105.
[29] D. Ríos-Covián, P. Ruas-Madiedo, A. Margolles, M. Gueimonde, C. G. de los Reyes-Gavilán, and N. Salazar, "Intestinal Short Chain Fatty Acids and their Link with Diet and Human Health," Front Microbiol, vol. 7, p. 185, 2016, doi: 10.3389/fmicb.2016.00185.
[30] Q. Zhou et al., "Association Between Gut Akkermansia and Metabolic Syndrome is Dose-Dependent and Affected by Microbial Interactions: A Cross-Sectional Study," Diabetes Metabolic Syndrome Obes Targets Ther, vol. 14, pp. 2177–2188, 2021, doi: 10.2147/dmso.s311388.

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright **2025** Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.