

TID: 3×40 mg/day (120 mg/day total) × 30 days | F=0.6 (FDA label) | kₐ=1.5 h⁻¹ (Hammarlund 1984) | Ki=0.35 mg/L (calibrated) | CKD-5
A IS — 3×40mg/day oral (TID), 30 days
IS plasma (mg/L)
Time (days)
Baseline 19.712 mg/L
ODE 24h-avg = 20.899 mg/L (+6.0%)
B pCS — 3×40mg/day oral (TID), 30 days
pCS plasma (mg/L)
Time (days)
Baseline 24.059 mg/L
ODE 24h-avg = 26.351 mg/L (+9.5%)
C Furosemide — last 24h cycle (TID: 3×40mg at 0/8/16h)
Furosemide plasma (mg/L)
Time after day start (h)
dose
0h
8h
16h
Analytical Css,avg = 3.215 mg/L
ODE 24h-avg = 0.836 mg/L
D IS final 24h cycle — sampling time effect
IS plasma (mg/L)
Time after day start (h)
No-drug baseline (19.712 mg/L)
ODE 24h-avg (+6.0%)
Analytical (+29.3%)
+6%
+6%
+6%
+6%
@1h
@2h
@3h
@4h
E ODE (TID: 3×40mg) vs Analytical (live PK) vs Clinical — >120 mg/day
% change vs no-drug baseline
ODE 24h-avg (TID)
Analytical (exact MM, live PK)
Clinical (Costes-Albrespic 2025)
+6.0%
+29.3%
+31.9%
+9.5%
+37.5%
+29.3%
Indoxyl Sulphate (IS)
p-Cresyl Sulphate (pCS)