# Historical source: CONCHOLGY

***continued from the previous page***

Of the Com Slesa, Misca Madera, were alo fujected to chemical examination, by the fame philosopher. When needed tested best in a crochets, they emitted a perpetraphie fnell of them or feathers, and by further examination, by foliation, the proportion of carbonic matter degraded appeared to be greater, and the proportion of carbonate time lobe, than what was indicated by the refut of the experiments on pencilhenures fields. When natural shells belonging to the fame species, were amended in nitric acid very much diluted, the epidermis separated, and the whole of the carbonate of lime we differed. A gelatinoid diffusence, yearly in a liquid that, remains, ed, but it did not retain the figure of the shell, and ex-philic no appearance at Alorsra Hardure. These fields, therefore, contains a larger portion of gelatinous matter than the porcellaneous fluffs, but the other com-parent part counts entirely of carbonate time. ~~—Health~~ **Shells** composed of ~~nuclear-depart—Shells~~ **mother-of-pearl.—Shells** of this description were subjected to ~~familiar~~ **similar** experiments with ~~this~~ **the** former. When the common oyster was ~~expelled~~ **exposed** to a red heat, the effects were the ~~fans~~ **fame** as those which were produced by the fame ~~processis~~ **process** on the ~~facies~~ **species** of patches from Madeira. The foliation of the ~~hallmark~~ **unburnt** shell was also ~~funlier, except~~ **similar, excepting** only ~~little~~ **that** the gelatinous ~~parts are~~ **part was** of a greater ~~confidence.~~ **confiderency.** When the ~~river,~~ **river** muffel was burnt **in** a ~~crushing, the melted~~ **crucible, it emitted** much ~~make,~~ **fmoke,** with a ~~frongu find all~~ **ftrong fmell of** burnt ~~born~~ **horn** or cartilage; the ~~shell become~~ **fhell became** of a dark gray colour, and exfoliated. By foliation in the acids, the proportion of carbonic matter ~~Separated~~ **separated** was ~~greatest,~~ **greater,** and ~~but~~ **that** of ~~carbonaceous oil line~~ **carbonate of lime** obtained was ~~left,~~ **lefs,** than from the other ~~shells~~ **fhells** on which experiments were made. ~~My Mother~~ **69 Mothers** of ~~peat.~~ **pearl.** When an unburnt ~~fluef~~ **fhell** of this ~~description~~ **defcription** was ~~im-mected~~ **immerfed** in diluted nitric ~~oxide,~~ **acid,** a rapid ~~foliation~~ **folution** and ~~effir-vicosecond~~ **effervescence** took ~~place,~~ **place;** and at the end of two ~~days~~ **days,** the whole of the carbonate of lime was nearly ~~diffused.~~ **diffolved.** A feries of ~~means~~ **membranes** now only ~~remnants,~~ **remained,** of which the ~~epidemics constituted~~ **epidermis confituted** the ~~full. These~~ **firft. Thefe** membranes fill retained the figure of the ~~shell.~~ **fhell.** The carbonate of lime was ~~a flat~~ **at firft** readily ~~diffused, because~~ **diffolved, becaufe** the acid came ~~easily to contain~~ **eafily in contact** with it; but the ~~proceeds~~ **procefs** became flower, as it was more difficult for the acid to ~~indicate itself~~ **infinuate itfelf** between the different membranes of which the ~~shell~~ **fhell** is ~~compelled.~~ **compofed.** The ~~halted~~ **haliotis** iris, and the turbo ~~olearies,~~ **olearius,** were found to ~~rethable~~ **refemble** this muffel, except that the membranaceous parts were more compact and ~~dendric.~~ **denfe.** When these fields are deprived by an acid of the carbonate of lime, which gives them their birders, they appear to be formed: of different membranes, ~~appled~~ **applied** fratum fuer fratum. Each membranes is furmilled with a corresponding coarse cruml of carbonate of lime, and it is foisted, that it is always between every two membranes, beginning with the epidermis, and ending with the internal membrane, which has been latformed. The animals which inhabit both fleaified flesh, precede their habitation by the addi-tion of a fratum of carbonate of lime, which is fe-cured by a new membrane. And as every additional fratum exceeds in extent that which was previously formed, the shell becomes stronger in properties as it is enlarged; and thus the growth and age of the animal may be denoted by the number of frata of which the thell is compold. Similar experiments were made

## CONCHOLGY

*[Margin: Chap 1]*

C O N C H O L G Y. on pieces of mother of pearl as they are imported from China, China, and with precisely the fame refute. They appeated to be compelled of the fame grafiations matter such as carbonate of lime. In all the thiefs of this description which were named in aside, the membranes parts retained the each figure of the field, and they appeared different to be compelled of fibres, arranged in a parallel direction, correponding to the configuration of the flank. _Pearl_–The constituent parts of pearl appear to be ~~Pearl~~ familiar to ~~thief~~ **those** of ~~mother-of-pent.~~ **mother-of-pearl.** They are ~~compelled concerned costs~~ **composed** of ~~membranes~~ **concentric coats of membrane** and ~~colonies~~ **carbonate** of ~~lines,~~ **lime,** and resemble in ~~firstature~~ **structure** the globular, calcareous ~~con-cretions~~ **concretions** which are known by the name of ~~—phylidae.—~~ **__peloids.__** The ~~influence~~ **iridescence** and ~~produced~~ **undulated** appearance of pearl and ~~another-of-past,~~ **mother-of-pearl,** evidently depend on their lamellated ~~fracture~~ **structure** and ~~femitran/parency.~~ **semitransparency.** From their experiments, it appears, that shells are complex composed of surface of lime and gluten. In formas as pure in the porcellaneous shells, the proportion of carbonate. Time is great, while that of the animal matter is finally, and the mean one regarded as the beginning of the feries; while shells that come under the definition of mother-of-post are to be placed into the extra-mity, having a smaller proportion of carbonate of lime, and a greater proportion of membranous fusblance. It is the first the carbonate of lime is merely consumed by the animal matter; in the better the carbonate of lime-ferives to harden the membranous fusblance. But between fields are enzymes, in the proportion of carbone-nate of lime and animal glutes, of which alltaculous fulblances are compelled, there are no doubt numerous intermediates, arising from the nature of the animal to which they from a covering, its

peculiar ha-bits, or mode of life. Sect. II. ~~—Of~~ **Of** the Formation of ~~Shells.— Tuz thehell~~ **Shells. The fhell** or covering of ~~telacous~~ **teftaceous** animals, has been ~~considered~~ **confidered** as in fome ~~marine analogues~~ **meafure analogous** to the bones of other animals, although ~~it~~ **its** formation and growth are very different, ~~since~~ **fince** it ~~feres~~ **ferves** as a ~~hole coupport~~ **bafe or fuppport** to the ~~mulches,~~ **mucles,** which are attached to its ~~inter-fortified.~~ **internal furface.** The principal ~~side~~ **ufe** of the ~~shell,~~ **fhell,** however, is to ~~force~~ **ferve** as a covering or ~~defense~~ **defence** to the animal. ~~Telacous~~ **Telcaceous** animals are not only extremely different in external form, ~~had alls~~ **but also** in the mode of their ~~problem. Limtion.~~ **production.** Some are viviparous, as the ~~motif~~ **mot** of ~~their~~ **tho** which ~~people~~ inhabit bivalve shells, multivalves, and even ~~forms~~ **fome** of the ~~unitvives,~~ **univalves;** while the others, which form the ~~fire greetate proportions~~ **far greater proportion,** are ~~ostropus.~~ **oviparous.** In ~~non~~ **one** point, ~~how ever,~~ **however,** they all agree, that whatever be the mode of ~~probilution,~~ **production,** whether from an egg, or directly from the uterus of the mother, the ~~fluid~~ **shell** is formed on the body of the young animal, and is ~~propontioned~~ **proportioned** to its bulk. The bell defenders who have yet been made, and made the most elaborate investigation which has hitherto ophawvel peace, concerning the formation and development of iron. balls, are holes of the colonized Heamur, which were published in the Memoirs of the Academy of Sciences for the year 1790. The fate should be beenpecuculand by other authors, but their orbits have been mainly the arms so thole of this distinguished naturalidh. Klein is almost the only author who has advanced a different opinion. In his dissertation concerning the formation

## Source record

- Page ID: `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:473`
- Scan: `page-0473-4ab88a7103cd.jpg`, 600 x 812 px
- Image SHA-256: `4ab88a7103cd8f6c0c4f49cde0370890e3cbca2af72a6a90f8e7db56d053c32d`
- Extraction: `nvidia/nemotron-3.5-parse`, raw reply `artifacts/extract/raw/0473.json`
- Text is Parse OCR output, normalized (long s as s, line-end hyphens joined). It is not corrected by hand.
- ~~Struck-through~~ words are Parse's reading; **bold** words are Super VL's correction, checked against the scan image. Both are model output.