Historical source: UPTICS

continued: UPTICS

Ortial ferer has not any mark to direct him: hence aftermomers are obliged to enlighten the crois-hairs artifi, in order to render them vifible, and this without letting the luminous body interfere with the objefe which they intend to view , and, for doing this, two ways are propofel. 1. The ~~object glafs~~ **object-glafs** of the telcfore may be obliquely enlightened by placing a could near usit in our oblique frition, to that its smoke or flame may not interfere with the object. But if the object-glafs should happen to be pretty deep in the tube, it cannot be difficultly enlightened by this means i said bedties, if the telefcope is above fix feet loop, there will be a considerable difficulty in throwing a sufficient quantity of high upon the crois hairs. 2. By fome an opening is made in the tide of the tube near the focus of the object-glafs, though which the crois hairs are illuminated by means of a candle. But this method alfo is attended with inconveniences for the oblorer is incommeded by the light being the near his eyes, and the hair harmless become liable to accidents through their expelled frustration. An error also attends this method i, which is, that, according to the position of the light illuminating these hairs, they will appear in different situations. For example, when the horizontal hair is enlightened always be perceive a luminos line which may be taken for the hair itself, and which appears at its upper superficies. On the other hand, when the hair is enlightened underneath, the luminos line will appear as the under the face of i, and the corners will be the diameter of the hair, which often amounts to more than fix records. M. de la Hire, however, found a remedy for this convenience. He often obterred, that in monodhine nights, when the weather was a little foggy, the crois hairs were diflically fees; but when the heavens were ferene, they could occur by her corrected. Here therefore covered with a piece of gauge, or fine filk-craps, that end of the tube next the object-glafs i and this method succeeded to find the task at a good dilution from the telefope to enlightened the crapes, that the crats-hairs diflically appeared, and the fight of the fans was by no means obforced. In making the oblorations, a model glafs mulb uefd for preserving the eyes i and which may be thus prepared. Take two equal and well political round prices of glafs into the farles of one of which, all round its limb, glue a palkboard ring: then put the other piece of glafs into the smoke of a lamp, taking it several times time, and putting it in again, left the heat thould break it, until the smoke be fo thick, that the lamp can favorably be free through it, but the smoke mult not be all over of the same thickets, that fito a place may be obtain anfering to the splendour of the fun. This being done, the glafs, thus blackened, mult e glued to the pathboard of abovementand, with its black fide next to the other glafs, that the smoke may not be ribbed off. There are two kinds of ~~observation~~ **observations** relating to the ~~flares,~~ **flares:** one is, when they are in the meridian; the other, when they are in vertical circles. If the ~~polition~~ **position** of the meridian ~~ke~~ **be** known, the quadrant ~~wohl~~ **with** which the ~~oblorates~~ **observation** is ~~much, mult he~~ **made, must be** placed in the plane of that circle, and then the meridian altitudes are ~~really obferred~~ **easily observed** by means of the plumb-line. The meridian altitude of a flat may likewise be had by a pendulum clock, if the exact time of the ~~flat's~~ **flat's** pulling by the meridian be known. It ~~mult~~ **must** be ~~obffered,~~ **observed,** however, ~~therehas lines~~ **that stars have** the ~~fans of latitude~~ **same altitude** a minute before and after their ~~puffling~~ **passing** the ~~meridians,~~ **meridian,** if they be not in ~~on~~ **or** near the ~~zenith i,~~ **zenith;** but if they be, their altitudes ~~mult~~ **must** be ~~takes~~ **taken** every minute when they are in or near the meridian, and then their greatest or ~~their~~ **least** altitudes will be those in question. The ~~polition~~ **position** of a given vertical circle ~~mult he~~ **must be** found by the following method. 1. The ~~quadratus~~ **quadrant** and telescope remaining in the ~~frame~~ **same** situation wherein it was when the altitude of a ~~flux,~~ **star,** together with the time of its ~~puffage~~ **passage** by the ~~interjection~~ **intersection** of the ~~crois~~ **cross** hairs in the focus of the ~~object-glafs,~~ **object-glass,** was taken, we observe the time when the ~~fan,~~ **star,** or ~~frame facet flar, whole~~ **fixed star, whose** longitude and latitude is known, arrives to the vertical ~~bar~~ **hair** in the ~~telefops~~ **telescope;** and from ~~those~~ **thence** the ~~potion~~ **position** of the first vertical circle will be ~~hard,~~ **had,** and ~~alfo~~ **also** the ~~obferred flar's~~ **observed star's** true place. But if the ~~fun,~~ **sun,** or ~~fome olertas,~~ **some other star,** does not ~~ask~~ **pass** by the mouth of the ~~tubes,~~ **tube;** and if a meridian ~~liebe~~ **line be** otherwise ~~will~~ **well** drawn upon ~~allows,~~ **a floor,** or very level ground, in the place of ~~obloration i~~ **observation;** a ~~plan-like mull~~ **plumb-line must** be ~~fuufpared~~ **suspended** from a fixed place, at about 18 or ~~30~~ **20** feet from the ~~quadrata y~~ **quadrant;** under which a ~~matmulb te mode~~ **mark must be made** on the floor, in a right line with the plumb-line. You ~~mult~~ **must** next put a thin piece of ~~trade of palefoard~~ **pasteboard** very near the ~~object-glafs,~~ **object-glass,** in the middle of which there is a ~~fits~~ **slit** vertically placed, and ~~puffing~~ **passing** through the centre of the circular figure of the ~~object-glafs.~~ **object-glass.** Now, by means of this ~~fills,~~ **slit,** the before-mentioned plumb-line ~~area to prevent,~~ **may be perceived** through the ~~telefcope,~~ **telescope,** which before could not be ~~feny,~~ **seen,** because of its ~~moments.~~ **nearness.** Then the plumb-line ~~mail~~ **must** be removed and fulpated, ~~to~~ **so** that it be ~~prevented~~ **perceived** in a right line with the vertical hair in the focus of the ~~object-glafs,~~ **object-glass,** and a point marked on the floor directly under it. And if a right-line ~~the~~ **be** drawn through this ~~points,~~ **point,** and that marked under the plumb-line before it was ~~removal,~~ **removed,** the ~~full~~ **said** line will meet the ~~meridian draw~~ **meridian drawn** upon the ~~floor i,~~ **floor;** and ~~do~~ **so** we shall have the ~~polition~~ **position** of the vertical circle in which the observed ~~flar's,~~ **star's,** with respect to the meridian, the angle ~~where'~~ **whereof** may be measured in ~~affleming~~ **affix-**

ing known lengths upon the two lines from the point of ~~concavity,~~ **concourse;** for ~~fi,~~ **if,** through the extremities of ~~thefe~~ **these** known lengths, a line or ~~bafe~~ **base** be drawn, we shall have a triangle, ~~which~~ **whose** three ~~tides~~ **sides** being known, the angle at the ~~verce~~ **vertex** may be found, which will be the angle made by the vertical circle and meridian. Under the parallel Qaz waves, is flew the proper method of fixing that obfluent exactly in the meridian, but where the oblorer has no convivency of this kind, it will be proper to use a guarda quadrant; by means of which the altitude of a flare mulb be ob-hered a little before its puffage over the meridian, every minute, if polibbs, until its greatest or lealt altitude be bad, by which means, though we have not the true polition of the meridian, yet we know the meridian altitude of the flar. But although this method is very good, yet if a flatter splits by the meridian near the zenith, we cannot have its meridian altitude by repeated observations every minute, unless by chance, because in every minutes can now, the altriche arguments (z minutes of a degree i) and in thefe kind of observations, the inconvenient limitation of the oblerver, the variation of the flar's azimuth feveral de-

Source record

- Page ID: `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:1945`
- Scan: `page-1945-d877e2c0e7d6.jpg`, 600 x 824 px
- Image SHA-256: `d877e2c0e7d6979331208cad696389c92af546a9707a21816f1962452815285f`
- Extraction: `nvidia/nemotron-3.5-parse`, raw reply `artifacts/extract/raw/1945.json`
- Text is Parse OCR output, normalized (long s as s, line-end hyphens joined). It is not corrected by hand.
- ~~Struck-through~~ words are Parse's reading; **bold** words are Super VL's correction, checked against the scan image. Both are model output.