Historical source: ANATOMY

continued: ANATOMY

[Margin: Company the Antony | Saral vertive. | Cocgeyal or catalan vertebre. | Occuges or catalans vertebre.]

The number of ~~sacert verbren~~ **sacral vertebrae** is still more various even in the species of the same genus. Thus, while in several of the ~~age genes,~~ **ape genus,** in the ~~left;~~ **lori,** in the ~~trampve hat,~~ **vampire bat,** the ~~colour (globallyblack);~~ **colugo (galepithecus),** the ~~coat),~~ **coati,** and ~~wre~~ **two** of the ~~dielab,~~ **didelphis,** there is one ~~sacert whether only~~ **sacral vertebra only;** most of the apes have ~~actors~~ **sacra** consisting of 2, 3, 4, 5, or 6 pieces; the majority of other animals have ~~a secret verbetering~~ **3 sacral vertebrae;** the ~~hedgebo; preciping,~~ **hedgehog, porcupine,** guinea ~~pigs place, bare,~~ **pig, paca, hare,** tiger, several of the murine ~~genes,~~ **genus,** the ~~anti-acute~~ **anteater,** rhinoceros, ~~canal,~~ **camel,** dromedary, ~~damos,~~ **chamois,** goat, ~~donkey,~~ **sheep,** and ~~es,~~ **ox,** have ~~1s~~ **4;** the ~~chiq phant~~ **elephant** has 5; the ~~di~~ **Ai** 6; the ~~Email:~~ **Unau** 7; and in the mole, ~~while bear any ange,~~ **white bear, and quagga,** they ~~sea~~ **also** amount to ~~.~~ **7.** The ~~opancy~~ **frequency** of the three sacral ~~verbeter than~~ **vertebrae in** the ~~bore~~ **lower** animals shows that Galen, who ascribes only 3 to the human subject, must have derived this ~~reference~~ **inference** from the ~~femalc.~~ **former.** These ~~verbeter~~ **vertebrae** are in the mammals narrower than in man, and their direction forms with the ~~speeies~~ **spine,** instead of receding ~~backwards.~~ **backwards,** a straight ~~line,~~ **line;** an arrangement ~~of a dently~~ **evidently** connected with the horizontal position of the ~~forner.~~ **former.** The ~~idea~~ **shape** of the ~~secum on~~ **sacrum in** the lower mammals is that of an elongated ~~triangle,~~ **triangle;** and it is further remarkable, that in those species which occasionally assume the erect ~~attribution~~ **attitude** on the ~~local~~ **hind** leg, ~~an apex,~~ **as apes,** bears, and ~~slobs,~~ ~~in~~ **sloths, the** width of the ~~spectrum~~ **sacrum** is proportionally greater. The ~~saerul~~ **sacral** spines, which are short in man and the ape, become longer in the ~~Zovatura~~ **Zoophaga,** and ~~transa container rigles~~ **form a continuous ridge** in the rhinoceros, most ruminants, and especially in the mole. In the ~~trampve and~~ **vampyre bat** the ~~secum form~~ **sacrum forms** a long ~~component one,~~ **compressed cone,** the extremity of which is united ~~by~~ **to** the ~~lethal tusberaltics,~~ **ischial tuberosities,** without sustaining a ~~coccyw.~~ **coccyx.** The seal has two ~~several bones just~~ **sacral bones; but** the ~~Czaraba, e, qh oldphol~~ **Cetacea, e. g. the dolphin** and porpoise, are void both of ~~secum~~ **sacrum** and ~~coccyc.~~ **coccyx.** The ~~coceyed~~ **coccygeal** bones constitute the tail of the lower ~~animal,~~ **animals,** and **in** many instances they ~~were~~ **are** extremely ~~paneo.~~ **numerous.** The smallest number is ~~8,~~ **3,** which is that of the ~~mapel (onia glabrata, pillows, ettanus) buttery type;~~ **magot (simia sylvanus, pithecus, et inuus) or Barbary ape;** and the greatest yet known is that of the ~~anti-steer,~~ **anteater, in** which they amount to 40, and the long-tailed ~~minis,~~ **manis,** in which they amount to 45. Next to these may be placed that of the ~~coutta~~ **coaita** 32, the ~~babon~~ **baboon** 31, the ~~philology (didialyksia menus (dialyksia menus)~~ **phalanger (didelphis orientalis) 30,** the ~~panganaln 38,~~ **marmoset (didelphis murina) 29, the pangoin 28,** the silky monkey **(simia rosalia)** and ~~knowing~~ **black rat 26, the weeping monkey and howling** ape **25; the panther, mouse, dormouse, and elephant, 24; the lion, beaver, water-rat, Norway rat, and field-rat,** 23; the ~~pankzic, mouse, diseases, and depuis 34; the tiny beaver, western, Norway rural, and fade ant; 3; the bigeut, pana,~~ **flying-cat, puma,** cat, dog, ~~marns,~~ **marmot,** and ~~rhinoceros 32;~~ **rhinoceros, 22;** the ~~otter 24;~~ **otter, 21;** the Chinese monkey, ~~reason,~~ **raccoon,** civet, ~~hair~~ **hare,** and rabbit, 20; the tiger and wolf, 19; the macauco, glutton, ~~marnes,~~ **marten,** fat ~~demons, domodary, graffa,~~ **dormouse, dromedary, giraffe,** and ~~qugger,~~ **quagga,** 18; the ~~tissues, shovel,~~ **tarsier, shrew,** camel, and horse, 17; and other genera and ~~species~~ **species,** without any ~~determination~~ **determinate** order, descending so low **as** to ~~5, %,~~ **9, 8,** 7, **6,** and ~~44.~~ **4.** The ~~quilted bacheli, children, orinthorkhyues kylazio]~~ **quilled duckbill (echidna, ornithorhynchus hystrix)** has only ~~18~~ **12** caudal ~~vertebra,~~ **vertebrae,** while the common one ~~(carrying the predarodon)~~ **(ornithorhynchus paradoxus)** has at least ~~30.~~ **20.** The ~~globos~~ **gibbon** and ~~compares far too~~ **vampyre bat are** the only ~~maamifice ous~~ **mammiferous** animals, excepting the ~~Czaraba,~~ **Cetacea,** in which there are no ~~cocyed~~ **coccygeal** bones. ~~I showed hope~~ **It sometimes happens** that a monkey or ~~opposite~~ **opossum** loses a portion of its tail, when the truncated end is converted into a ~~hustlycence,~~ **knotty excrescence,** sometimes ~~carrato,~~ **curious,** always ~~differs~~ **different** from the taper point of the last ~~cocygeal vertebrae;~~ **coccygeal vertebra;** and in **this** case it is ~~different~~ **difficult** to ~~do terminate~~ **determine** the exact species. In the Czaraba, in which the absence of pelvis affords no mark to distinguish the lower vertebro into lumbar, sacral, and coccygeal, those below the dorsal may be ~~reprovided~~ **reregarded** as ~~fundso-coccpond;~~ **lumbo-coccygeal;** and their number is estimated by deducting that of the cervical and ~~coastal~~ **costal** from the total number. The following table, which shows the number of the ~~cost,~~ **costal,** the ~~lumber-eccygeal,~~ **lumbo-coccygeal,** and the total number of ~~vertebral~~ **vertebrae,** indicates that their number varies much in various genera of this family. 4. 1. 14 16 1c - Lanantin...16 24 46 Perpoie..13 40 60 Doping–10 14Vila...13 47 67 Hyperodon...3 30 54 Diplini...13 47 67 Hyperodon...3 30 45 Tories...19 30 Cedekar or White by Globeper –11 37 56 Whalesite 14-130 30 60 On Grisques...11 24 42 60 Greenland Walla...12 75 69 to Congress...12 24 46 Repopial...11 19 32 In general, however, if we reckon the first 23, 3, or 5 vertebres after the coastal, buther, it may be said that the cualth vertebron of the Czaraca was from 30 or 33, which are the numbers respectively in the humanants and dungory, on 34, 38; and 42, an adult they may be

estimain in the dolphin. We shall see that, in the dungog at least, we are guided in this estimate by the rudimentary bones of the polvis. The ~~coceyed~~ **coccygeal** or ~~cranial~~ **caudal** vertebrae of the ~~Mamatta~~ **Mammalia** may be distinguished into two kinds, those which ~~conma in~~ **contain a** canal in continuity with that of the vertebral ~~cohauman~~ **column** and ~~stern,~~ **sacrum,** and those in which the ~~places~~ **pieces** are ~~sold.~~ **solid.** The first, which are next the ~~secum,~~ **sacrum,** have articular, transverse, and spinous processes, distinct in proportion as the animal move ~~over the~~ **their** tails. The latter are generally ~~promised~~ **prismatic** in shape, diminish in size towards the ~~verymity,~~ **extremity,** and have only slight tubercles for ~~macular~~ **muscular** attachments. ~~Atimals~~ **Animals** with ~~prebiotics~~ **prehensile** tails, as the American ~~que (_ampeyona_),~~ **ape (sapajou),** have above, at the base of the body of each vertebra, two ~~and attacheders,~~ **small tubercles,** between which pass ~~he~~ **the** tendons ~~at~~ **of** the flexor muscles. By means of this mechanism these animals can twine the tail round the branch of a tree with sufficient force to support the weight of the body. The ~~Mareaama~~ **MAMMALIA** with long mobile ~~talk~~ **tails** have often ~~to~~ **two** or ~~these~~ **three** small ~~superheavy~~ **supernumerary** bones placed ~~in~~ **on** the lower surface of the junctions of several of the ~~coveged~~ ~~verteères,~~ **coccygeal vertebrae,** from the ~~six 44 sth~~ **3d or 4th** to the 7th or ~~Still.~~ **8th.** These ~~seasonal~~ **sesamoid** bones give attachment to muscles. In the beaver, which employs its tail as a ~~travel,~~ **trowel,** the transverse processes are remarkable for size, while the ~~bone~~ **lower** spinous processes are ~~large~~ **larger** than the ~~upperes—and arrangements~~ **upper ones,—an arrangement** which enables it to depress the ~~tall~~ **tail** forcibly when it beats the ground. The shape of the chest in the ~~Mamatta'va~~ **Mammalia** varies in ~~the chest.~~ general ~~is~~ **as** the clavicles are present ~~the waiting.~~ **or wanting.** In animals provided with clavicles, as the ~~QaradaraNA, falls,~~ **Quadrumana, bats,** the ~~spiral,~~ **squirrel,** beaver, mole, ~~ante-anteer,~~ **ant-eater,** hedgehog, and sloth, the shape of the ~~crosshead~~ **chest approaches** to the human, ~~we~~ **or** is ~~possible,~~ **conical,** and flattened before and behind. In those void of ~~clavicle~~ **clavicles** it is ~~compasses~~ **compressed** laterally, from the smaller ~~incouration~~ **incurvation** of the ~~tibia,~~ **ribs;** and the ~~titanium~~ **sternum** makes a ~~remain. The sterable~~ **remarkable** prominence, so that the transverse or ~~intercross disnum, under~~ **intercostal diameter** is less proportionally, and the ~~dense vertebral~~ **sterno-vertebral** is greater proportionally, ~~that~~ **than** in man and the claviculated ~~stimulus.~~ **animals.** In the ~~long-sized~~ **long-legged** animals, as the giraffe and those of the ~~stunk,~~ **stag kind,** this prominence of the ~~starum~~ **sternum** is sufficient to give it a ~~level-like~~ **keel-like** appearance ~~(_borne cerima-_tas).~~ **(thorax carinatus).** In the ~~converse~~ **carnivorous** animals the chest presents its greatest longitudinal extent. The number and shape of the ribs ~~varied~~ **varies** in the ~~differ. The ribs did tribes, in~~ **differ-ent tribes. In** number, indeed, the ~~riba~~ **ribs** always correspond with that of the ~~coral cerebretne.~~ **costal vertebrae.** Thus, in the ~~Qaradara NAJZODRAGA, RODSTATA, EENSTATA,~~ **QUADRUMANA, ZOOPHAGA, RODENTIA, EDENTATA,** and ~~DCIMAMATIA,~~ **RUMINANTIA,** they vary from ~~18~~ **12** to ~~10 pair~~ **15 pair,** with ~~one~~ **only** three exceptions, the ~~glutton byrena,~~ **glutton, hyena,** and ~~ante-teier.~~ **ant-eater.** In the Chinese monkey, common bat, and armadillo, they are a pair less than in

Source record

- Page ID: `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:2402`
- Scan: `page-2402-5030b31ea901.jpg`, 600 x 787 px
- Image SHA-256: `5030b31ea90173a447b4749960488cdeab085e4ff2f4bf49cf473614e50ce29e`
- Extraction: `nvidia/nemotron-3.5-parse`, raw reply `artifacts/extract/raw/2402.json`
- Text is Parse OCR output, normalized (long s as s, line-end hyphens joined). It is not corrected by hand.
- ~~Struck-through~~ words are Parse's reading; **bold** words are Super VL's correction, checked against the scan image. Both are model output.