Historical source: ANATOMY

ANATOMY

A N A T O M Y. Of the muticles 12 12, The first cervical nerves; 13 13, The Brain and femoral cervical nerves both front upon in muticles that lie on the neck, and teguration of the neck and head. 14 14. The third cervical nerves, which after feeding off (12 15, Scc) the placenta nerve to the diaphragm, supply the mufles and tegurements that lie on the fide of the neck and top of the shoulder. 16 16, The brachial plexus, formed by the fourth, fifth, forth, towards cervical, and frit dorial nerves–which lupply the mufles, and tegurement of the effective extremity. 17 17, The twelve dorial, or proper intercedul nerves, which is peut upon the intercedul muscles and tone of the large mufles which are upon the thorax. 18 18, The five lumbar part of the nerves, which puppy the lumbar and abdominal muscles, and forms of the tegurism and mufles of the inferior extremity. 10 17, The fourscoles, as preferred cranial nerve, formed by the two inferior lumber, and three superior of the os Herman. This large nerve fulpthe the general part of the medicale and teguements of the inferior extremity. 20. The thoracic plexus, formed by the eighth pair. 21 21, Branches of the follor or callic plexus, formed by the eighth pair and interCHAP. VI. OF THE SENSES, AND THEIR ORGANS. 140 IN treating of the ~~feeling,~~ **fenfes,** we mean to confine ~~ourselves~~ **ourfelves** to the external ones of touch, ~~triple, fineing,~~ **tafte, fmelling,** hearing, and vision. The word ~~[only],~~ **fenfe,** when applied to ~~their lives, fears us~~ **thefe five, feems to** imply not only the ~~forlation occurred~~ **fenfation excited** in the ~~limited~~ **mind** by certain ~~impurities~~ **impreffions** made on the body, but ~~likewise~~ **likewife** the organ defined to receive and ~~transmit basic ingredients in~~ **tranfmit thefe impreffions to** the ~~forbidden.~~ **fenforium.** Each of ~~these~~ **thefe** organs being ~~at~~ **of** a peculiar ~~threat,~~ **ftructure,** is ~~if dependible~~ **fufceptible** only of particular ~~impretions,~~ **impreffions,** which will be pointed out as we proceed to ~~define~~ **defcribe** each of them fe parately. Sect. I. _Of Touch._ ~~collals,~~ **cofials,** which ~~lupply~~ **fupfly** the flomach and ~~chylopotic vience.~~ **chylopoetic vifcera.** 22 ~~19. Branch~~ **22, Branches** of the ~~figure~~ **fuperior** and ~~anterior mesenteric plexus,~~ **inferior mefenenteric plexufes,** formed by the eighth pair and ~~fifth tercedula,~~ **intercostal,** which ~~lupply~~ **fupfly** the ~~chyopotic vience. With~~ **chylopoetic vifcera, with** part of the organs of urine and generation. 23 23, ~~Nerve~~ **Nerves** which accompany the ~~fermatic~~ **fpermatic** cord. 24 24, The ~~hypogotic~~ **hypogaftric** plexus, ~~while fusplies~~ **which fupplies** the organs of urine and ~~generates~~ **generation** within the ~~pelvis.~~ **pelviis.** Fig. 2, 3, ~~4.~~ **4, 5.** Show different views of the Inferior part of the Brain, cut perpendicularly through the ~~Middle-with,~~ **Middle—with** the Origin and large portions of all the ~~Nerve~~ **Nerves** which ~~puts~~ **pass out** through the Bones of the Cranium,—and the three ~~full~~ **first** Cervicals. The anterior lobe, B, The lateral lobe of the cerebellum. C, One of the lobes of the cerebellum. ~~To Tube aunture, K,~~ **D, Tuber annulare, E,** Corpus ~~pyramidal,~~ **pyramidale,** in the middle of the medulla oblongata. ~~G,~~ **F,** The corpus olivare, in the ~~fide~~ **side** of the medulla oblongata. G, The medulla oblongata. H, ~~Hamilton mediala insula. Naxxta - 2, 3,4,~~ **The medulla spinalis. 2 3 4 5** 6 ~~5~~ **7 8** and ~~3 Plate~~ **9, Pairs** of nerves. 10 ~~16,~~ **10,** Nervus ~~accoltionis,~~ **accessorius,** which comes ~~from—11,~~ **from—11,** 12, and 13, the three first cervical nerves. Sect. II. ~~_Of Teife._ This feeds~~ **Of Taffe. The fenfe** of ~~safe~~ **tafe** is ~~faced~~ **feated** chiefly in the ~~tongue,~~ **tongue ;** the ~~fixation~~ **fituation** and figure of which are ~~sufficiently~~ **fufficiently** known. On the upper further of this organ was may observe a great number of pupils, which, on occasion of their difference in free and hays, are commonly divided into three clades. The largest are trusted towards the final of the tongue. Their main community varies from fever to nine, and they form to be mucous folicles. Thief of the second clats and tunnels that matter, and of a cylindrical shape. They are most humorous about the middle of the tongue. Those of the third clats are very minute, and of a conical shape. They are 141 In this feature of touch may be defined to be the faculty of ~~diltinguling~~ **distinguishing** certain properties of bodies by the feel. In a general accepted, this definition might perhaps not properly be extended to every part of the body political of identifying ( ), it is commonly combined to the nervous pupils, and consequently of exciting differences in the fenision communes. But the origin of touch, like all the other feels, it not equally distinct in every part of the body, or in every subjects, being in force much more equitable than it to others. (v) In the course of this article, mention has often been made of the feasibility or ~~interfilability~~ **infeasibility** of different parts of the body: it will therefore, perhaps, not be ~~amido~~ **amiss** to ~~oblive~~ **observe** in this place, that many parts which were formerly ~~tipped~~ **supposed** to ~~polls~~ **possess** the ~~molt exquiteite fenle,~~ **most exquisite feeling,** are now known to have but little or no feeling, as leaf in a found story, for its inflamed there, even the bones, the ~~molt~~ **most** infertible part of any, become susceptible of the ~~molt~~ **most** painful ~~fentiations.~~ **sensations.** This curious discovery is due to the late late Haller. His ~~experiment~~ **experiments** prove, that the ~~females, cartilage,~~ **bones, cartilages,** ligaments, tendons, ~~epadenies,~~ **epidermis,** and membranes (as the ~~plantar, periardium, diam~~ **pleura, pericardium, dura** and just master, peripheral, ~~Sec)~~ **&c.)** may in a healthy thus ~~he~~ **be** considered in infinities. As ~~fertilising~~ **sensibility** depends on the brain and nerves, of ~~courie~~ **course** different parts will ~~polls~~ **possess** a greater or ~~lefs~~ **less** degree of feeling, in proportion as they are ~~fluppled~~ **supplied** with a greater or ~~insufficient~~ **smaller** number of nerves. Upon this principle it is, that the ~~film, mulcles, domonic, interfere,~~ **skin, muscles, romench, intestines,** urinary bladder, ureters, ~~vagues, pens,~~ **uterus, vagina, penis,** tongue, and ~~setina,~~ **retina,** are extremely feasible, while the lungs and glands have only an obscure degree of feeling.

Source record

- Page ID: `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:2277`
- Scan: `page-2277-cdde2fc730b9.jpg`, 600 x 783 px
- Image SHA-256: `cdde2fc730b9b43e147b41385c61a207df3d0519a0ec368e9bbc3b3ecd4432de`
- Extraction: `nvidia/nemotron-3.5-parse`, raw reply `artifacts/extract/raw/2277.json`
- Text is Parse OCR output, normalized (long s as s, line-end hyphens joined). It is not corrected by hand.
- ~~Struck-through~~ words are Parse's reading; **bold** words are Super VL's correction, checked against the scan image. Both are model output.