

Theoretical Astroparticle Physik

Homework 2

Chenhuan Wang and Koay Yong Sheng

May 10, 2020

1 Quickies

- (a) What is the definition of the cosmological redshift?

Cosmological redshift is the redshift caused by the expansion of the universe itself. If a photon is emitted at time t_i with a given wavelength, λ_i , then the observed wavelength of the photon at present time (with the cosmological redshift, z) is

$$\lambda_0 = \lambda_i \frac{a_0}{a(t_i)} = \lambda_i [1 + z(t_i)] \quad (1.1)$$

where a_0 is the scale factor at present time and $a(t_i)$ is the scale factor at time t_i .

- (b) Explain the notion of co-moving coordinates.

Since in GR we learned that physics doesn't care what coordinate system we choose, we can choose one that is "natural"/easiest. In the comoving frame, the worldlines of particles at rest in this frame are geodesics, i.e. particles initially at rest will remain at rest. Particles that are initially moving with respect to this frame will eventually come to rest in it. The comoving coordinates assign constant spatial coordinate values to observers who perceive the universe as isotropic. The comoving time coordinate is just the proper time by an observer at rest in the comoving frame. It is also the elapsed time since the Big Bang according to a clock of a comoving observer and is a measure of cosmological time. Comoving distance is the distance between two points measured along a path defined at the present cosmological time. The comoving spatial coordinates tell where an event occurs while cosmological time tells when an event occurs.

2 Equation of state for matter and radiation

- (a) Compute the number density of non-relativistic particles

Maxwell-Boltzmann distribution can be approximated as

$$f_{\text{MB}}(\vec{p}) = \frac{1}{(2\pi)^3} \exp\left(-\frac{E(\vec{p}) - \mu}{T}\right) \approx \frac{1}{(2\pi)^3} \exp\left(\frac{\mu - m}{T} - \frac{|\vec{p}|^2}{2mT}\right)$$

For now on $|\vec{p}| = p$. The number density is defines as

$$\begin{aligned} n &= g_i \int d^3p f(\vec{p}) \\ &= \frac{g_i}{(2\pi)^3} \exp\left(\frac{\mu - m}{T}\right) \int d^3p \exp\left(-\frac{|\vec{p}|^2}{2mT}\right) \\ &= \frac{4\pi g_i}{(2\pi)^3} \exp\left(\frac{\mu - m}{T}\right) \int d|\vec{p}| |\vec{p}|^2 \exp\left(-\frac{|\vec{p}|^2}{2mT}\right) \\ &= \frac{4\pi g_i}{(2\pi)^3} \exp\left(\frac{\mu - m}{T}\right) \frac{\sqrt{\pi}}{2} (2mT)^{3/2} \\ n &= g_i \exp\left(\frac{\mu - m}{T}\right) \left(\frac{mT}{2\pi}\right)^{3/2} \end{aligned} \tag{2.1}$$

- (b) Determine the pressure and find the equation of state

The pressure is given as

$$\begin{aligned} P &= g_i \int d^3p \frac{p^2}{3E} f(\vec{p}) \\ &= \frac{g_i}{(2\pi)^3} \frac{4\pi}{3} \exp\left(\frac{\mu - m}{T}\right) \int dp p^2 \frac{p^2}{E} \exp\left(-\frac{p^2}{2mT}\right) \\ &\approx \frac{g_i}{6\pi^2 m} \exp\left(\frac{\mu - m}{T}\right) \int dp p^4 \exp\left(-\frac{p^2}{2mT}\right) \\ &= \frac{g_i}{8\pi^2 m} \exp\left(\frac{\mu - m}{T}\right) (2mT)^{5/2} \sqrt{\pi} \\ &\stackrel{2.1}{=} nT \end{aligned} \tag{2.2}$$

In the limit $T \ll m$, the equation of state is

$$\omega = \frac{P}{\rho} = \frac{T}{m} \approx 0 \tag{2.3}$$

(c) Determine the energy density in the limit $T \gg E \gg m$

First to show $E \, dE = p \, dp$

$$\frac{dE}{dp} = \frac{d}{dp} \sqrt{m^2 + p^2} = \frac{E}{p}$$

$$\begin{aligned} \rho &= g_i \int d^3p \, E(\vec{p}) f(\vec{p}) \\ &= \frac{g_i}{(2\pi)^3} \int \frac{d^3p \, E(\vec{p})}{\exp(E(\vec{p})/T) \mp 1} \\ &= \frac{g_i}{2\pi^2} \int_0^\infty \frac{dp \, p^2 E}{\exp(E(\vec{p})/T) \mp 1} \\ &= \frac{g_i}{2\pi^2} \int_m^\infty \frac{dE \, E^2 \sqrt{E^2 - m^2}}{\exp(E(\vec{p})/T) \mp 1} \\ &\approx \frac{g_i}{2\pi^2} \int_0^\infty \frac{dE \, E^3}{\exp(E(\vec{p})/T) \mp 1} \\ &= \begin{cases} \frac{g_i}{2\pi^2} T^4 \zeta(4) \Gamma(4) & \text{bosons} \\ \frac{g_i}{2\pi^2} T^4 \frac{7\pi^4}{120} & \text{fermions} \end{cases} \\ &= (1; 7/8) \cdot g_i \frac{\pi^2 T^4}{30} \end{aligned}$$

Here $(1; 7/8)$ denotes that it is 1 for bosons and $7/8$ for fermions. For boson case, the integral representation of riemannian zeta function is used. Mathematic did the calculation of fermions.

(d) Determine the pressure and further the equation of state

Following the same recipe as before

$$\begin{aligned} P &= \frac{g_i}{(2\pi)^3} \int d^3p \, \frac{p^2}{3E} \frac{1}{\exp(E/T) \mp 1} \\ &\approx \frac{g_i}{6\pi^2} \int_0^\infty \frac{dE \, E^3}{\exp(E/T) \mp 1} \\ &= \rho/3 \end{aligned} \tag{2.4}$$

This is the equation of state for ultra-relativistic particles with $\omega = 1/3$.

3 Equation of state for an exotic scalar field

(a) Using the identity,

$$\frac{\delta(\det A)}{\det A} = \det A \operatorname{tr} \left(\frac{\delta A}{A} \right) \quad (3.1)$$

One can show that

$$\frac{\delta g}{g} = g^{\mu\nu} \delta g_{\mu\nu} \quad (3.2)$$

$$\delta g = g g^{\mu\nu} \delta g_{\mu\nu} \quad (3.3)$$

Then,

$$\delta \sqrt{-g} = -\frac{1}{2\sqrt{-g}} \delta g \quad (3.4)$$

$$= -\frac{1}{2\sqrt{-g}} g g^{\mu\nu} \delta g_{\mu\nu} \quad (3.5)$$

$$= -\frac{1}{2} \sqrt{-g} g^{\mu\nu} \delta g_{\mu\nu} \quad (3.6)$$

Now, our Energy-momentum Tensor:

$$T_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L})}{\delta g^{\mu\nu}} \quad (3.7)$$

$$= \frac{2}{\sqrt{-g}} \left[\frac{\delta(\sqrt{-g})}{\delta g^{\mu\nu}} \mathcal{L} + \sqrt{-g} \frac{\delta \mathcal{L}}{\delta g^{\mu\nu}} \right] \quad (3.8)$$

$$= \frac{2}{\sqrt{-g}} \left(-\frac{1}{2} \sqrt{-g} g_{\mu\nu} \right) \mathcal{L} + 2 \frac{\delta \mathcal{L}}{\delta g^{\mu\nu}} \quad (3.9)$$

$$= -g_{\mu\nu} \mathcal{L} + 2 \frac{\delta \mathcal{L}}{\delta g_{\mu\nu}} \quad (3.10)$$

Given that the lagrangian is

$$\mathcal{L} = -V_0 \sqrt{1 - g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi} \quad (3.11)$$

$$(3.12)$$

Then,

$$\frac{\delta \mathcal{L}}{\delta g^{\mu\nu}} = -V_0 \left(\frac{1}{2} \right) \left(\frac{-\delta_\mu^\alpha \delta_\nu^\beta \partial_\alpha \phi \partial_\beta \phi}{\sqrt{1 - g^{\lambda\rho} \partial_\lambda \phi \partial_\rho \phi}} \right) \quad (3.13)$$

$$= \frac{V_0}{2} \frac{\partial_\mu \phi \partial_\nu \phi}{\sqrt{1 - g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi}} \quad (3.14)$$

Coming back to our Energy-momentum Tensor:

$$T_{\mu\nu} = -g_{\mu\nu}\mathcal{L} + \frac{V_0\partial_\mu\phi\partial_\nu\phi}{\sqrt{1 - g^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi}} \quad (3.15)$$

$$= V_0 \frac{g_{\mu\nu}(1 - g^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi) + \partial_\mu\phi\partial_\nu\phi}{\sqrt{1 - g^{\alpha\beta}\partial_\alpha\phi\partial_\beta\phi}} \quad (3.16)$$