COMPONENT · DEMONSTRATION KITCHEN

SESAME SALT GARNISH

A loose, lightly crushed sesame finish for grilled vegetables. White sesame brings the body; whole black seeds add contrast. Stop crushing while the mixture still falls freely from a spoon.

TOTAL TIME
3 minutes
Active 3 min
Inactive 0 min

DIFFICULTY
Basic
Controlled mortar crushing

YIELD / PORTION
128 g usable
64 portions at 2.0 g

STORAGE
Unconfirmed
House SOP required

R&D — NOT RELEASED FOR PRODUCTION. Approved holding and storage conditions, container labeling, use-by basis and discard rule; Second-cook reproduction with chef-approved tolerances for yield, crushing endpoint and portioning; Explicit production release for this exact final version.

DEMO-001 · v0.4 · 2026-10-26
Equipment / tested load: 300 mL mortar and pestle, spoon, dry scraper, dry receiving container; scale readable to 0.1 g. One 130 g input batch only.
Allergens: Sesame. Simulated label review lists no additional declared allergens. Shared utensils need house cross-contact controls; no free-from claim.
Percentage basis: Ready-to-use roasted white sesame = 100% = 100 g.

INGREDIENTS

INGREDIENT	WEIGHT	VOLUME	SCALING %
For the garnish			
I1 · Purchased roasted white sesame Ready-to-use; label checked in the fictional trial reply; step 1. Volume is a labeled proxy; weight authoritative.	100 g	≈ 11.3 tbsp (US)	100%
I2 · Purchased roasted black sesame Ready-to-use; step 3. Volume is a labeled proxy; weight authoritative.	20 g	≈ 2.25 tbsp (US)	20%
I3 · Fine salt Step 3. Volume is a labeled proxy; weigh the salt; weight authoritative.	10 g	≈ 1.7 tsp (US)	10%
Measured usable yield	128 g	—	—

Basis: Ready-to-use roasted white sesame = 100% = 100 g. Input 130 g. Measured usable output 128 g. Transfer loss 2 g. Complete portions 64. Volumes are researched approximations from cited sources, not measured spoonfuls; weigh ingredients.

METHOD

CRUSH

1. Weigh white sesame (I1) into the 300 mL mortar. Use 10 press-and-turn strokes, approximately 30 seconds from the first stroke.
2. Check the texture: roughly half the seeds should split. The mixture must remain loose, with no oily paste.

   Roughly half the seeds should split. The mixture must stay loose, with no oily paste.

MIX

3. Add black sesame (I2) and salt (I3). Stir with a spoon for 20 turns, approximately 20 seconds from the first turn. Total mortar load: 130 g.

   Settings — speed rpm spoon stirring, 20 turns
4. Check that black seeds are distributed through the mixture and no salt pockets remain. Log any failed endpoint for trial review.

WEIGH & PORTION

5. Scrape into a tared, dry container. Record actual usable mass and transfer loss. Reference trial: 128 g usable from 130 g input.
6. For the simulated plating trial, weigh 2.0 g per plate and sprinkle evenly across the vegetable surface immediately before dispatch. No bowl garnish.

   Three minutes total active preparation. Do not force actual yield to match the reference result.

LABELING

product name: Sesame salt garnish

batch or date code: NEEDS CHEF CONFIRMATION - date code convention is not set

use by basis: NEEDS CHEF CONFIRMATION - no approved shelf-life basis located

storage instruction: NEEDS CHEF CONFIRMATION - approved storage conditions and container are unresolved

allergen declaration: Sesame

preparer or station: NEEDS CHEF CONFIRMATION - preparer and station are not recorded

SERVICE & PLATING

This card standardizes the garnish only. Vegetable quantity, plate dimensions and finished-dish service temperature belong in the linked assembly card,

HOLDING & STORAGE

Unresolved: approved conditions, container, label, use-by and discard rule. No shelf life is assigned. Cooling and reheating are not

which has not been supplied. No verified plating photograph is available.

SCALING & VARIATIONS

At factor 2: 200 g white sesame, 40 g black sesame and 20 g salt. Run two separate reference-size batches; weigh each output. Do not double mortar load or crushing time. Finished yield is not guaranteed. No substitutions are approved.

VOLUME RESEARCH AND SOURCES

Volumes are researched approximations, not measurements; gram weights stay authoritative. Source 1: King Arthur Baking, Ingredient Weight Chart, https://www.kingarthurbaking.com/learn/ingredient-weight-chart, accessed 2026-10-26, listing sesame seeds at 1/2 US cup = 71 g, that is 8.875 g per US tablespoon, with US measures (1 US cup = 236.6 mL, 1 US tbsp = 14.79 mL, 1 US tsp = 4.93 mL). Source 2: USDA FoodData Central SR Legacy 173468, Salt, table, https://fdc.nal.usda.gov/food-details/173468/nutrients, accessed 2026-10-26, listing 1 tsp = 6 g, 1 tbsp = 18 g and 1 cup = 292 g in US measures. Units convention: US tablespoon, teaspoon and cup throughout. An Australian 20 mL tablespoon is not mixed with these figures, and no metric measure is used on this card. Proxy status: rows I1 and I2 use the generic whole-sesame-seed ratio because the purchased brand and roast state were not provided, so both are labeled proxies rather than exact matches; row I3 uses the generic table-salt ratio because the salt type was not provided. Fine and kosher salts are not interchangeable - table salt at 18 g per US tbsp, Morton's kosher at 16 g and Diamond Crystal at 8 g, a spread of about 2.25 times for the same 10 g - so the salt volume is a cross-check only and the 10 g weight remains the production control.

applicable to this unheated component. Production remains blocked.

CONTROLS & CORRECTIVE ACTION

Check seed breakage, absence of paste, mixing and output on every trial. Record deviations for chef review. Recheck current supplier labels and house utensil controls. No validated food-safety critical control point is asserted; applicable controls await house SOP review.

VOLUME ARITHMETIC AND ROUNDING

I1: 100 g divided by 8.875 g per US tbsp = 11.27 tbsp, shown as approximately 11.3 tbsp, rounded to the nearest 0.1 tbsp. I2: 20 g divided by 8.875 g per US tbsp = 2.25 tbsp, shown as approximately 2.25 tbsp, rounded to the nearest 0.05 tbsp. I3: 10 g divided by 6 g per US tsp = 1.67 tsp, shown as approximately 1.7 tsp, rounded to the nearest 0.1 tsp. Cup equivalents at 142 g per US cup are approximately 0.70 cup for I1 and approximately 0.14 cup for I2. Every volume on this card is a derived approximation: where an exact measure is needed, weigh the ingredient and measure the spoon.

REQUIRED BEFORE PRODUCTION RELEASE

- Confirm holding/storage, labeling, use-by and discard SOP.
- Complete second-cook reproduction; approve yield, texture and portion tolerances.
- Obtain explicit chef release for the exact final version after controls are resolved.

Evidence: Fictional demonstration card. Original fictional notes and chef replies; no actual photo inspected. Ingredient weights, method and usable yield are simulated statements. Percentages, 64 portions and factor-2 quantities are calculations. Household volumes are researched approximations derived from the cited public sources with their access dates, not measured spoonfuls.

Validation: One simulated batch; no real cook test and no production approval. New trial with no predecessor. Local demonstration only.

Filing: Local demonstration only. Proposed relative route: R&D / needs-confirmation. No remote write attempted.