

Free Amino Acids in Human Milk: Time and Infant Gender

Federico J. Zertuche

March 28, 2019

Abstract

This report has the objective of comparing our results with those in the litterature, specifically those found here:

<https://www.mdpi.com/2072-6643/10/9/1233>

It is divided into 3 parts:

The first is about understanding the data. I try to describe and motivate the models, significant results and thier relationship with existing literature and medical knowledge.

In the second, the models are used to understand what should we be looking forward to in future experiments. Here, I take the models seriously and device possible predictions that are compatible with current knowledge.

The third part is about missing data. I mention what I did with the missing patients.

Part I

Three Objectives, Three Models

I am interested in answering 3 questions:

1. Do amino acid levels change over time?

2. Are these levels different in the milk for boys and girls?
3. If the concentration changes over time, is it different for boys and girls?

I use 3 regression models to summarize the effects for each one of the questions. The answers are given as coefficients with confidence intervals.

Before analyzing the results let me introduce some notation.

I will write AA for the concentration level of a particular amino acid. The data contains the concentration of several amino acids. In general, I will study them individually. I use the essential and non essential classification since I am interested in the difference between nourishment and body production.

The measurements were taken at weeks 1, 2, 8 and 16 so the variable corresponding to time is called *week*. This is not a categorical variable.

I note the infant genders *boy* and *girl* and the variable will be called *sex*.

There is an additional variable called *group* that tells us if the mother was a *teen* or an *adult* when she gave birth.

1 Do AA levels change over time?

Here the model used is:

$$AA = \alpha_0 + \alpha_1 \text{ week} + \alpha_{id} \quad (1)$$

Each coefficient is the summary of an effect.

The first, α_0 , is the concentration at week 0, I will not talk about this coefficient. The second one has units *concentration/time*, it is the main focus in this part of the analysis.

The last one represents a random effect that is different for each one of the participants, think about it as all the patients differences that were not controlled for during the data collection process.

1.1 Essential Amino Acids

The list of amino acids with significant effects (figure 1) over time is: THR, VAL, LUE, PHE. As you can see in figure 2, in all of these amino acids, there is a general increasing or decreasing trend. The coefficients are meant to capture this. This model does not capture phenomena like the decrease from




Figure 1: Week coefficients of model (1) for essential amino acids.

week 8 to 16 in PHE and VAL. Think of the model as a first approximation that helps us with classifying the concentrations.

Four amino acids have non significant effects: HIS, ILE, MET and TRP. Figure 3 shows their mean concentration over time. In each of the figures there is an abrupt change. For example, TRP jumps between positive and zero concentrations, probably due to the size of the measurements (the high-




Figure 2: Mean concentration per week with a linear regression line in blue. Essential amino acids with significant week effects.

est vaule is 0.03). In ILE and MET there is an abrupt decrease from month 8 to 16. Finally, HIS has a big increase from week 1 to 2 and a steady decrease from there on. Once again, all these non linear patterns are not very well described by model (1).




Figure 3: Mean concentration per week with a linear regression line in blue. Essential amino acids with non significant week effects.

1.2 Non Essential Amino Acids

The effects for the non essential amino acids are shown in figure 4. Figure 5 shows the small coefficients in detail.

The concentrations of *ARG* and *PRO*, in figure 6, decrease in a similar pattern as that of *LYS* in figure 2. First a big drop from week 1 to 2 and a

slower decrease afterwards.

All the other significant effects are positive, the concentration of GLU, GLN, ALA, SER, GLY, ASP, CYS and TYR increases over time as shown in figure 6.

The only non significant effect is that of ASN. Figure 7 shows how its concentration changes over time: first a big drop and then a big increase. Studying what happens between weeks 2 and 8 might help in understanding how this concentration varies over time.




Figure 4: Week coefficients of model (1) for non essential amino acids.




Figure 5: Detail of the week coefficients of model (1) for the non essential amino acids with small concentrations.




Figure 6: Mean concentration per week with a linear regression line in blue. Non essential amino acids with significant week effects.




Figure 7: Mean concentration per week with a linear regression line in blue. Non essential amino acid ALA has a non significant week effect.

2 Are AA levels different in the milk for boys and girls?

There is a time effect for most amino acids. Here I am interested in studying whether the milk for boys and girls has different concentrations. The model:

$$AA = \alpha_0 + \alpha_1 \text{ week} + \alpha_2 \text{ sex} + \alpha_{id} \quad (2)$$

has an additional effect that depends on the infant gender - α_2 . The rest of the coefficients are the same as before. This effect is only present for boys. So another way of writing model 2 is:

$$AA = \begin{cases} \alpha_0 + \alpha_1 \text{ week} + \alpha_2 + \alpha_{id} & \text{if boy,} \\ \alpha_0 + \alpha_1 \text{ week} + \alpha_{id} & \text{if girl.} \end{cases}$$

The coefficient for sex describes the difference in concentration between boys and girls. If $\alpha_2 > 0$, then boys have a higher concentration of that particular amino acid, if it is 0, there is no difference and when the coefficient is negative, girls have a higher concentration.

2.1 Essential Amino Acids

There is no significant differences for all of the essential amino acids. Figure 8 shows the estimated coefficients. Most of the amino acids have a higher concentration for boys except for THR. Once again TRP concentrations are very small and sometimes 0, so its coefficient is close to 0 too.

The effects of VAL, HIS, ILE, LEU are similar to those in [1] and the rest have an opposite sign. In [1] all of the effects were non significant too.

2.2 Non Essential Amino Acids

For non essential amino acids some effects are significant and their sign coincide with the estimations found in [1].

The effects of GLU, GLN, GLY, ALA, SER, ASP have the same sign as those in [1]. On the other hand, TYR, ARG, ASN have opposite signs.

In figures 9 and 10, there are 4 significant differences for girls and boys. They correspond to amino acids GLU, GLY, TYR and CYS. The effects




Figure 8: Sex coefficients of model (2) for essential amino acids.

of GLU and GLY have the same sign as in article [1] but here they are significant.

In the study [1], they could not measure CYS. Here it has a significant difference.

Finally, the sign of TYR is different from that in [1] but here it has a significant effect.




Figure 9: Sex coefficients of model (2) for non essential amino acids.





Figure 10: Detail of the sex coefficients of model (2) for the non essential amino acids with small concentrations.



(D)

Figure 11: Sex coefficients found in article [1].

3 If the concentration changes over time, is it different for boys and girls?

Now we know that there is a time effect for almost all amino acids and a sex effect for 4 non essential amino acids (GLU, GLY, TYR and CYS). In this section I study whether the variation of concentration over time changes for girls and boys.

For the model I add yet another effect:

$$AA = \alpha_0 + \alpha_1 \text{ week} + \alpha_2 \text{ sex} + \alpha_3 \text{ week} \times \text{sex} + \alpha_{id} \quad (3)$$

where the coefficient α_3 , that corresponds to the interaction term $\text{week} \times \text{sex}$, can be interpreted in two different ways. First by rewriting model 3 as:

$$AA = \alpha_0 + (\alpha_1 + \alpha_3 \text{ sex}) \times \text{week} + \alpha_2 \text{ sex} + \alpha_{id} \quad (4)$$

you can read $\alpha_1 + \alpha_3$ as the time effect for boys and α_1 as the time effect for girls. With this notation α_3 is the time difference between boys and girls.

The second interpretation follows from the other possible arrangement, namely:

$$AA = \alpha_0 + \alpha_1 \text{ week} + (\alpha_2 + \alpha_3 \text{ week}) \times \text{sex} + \alpha_{id} \quad (5)$$

here $\alpha_2 + \alpha_3$ is the sex effect per week. So for a given week, the difference between boys and girls is $\alpha_2 + \alpha_3$. As before this difference can be positive, negative or zero.

Either way the interaction coefficient α_3 represents a difference over time. So I will focus on this coefficient first.

3.1 Essential Amino Acids

As expected, all of these effects are smaller than the main effects. Also the coefficient are non significant. However, the trends suggest that the concentration increases faster for girls over time for all amino acids except HIS. The variation in concentration of essential amino acids over time is similar for boys and girls.

Figure 13 shows the amino acids whose concentration increases more rapidly over time for girls. For HIS they concentration for boys is bigger over time except at week 8.

Finally, figure 15 shows TRP. As before it jumps between positive values and 0.




Figure 12: $Sex \times week$ coefficients of model (3) for essential amino acids.




Figure 13: Mean concentration per week with a linear regression line in blue for boys and pink for girls. Essential amino acids with a bigger interaction effect for girls.




Figure 14: Mean concentration per week with a linear regression line in blue for boys and pink for girls. Essential amino acids with a bigger interaction effect for boys.




Figure 15: Mean concentration per week with a linear regression line in blue for boys and pink for girls for TRP.

3.2 Non Essential Amino Acids

Now, most of the effects are bigger for boys although they remain non significant. The concentration increases in a similar fashion for both sexes as can be seen in figures 17 and 18.

In both ARG and PRO (figure 18), the concentration for girls decreases

slower mostly due to a jump in week 8.




Figure 16: $Sex \times week$ coefficients of model (3) for non essential amino acids.




Figure 17: Mean concentration per week with a linear regression line in blue for boys and pink for girls. Non essential amino acids with a bigger interaction effect for boys.




Figure 18: Mean concentration per week with a linear regression line in blue for boys and pink for girls. Non essential amino acids with a bigger interaction effect for girls.

Part II

What to look forward in future experiments?

Some aspect of the data are not discussed in part I. First I explore if the AA concentration is due to differences in baby weights and sizes. Then, I study the role of the group variable in the concentration.

4 Baby weights and sizes

Besides the concentrations, sex and group, baby weights and head circumferences were measured too.

Figures 19 and 20 show the mean weight and mean head circumference over time for girls and boys. There are some differences but they are non significant. For example, the estimated coefficients for the regression models:

$$weight = \alpha_0 + \alpha_1 \text{ week} + \alpha_2 \text{ sex} + \alpha_3 \text{ week} \times \text{sex}$$

and

$$head = \alpha_0 + \alpha_1 \text{ week} + \alpha_2 \text{ sex} + \alpha_3 \text{ week} \times \text{sex}$$

are displayed in tables 1 and 2. Notice that these models do not have an id effect since we are analyzing mean weights and head circumferences.

	Estimate	Std. Error	t value	P(> t)
(Intercept)	3365.70	174.91	19.243	4.3e-05 ***
week	223.54	19.40	11.520	0.000324 ***
sex (boy)	-185.37	247.35	-0.749	0.495267
week:sex (boy)	32.75	27.44	1.194	0.298597
<hr/>				
Signif. codes:	0 '***'	0.001 '**'	0.01 '*'	0.05 '.' 0.1 ' ' 1

Table 1: Estimated coefficients for weight.




Figure 19: Mean weight in [g] per week in blue for boys and pink for girls.




Figure 20: Mean head circumference in [cm] per week in blue for boys and pink for girls.

	Estimate	Std. Error	t value	P(> t)
(Intercept)	34.07687	0.66572	51.188	8.72e-07 ***
week	0.36628	0.07385	4.959	0.00771 **
sex (boy)	0.12198	0.94147	0.130	0.90317
week:sex (boy)	0.08387	0.10445	0.803	0.46700
—				
Signif. codes:	0 '***'	0.001 '**'	0.01 '*'	0.05 '.' 0.1 ' ' 1

Table 2: Estimated coefficients for head circumference.

4.1 Could baby size and weight explain the difference in concentration?

We established that, in our study, baby weight and head circumference do not depend on gender, so these variables do not explain the gender differences in concentration. Nonetheless, it could be that bigger babies have higher concentrations - in spite of their gender. Consider the following models for the concentration:

$$AA = \alpha_0 + \alpha_1 \textit{ week} + \alpha_2 \textit{ sex} + \alpha_3 \textit{ weight} + \alpha_{id} \quad (6)$$

and

$$AA = \alpha_0 + \alpha_1 \textit{ week} + \alpha_2 \textit{ sex} + \alpha_3 \textit{ head} + \alpha_{id}. \quad (7)$$

In these models I study the effect of weight and head circumference while controlling for two known effects: sex and week.

References

- [1] Longitudinal Variation of Amino Acid Levels in Human Milk and Their Associations with Infant Gender, Joris H. J. Van Sadelhoff, Bert J. M. Van de Heijning, Bernd Stahl, et al., *Nutrients* 2018, 10(9), 1233; <https://doi.org/10.3390/nu10091233>