

MATH 425A HW5, DUE 09/27/2022, 6PM

JUAN SERRATOS

CHAPTER 3. §2.

Exercise 0.1 (2.5.). Finish the proof of Proposition 2.6., by proving that

$$\text{Int}_Y(U) \cap \text{Int}_X(Y) \subseteq \text{Int}_X(U) \quad (U \subseteq Y \subseteq X)$$

Proof. Suppose $U \subseteq Y \subseteq X$, where (X, d) is a metric space. Now take $t \in \text{Int}_Y(U) \cap \text{Int}_X(Y)$. Then we have that there is some $t \in U \cap Y$ such that $r_\alpha > 0$ with $B_Y(t, r_\alpha) \subseteq U$, and there is some r_β such that $B_X(t, r_\beta) \subseteq Y$. Using ball notation, we know that $t \in U \cap Y \cap X$, and that $B_X(t, r_\beta) = \{y \in X : d(t, y) < r_\beta\}$ and $B_Y(t, r_\alpha) = \{q \in Y : d(t, q) < r_\alpha\}$. So we need to show that $t \in \text{Int}_X(U)$, i.e. $t \in X$ and there is some $r_\gamma > 0$ such that $B_X(t, r_\gamma) \subseteq U$. \square

Exercise 0.2 (2.6.).

Exercise 0.3 (2.7.). Let (X, d) be a metric space, and let U be a subset of X . Use Proposition 2.9 to prove that $\text{Int}_X(U)$ is open in X .

Proof. We want to show that $\text{Int}_X(U) \subseteq \text{Int}_X(\text{Int}_X(U))$. So let $t \in \text{Int}_X(U)$. Then $t \in U$ with $B(t, r) \subseteq U$ for some $r > 0$. Now let $y \in B(t, r)$, but as open balls are open then $y \in \text{Int}(B(t, r))$, and so we can find another open ball $B(t, \epsilon) \subseteq B(t, r)$. All together, we have that $B(t, \epsilon) \subseteq B(t, r) \subseteq U$. Thus if we have some point $y \in B(t, r)$ then $y \in \text{Int}(U)$, and hence $B(t, r) \subseteq \text{Int}(U)$. Therefore $\text{Int}(U)$ is an open set. \square

Exercise 0.4 (2.8.).

Email address: jserrato@usc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90007