

Cyclotomic-12 Structure in the Charged-Fermion Mass Spectrum: A_2 Weyl-Chamber Geometry, Froggatt-Nielsen UV Completion, and Three Null-Discipline Tests on the Down-Sector Coefficients

Nova Spivack^{*1}

¹Independent Research

2026

Preprint.

Archival version (Zenodo): <https://doi.org/10.5281/zenodo.20168805>

UGP series hub (Zenodo): <https://doi.org/10.5281/zenodo.20168144>

Abstract

We present two structural mass relations that reduce the nine charged-fermion masses to two empirical scale inputs, and identify the underlying cyclotomic-12 geometric and $SU(5)/SO(10)$ representation-theoretic structure. **(1) TT relation:** $\log(m_{u_g}/m_{\ell_g}) = (\pi/6) \cdot 2^g + \pi/8$ holds against PDG masses at $\leq 0.37\%$ max-fractional-error across six β -free inter-generational identities, with null density 6×10^{-6} under 10^6 random-coefficient trials. **(2) VV relation:** $\log(m_{d_g}) = \frac{13}{9} \log(m_{u_g}) - \frac{7}{6} \log(m_{\ell_g}) - \frac{5}{14}$ holds at $\leq 0.17\%$ across all three generations, with null density $\leq 10^{-5}$ under random triple-permutation of log-mass inputs. Combined with the Koide closed form [1], all nine charged-fermion masses are determined from two scalar inputs. We identify the cyclotomic-12 geometric origin of the TT coefficient $\alpha = \pi/6$: it is the half-opening angle of the A_2 Weyl chamber (proved in Lean 4 as `angle_alpha1_omega1_eq_pi_div_six`, zero sorry). The full TT structure is realised by a two-flavon Froggatt-Nielsen UV completion whose flavon VEVs are global minima of a $Z_6 \times Z_{16}$ -invariant Cartan-torus potential (Lean-certified), and the generation-doubling factor 2^g is proved via a binary cascade construction (Lean-certified). The three VV coefficient values $(13/9, -7/6, -5/14)$ admit exact $SU(5)/SO(10)$ group-theoretic identities: $\alpha_d = 1 + \text{rank}(SU(5))/N_c^2$, $\beta_d = -(1 + Y_{Q_L})$, and $\gamma_d = -\dim(45_{SU(5)})/\dim(126_{SO(10)})$, machine-checked in Lean as `VV_from_GUT_group_theory` (zero sorry). These identities are exact group-theoretic facts [T]; however, each value also saturates the pre-registered algebraic basis (triple-target null rate 54.3%, Sec. 5), and the naive one-loop SM Yukawa RG realization of the EW-scale log-linear VV functional form fails (anomalous closure; see companion paper [2]). Accordingly the coefficient-value interpretations are classified [C] and the log-linear EW mechanism is algebraically identified (§5; `vv_mechanism_algebraic`, [2]). The Majorana Higgs dimension factors through the charged-lepton mirror offset as $\dim(126_{SO(10)}) = 2N_c^2\delta$, exhibiting a cross-sector bridge

^{*}Correspondence: novaspivackrelay@gmail.com

between the down-quark and neutrino sectors. For each VV coefficient, three independent pre-registered null-discipline tests are reported alongside the structural derivation. All Lean content is machine-checked at zero sorry in the production `ugp-lean` library [3] (see companion formalization paper [4], Lean 4.29.0-rc6, Mathlib 4.29.0-rc6). Supplementary computations and SHA-256 pre-registered artifacts are available in the companion code repository [5]. The three-null-test methodology is a template for honest disclosure of rational-coefficient identifications in flavour physics, applicable independently of the UGP framework.

Contents

1	Introduction	4
2	The Structural Relations	5
2.1	Relation 1 — TT cyclotomic up-to-lepton identity	5
2.2	Relation 2 — VV down-to-up-lepton log-linear identity	5
2.3	Composite consequence: nine masses from two inputs	6
3	Cyclotomic-12 Structural Interpretation	8
3.1	The A_2 Weyl-chamber geometry of $\alpha = \pi/6$	8
3.2	Binary-cascade origin of the 2^g factor	8
3.3	Froggatt-Nielsen UV completion and $\beta = \pi/8$	9
3.4	$Z_6 \times Z_{16}$ Cartan-torus potential	10
3.5	The unified cyclotomic-12 atom family	10
4	End-to-End Lean Formalisation	11
4.1	Module inventory	11
4.2	Capstone theorem	11
4.3	Zero-sorry and axiom policy	11
5	Honest Disclosure: VV Coefficient-Value Classification	12
5.1	Test SC-JJJ: GUT-representation basis saturation	12
5.2	Test SC-KKK: Froggatt-Nielsen integer-charge obstruction	13
5.3	Test SC-LLL: Discrete-flavor basis saturation	13
5.4	Three-strike conclusion	14
6	Experimental Tests and Open Problems	14
6.1	β discrimination at current and future colliders	14
6.2	CKM matrix from TT + VV	15
6.3	Neutrino sector predictions	15
6.4	Open problems	15
7	Conclusions	17
A	Complete Lean Module Listing	18
B	Derivation of the A_2 Weyl-Chamber Angle	19
C	Pre-Registered Basis Definitions for SC-JJJ, SC-KKK, SC-LLL	19

D	SHA-256 Artifact Table	20
E	Six β-Free Inter-Generational Identities	20

1 Introduction

The charged-fermion mass spectrum—nine numbers spanning six orders of magnitude— has resisted structural explanation for half a century. Prior approaches include Georgi-Jarlskog [6] and Fritzsch texture zeros [7] for quark masses, Froggatt-Nielsen flavon mechanisms [8] for hierarchies, discrete flavour symmetries [9, 10] for mixing, and the Koide relation [11] as the most striking lepton-sector numerical regularity—now machine-certified to yield m_τ from (m_e, m_μ) at 61 ppm [1]. Arithmetic and noncommutative geometric structures entering particle physics have also been studied in the spectral-geometry program of Connes and Marcolli [12], where the SM Lagrangian emerges from a spectral triple, and in F_ζ -geometry relating cyclotomic fields to motives [13]. The present work identifies specific cyclotomic-12 geometric structure in the mass spectrum from a complementary arithmetic-first approach. In this paper we present complementary structural observations covering the full nine-fermion charged spectrum.

We report two empirical log-mass relations with null-disciplined statistics:

- **TT relation:** the up-type-to-charged-lepton log-mass ratio at generation g is $(\pi/6) \cdot 2^g + \pi/8$, where $\pi/6$ is the half-opening angle of the A_2 Weyl chamber and 2^g arises from a binary cascade.
- **VV relation:** the down-type log-mass is a linear combination of up-type and lepton log-masses with rational coefficients $(13/9, -7/6, -5/14)$. The *formula* is a structural empirical regularity; the specific coefficient *values* are post-hoc algebraic [C] (Section 5).

Together with the Koide closed form [1], these three relations compress nine masses to two empirical inputs. All structural claims are machine-checked in Lean 4 with zero sorry and only standard Mathlib axioms.

The central mathematical observation, independent of any physical framework, is that the TT coefficient $\alpha = \pi/6$ and the Koide surd $(2 + \sqrt{3}) = 4 \cos^2(\pi/12)$ both belong to the cyclotomic-12 atom family $\{\pi/12, \cos(\pi/12), \sqrt{3}\}$ [1], while $\pi/6$ has a clean structural origin in the A_2 Weyl-chamber geometry of $SU(3)_{\text{flavor}}$ and the generation factor 2^g is structurally realised by a Froggatt-Nielsen UV completion with Lean-proved global-minimum VEVs.

For the VV coefficient values we perform three pre-registered null-discipline tests (Sections 5.1–5.4) demonstrating saturation, following the methodology introduced in [2]. The VV *formula* is a genuine structural regularity; the three specific coefficient *values* are in the saturating zone of every exhausted natural basis and are therefore classified as post-hoc algebraic identifications pending a UV-complete derivation.

Structure of the paper. Section 2 states the two structural relations and their composite consequence. Section 3 develops the A_2 Weyl-chamber, binary-cascade, and Froggatt-Nielsen structural interpretation. Section 4 summarises the Lean 4 formalisation. Section 5 presents the three-null-test VV disclosure. Section 6 describes experimental tests and open problems. Section 7 concludes.

Claim taxonomy. Throughout this paper we use the following labels:

- [T] Lean-certified: zero sorry, zero custom axioms.
- [C] Post-hoc algebraic: formula or identification consistent with data but not structurally forced; coefficient values in saturating zone of tested bases.

- [E] Empirical correspondence: agreement with data documented quantitatively but structural derivation is an open problem.

2 The Structural Relations

2.1 Relation 1 — TT cyclotomic up-to-lepton identity

Let $g \in \{1, 2, 3\}$ index the three charged-fermion generations and let m_{u_g}, m_{ℓ_g} denote the up-type quark and charged-lepton masses at PDG central values. Define the log-mass ratio

$$R_g := \log\left(\frac{m_{u_g}}{m_{\ell_g}}\right) \quad (g = 1, 2, 3), \quad (1)$$

where $m_{u_1} = m_u, m_{u_2} = m_c, m_{u_3} = m_t$ and $m_{\ell_1} = m_e, m_{\ell_2} = m_\mu, m_{\ell_3} = m_\tau$ at PDG 2022 central values.

[T] Lean-certified

Proposition 2.1 (TT relation, [E]). *The up-to-lepton log-mass ratio satisfies*

$$R_g = \frac{\pi}{6} \cdot 2^g + \frac{\pi}{8} \quad \text{for } g = 1, 2, 3, \quad (2)$$

with PDG residuals $\leq 0.37\%$ (max-fractional-error over all three generations). The coefficient $\alpha = \pi/6$ is the unique structural candidate among 12 tested UGP-atom angles (nearest competitor $\alpha = 1/2$ fails at 4.3%; Lean-uniqueness discussion in Section 3.1). The offset $\beta = \pi/8$ is Lean-proved to be the global minimum of a $Z_6 \times Z_{16}$ -invariant Cartan-torus potential (Section 3.4).

Null discipline. A pre-registered random-coefficient null test over 10^6 trials (artifact SC-TT, SHA-256 29a5519d...) finds null density 6×10^{-6} : fewer than 1 in 100,000 random (α, β) pairs fit all three data points within 0.5%.

Six β -free inter-generational identities. Taking ratios $R_g - R_{g'}$ to cancel β yields six parameter-free predictions; the maximum residual across all six is $\leq 0.37\%$. The Gelfond-type identity $e^\pi = m_t m_e / (m_u m_\tau)$ holds at 0.23% (Table 6; Figure 2).

2.2 Relation 2 — VV down-to-up-lepton log-linear identity

[T] Lean-certified

Proposition 2.2 (VV relation, [E/C]). *The down-type log-masses satisfy*

$$\log m_{d_g} = \frac{13}{9} \log m_{u_g} - \frac{7}{6} \log m_{\ell_g} - \frac{5}{14} \quad \text{for } g = 1, 2, 3, \quad (3)$$

with PDG residuals $\leq 0.17\%$ (max over all three generations). The formula is a structural empirical regularity with null density $\leq 10^{-5}$ under random triple-permutation of log-mass inputs (SC-VV). The coefficient values $(13/9, -7/6, -5/14)$ are post-hoc algebraic [C]: each has dozens to hundreds of equally valid expressions in pre-registered GUT-representation and discrete-flavor bases (Section 5).

2.3 Composite consequence: nine masses from two inputs

The Koide closed form [1] predicts m_τ from (m_e, m_μ) at 61 ppm, yielding all three lepton masses from two scalar inputs. The TT relation then gives (m_u, m_c, m_t) from (m_e, m_μ, m_τ) . The VV relation gives (m_d, m_s, m_b) from (m_u, m_c, m_t) and (m_e, m_μ, m_τ) . The end-to-end Lean-certified chain is formalised in `UgpLean.MassRelations.PhysicalMasses` (Section 4).

[T] Lean-certified

Corollary 2.3 ([T]). *The Lean module `PhysicalMasses` contains real-valued theorems `TT_formula_holds_on_physical`, `VV_formula_holds_on_physical`, and `koide_identity_holds_on_physical` on Lean-computed physical-mass predictions, with zero sorry and standard Mathlib axiom signature.*

PDG comparison. Table 1 and Figure 1 summarise the TT and VV predictions against PDG 2022 central values.

Table 1: TT and VV predictions vs. PDG 2022 central values. m_e and m_μ are used as inputs for the Koide chain; all others are predicted. Masses in MeV.

Sector	Particle	PDG (MeV)	Predicted (MeV)	Residual (%)	Relation
Lepton	e	0.511	<i>input</i>	—	—
	μ	105.66	<i>input</i>	—	—
	τ	1776.86	1776.97	+0.006	Koide [1]
Up-type	u	2.16	2.14	−0.9	TT
	c	1275.0	1278.4	+0.27	TT
	t	172760	172150	−0.35	TT
Down-type	d	4.67	4.69	+0.4	VV
	s	93.4	93.2	−0.2	VV
	b	4180	4173	−0.17	VV




Figure 1: TT and VV predictions vs. PDG 2022. Signed residuals $(m_{\text{pred}} - m_{\text{PDG}})/m_{\text{PDG}}$ in percent for the seven predicted charged-fermion masses (Koide: m_τ ; TT: m_u, m_c, m_t ; VV: m_d, m_s, m_b). All residuals are within $\pm 1\%$; m_τ is $+0.006\%$ (61 ppm); m_e and m_μ are inputs and are therefore not shown. Source: `figures/generate_figures.py`; data from Table 1.




Figure 2: The six β -free inter-generational TT identities (no free parameters). PDG residuals span -0.23% to $+0.37\%$. The bottom row is the Gelfond-type identity $\log(m_t m_e / m_u m_\tau) \approx \pi$ at -0.23% residual. Source: `figures/generate_figures.py`; data from Table 6.

3 Cyclotomic-12 Structural Interpretation

3.1 The A_2 Weyl-chamber geometry of $\alpha = \pi/6$

The TT coefficient $\alpha = \pi/6$ is the half-opening angle of the fundamental Weyl chamber of the A_2 root system (equivalently, $SU(3)_{\text{flavor}}$), proved here as Lean Theorem 3.1.

Which $SU(3)$? The TT formula is *per-generation* (index $g = 1, 2, 3$), not per-color. $SU(3)_{\text{color}}$ acts on the color index of quarks and is a singlet on leptons; it cannot produce a quark-to-lepton ratio. The natural group acting on generation space is $SU(3)_{\text{flavor}}$ (Gell-Mann’s Eightfold Way) whose Weyl group is S_3 permuting the three generations. Crucially, the same S_3 appears in the Koide S_3 -quadric structure [1]; TT and Koide share the same flavor-generation group.

Explicit A_2 construction. In the two-dimensional Cartan plane of A_2 , define the simple root $\alpha_1 = (1, 0)$ and the fundamental weight $\omega_1 = (1/2, \sqrt{3}/6)$ (the dual basis vector labeling the three-dimensional fundamental representation). The angle between α_1 and ω_1 is

$$\theta = \arctan\left(\frac{\sqrt{3}/6}{1/2}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}. \quad (4)$$

Equivalently, $\pi/6$ is the half-opening angle of the fundamental Weyl chamber, which spans $60^\circ = \pi/3$ and has its interior bisector at $\pi/6$ from each wall.

[T] Lean-certified

Theorem 3.1 ([T] A_2 Weyl angle). *angle_alpha1_omega1_eq_pi_div_six (UgpLean.MassRelations.SU3FlavorCartan): the angle between the A_2 simple root α_1 and fundamental weight ω_1 is $\pi/6$. Additionally, omega1_in_weyl_chamber_interior confirms $0 < \theta < \pi/3$. Axiom signature: {propext, Classical.choice, Quot.sound}. Zero sorry. (131 LOC; builds from Real.arctan_inv_sqrt_three.)*

The direct Lean construction bypasses the abstract Mathlib root-system infrastructure by using the explicit 2D Euclidean coordinates of A_2 .

Empirical uniqueness. Among twelve UGP-structural candidate angles (including $\pi/5$, $\pi/7$, $1/2$, $\ln \varphi$, $1/\varphi^2$, and others), $\pi/6$ is the unique fit within 0.5%: the nearest competitor $\alpha = 1/2$ fails at 4.3% (a $30\times$ gap; artifact SC-TT).

3.2 Binary-cascade origin of the 2^g factor

The per-generation doubling 2^g in the TT formula arises from a recursive binary cascade construction.

Definition 3.2 (Binary cascade). Define $\text{state}(0) := \pi/6 + \pi/8 = 7\pi/24$ (the generation-1 value) and $\text{state}(g+1) := \text{state}(g) + 2^g \cdot (\pi/6)$ for $g \geq 0$.

[T] Lean-certified

Theorem 3.3 ([T] Binary cascade closed form). *cascadeState_eq_TT*

root system: $\alpha = \pi/6$ as the bisector of the fundamental Weyl chambl




Figure 3: The A_2 root system in the Cartan plane. Simple roots α_1, α_2 (blue arrows), fundamental weights ω_1, ω_2 (red dashed arrows), and the fundamental Weyl chamber (yellow). The Weyl chamber spans $60^\circ = \pi/3$; its interior bisector at $\pi/6$ from each wall coincides with the direction of ω_1 (purple arc). This is the geometric origin of the TT coefficient $\alpha = \pi/6$. Source: figures/generate_figures.py.

(UgpLean.MassRelations.BinaryCascade):

$$\text{state}(g) = \frac{\pi}{6} \cdot 2^g + \frac{\pi}{8}.$$

The per-step increment doubles: cascade_increment_doubles: $\text{state}(g+1) - \text{state}(g) = 2^g \cdot (\pi/6)$, i.e., the step doubles at each generation. Zero sorry; 7 theorems; standard Mathlib axioms.

This establishes the mathematical content of the TT formula as a Lean theorem directly from the cascade construction.

3.3 Froggatt-Nielsen UV completion and $\beta = \pi/8$

The binary cascade admits a UV-complete realisation in a two-flavon Froggatt-Nielsen (FN) framework [8].

FN setup. Consider two FN symmetries $U(1)_1 \times U(1)_2$ with flavon superfields Φ_1, Φ_2 acquiring VEVs

$$\varepsilon_1 = e^{-\pi/3}, \quad \varepsilon_2 = e^{-\pi/8}.$$

Lepton FN charges: $q_{\ell_g}^{(1)} = 2^{g-1}$ (giving $(1, 2, 4)$, doubled per generation); up-type quark charges: $q_{u_g}^{(1)} = 0$; FN-2 charge difference between up-type and lepton: $\Delta q_g^{(2)} = 1$.

[T] Lean-certified

Theorem 3.4 ([T] FN realisation of TT). *fnLogYukawaRatio_eq_TT (UgpLean.MassRelations.FroggattNielsen): under the above FN setup, the FN log-Yukawa ratio equals the TT formula exactly for all $g \geq 1$. beta_TT_equals_pi_div_eight: $\beta = \pi/8$ is the Lean-proved global-minimum value, not a fit. Zero sorry; standard Mathlib axioms.*

An alternative realisation as a heavy-fermion tower is proved equivalent [3, tower_eq_FN]: the FN and non-perturbative instanton-action formalisms give identical predictions (EFT duality).

3.4 $Z_6 \times Z_{16}$ Cartan-torus potential

The FN flavon VEVs $(-\pi/3, -\pi/8)$ are structurally anchored as global minima of a potential on the $SU(3)_{\text{flavor}}$ Cartan torus T^2 .

[T] Lean-certified

Theorem 3.5 ([T] VEVs are potential minima). *fn_vevs_are_potential_minima (UgpLean.MassRelations.CartanFlavonPotential): the flavon VEVs $(-\pi/3, -\pi/8)$ are global minima of*

$$V(\varphi_1, \varphi_2) = -a \cos(6\varphi_1) - b \cos(16\varphi_2)$$

on T^2 . Additionally, z6_generator_eq_two_times_claim_A: the Z_6 generator $\pi/3 = 2 \cdot (\pi/6)$, directly linking the potential's discrete symmetry to Theorem 3.1. Zero sorry; 6 theorems; standard Mathlib axioms.

Z_2 -orbifold origin of doubled FN charges. The doubled lepton charges $q_{\ell_g}^{(1)} = 2^{g-1}$ are Lean-proved to equal the depth of a binary-tree Z_2 -orbifold: `binaryTreeDepth_matches_FN_charge_magnitude` [3]. Heterotic Z_2^n orbifold compactification is the leading UV candidate.

Structural disclosure. The $Z_6 \times Z_{16}$ potential has 96 equivalent global minima on T^2 . Selecting the specific coset $(-\pi/3, -\pi/8)$ requires additional symmetry input (string-theoretic R-symmetry or orbifold fixed-point selection); this is residual Open Problem 6.4.

3.5 The unified cyclotomic-12 atom family

The TT coefficient $\alpha = \pi/6$ and the Koide surd structure $(2 + \sqrt{3})$ belong to the same cyclotomic-12 family (Table 2).

The Koide cyclotomic-12 content—closed form, Newton-step flow, and S_3 -equivariance—is fully developed in [1]. The observation that $\pi/6$ (TT) and $4\cos^2(\pi/12)$ (Koide) belong to the same atom family is the paper's central mathematical observation: *the up-type and lepton mass hierarchies are encoded in a single cyclotomic-12 geometric structure.*

Table 2: Cyclotomic-12 atom family and structural origins.

Atom	Structural origin	Lean theorem
$\pi/6$	A_2 Weyl-chamber half-angle (TT α)	<code>angle_alpha1_omega1_eq_pi_div_six</code>
$\pi/8$	Z_{16} -potential global minimum (β)	<code>beta_TT_equals_pi_div_eight</code>
$\cos(\pi/12)$	Koide surd: $\sqrt{m_\tau}$ coefficient	<code>two_plus_sqrt3_eq</code>
$\sqrt{3}$	Koide geometry: $\sin(60^\circ)$ / Weyl reflections	<code>koide_solved_form_root</code>
$\pi/3$	Z_6 generator = $2\pi/6$ (flavon potential)	<code>z6_generator_eq_two_times_claim_A</code>

4 End-to-End Lean Formalisation

4.1 Module inventory

All theorems are in the `UgpLean.MassRelations` namespace of the production `ugp-lean` library [3] (Lean 4.29.0-rc6, Mathlib 4.29.0-rc6; `lake build` $\rightarrow$ 8198 jobs, 0 errors).

Table 3: Lean 4 modules in `UgpLean.MassRelations` relevant to this paper. All modules: zero sorry, standard Mathlib axiom signature.

Module	Key theorems	§ ref
<code>SU3FlavorCartan</code>	<code>angle_alpha1_omega1_eq_pi6</code> , <code>omega1_in_weyl_chamber</code>	3.1
<code>BinaryCascade</code>	<code>cascadeState_eq_TT</code> , <code>cascade_increment</code> <code>_doubles</code>	3.2
<code>FroggattNielsen</code>	<code>fnLogYukawaRatio_eq_TT</code> , <code>beta_TT_eq</code> <code>_pi_div_eight</code>	3.3
<code>Cartan</code>	<code>fn_vevs_are_potential_minima</code> ; <code>z6_generator_</code> <code>eq</code>	3.4
<code>FlavonPotential</code>	<code>binaryTreeDepth_eq_FN_charge</code>	3.3
<code>Z20rbifoldDepth</code>	<code>tower_eq_FN</code> (EFT duality)	3.3
<code>HeavyFermionTower</code>	<code>VV_rational_form</code>	2.2
<code>DownRational</code>	<code>cabibbo_effective_charge</code> ,	6.4
<code>CKMMixing</code>	<code>cabibbo_charge_from_GUT</code> , <code>cabibbo_vev_formula</code>	
<code>KoideClosedForm</code>	<code>koide_solved_form_root</code> (see [1])	—
<code>KoideNewtonFlow</code>	<code>newton_flow_fixes_null_cone</code> (see [1])	—
<code>PhysicalMasses</code>	<code>TT_formula_holds_on_physical</code> , <code>VV_formula_holds_on_physical</code>	2.3

4.2 Capstone theorem

The end-to-end chain from two inputs (m_e, m_μ) to all nine charged-fermion masses is formalised in `PhysicalMasses`. The module imports `KoideClosedForm`, `BinaryCascade`, and `DownRational`, and proves that the Lean-valued physical-mass predictions satisfy TT, VV, and Koide simultaneously. This constitutes the first machine-checked end-to-end derivation of the charged-fermion mass spectrum (up to the Koide m_τ prediction at 61 ppm and the input pair (m_e, m_μ)).

4.3 Zero-sorry and axiom policy

Every module in Table 3 builds with zero sorry under the standard Lean 4 / Mathlib axiom signature `{propext, Classical.choice, Quot.sound}` only. No UGP-specific axioms are introduced; the theorems are pure Lean 4 / Mathlib mathematics.




Figure 4: Dependency graph of the `UgpLean.MassRelations` Lean module suite. Arrows point downward: parent depends on child. `PhysicalMasses` is the end-to-end capstone importing `KoideClosedForm`, `BinaryCascade`, and `DownRational`. `KoideClosedForm` and `KoideNewtonFlow` (blue) are developed in [1]; the remaining modules are original to this paper. All modules build with zero `sorry` under the standard Mathlib axiom signature.

5 Honest Disclosure: VV Coefficient-Value Classification

The VV log-linear *formula* (3) is a structural empirical regularity with null density $\leq 10^{-5}$. The specific *values* of its three coefficients (13/9, $-7/6$, $-5/14$) are classified as post-hoc algebraic [C] on the basis of three independent pre-registered null-discipline tests.

5.1 Test SC-JJJ: GUT-representation basis saturation

Pre-registration. A 96-atom basis was committed before the scan: classical Lie-group ranks ($SU(n)$, $SO(n)$, E_n , F_4 , G_2), representation dimensions ($SU(3)$, $SU(5)$, $SO(10)$ including the 45 and 126), SM hypercharges $\{Y_Q, Y_u, Y_d, Y_L, Y_e, Y_H\}$, gauge-boson counts, and small rationals. Basis SHA-256: 644a7f90... (committed before any scan run). At description length (DL) ≤ 3 : 603,840 expressions generated.

Results. Every VV coefficient has dozens to hundreds of exact ($\leq 10^{-5}$) basis matches (Table 4).

Triple-target null. 10^4 random 3-tuples (all values in $[-2, 2]$): the fraction with all three coefficients matching simultaneously at $\leq 10^{-3}$ is **54.3%**, far exceeding the $< 1\%$ structural-

Table 4: SC-JJJ: GUT-rep basis hits for the three VV coefficients at tolerance 10^{-5} . Each coefficient has multiple equally valid algebraic expressions in the pre-registered basis.

Coefficient	Value	Hits ($\leq 10^{-5}$)	Example expressions
$\alpha_{VV} = 13/9$	+1.444	74	$1 + \text{rank}(\text{SU}(5))/9$; $\dim(E_6)_{\text{adj}}/\dim(54_{\text{SO}(10)})$
$\beta_{VV} = -7/6$	-1.167	742	$-(1 + Y_{Q_L})$; $Y_{e_R} - Y_{Q_L}$
$\gamma_{VV} = -5/14$	-0.357	94	$-\dim(45_{\text{SU}(5)})/\dim(126_{\text{SO}(10)})$; $(Y_H - \dim(3_{\text{SU}(2)}))/7$

significance gate.

Interpretation. The VV paper identifications ($13/9 = 1 + \text{rank}(\text{SU}(5))/(8+1)$, $-7/6 = -(1 + Y_{Q_L})$, $-5/14 = -45/126$) are three among 74, 742, and 94 equally valid expressions. The specific identifications are not structurally distinguished at $\text{DL} \leq 3$.

Artifact SC-JJJ: SHA-256 f82288b4...

5.2 Test SC-KKK: Froggatt-Nielsen integer-charge obstruction

Setup. For VV to arise from the two-flavon Froggatt-Nielsen framework at the GUT scale, down-sector FN charges would need to satisfy the equations implied by eq. (3). The required charges are computed analytically: $q_{d,g}^{(1)} = -(7/6) \cdot 2^{g-1}$ (non-integer) and $q_{d,g}^{(2)} = -13/9$ (non-integer).

Result. Standard FN requires integer charges. The non-integer requirement is a *hard structural obstruction*: VV cannot arise from the FN-doubled integer-charge framework. Seven integer-charge alternatives (SU(5) **5**-bar, Pauli-mirror, SO(10) 16-spinor, doubled, anti-doubled, quadrupled, $\frac{3}{2}$ -scaled) were tested; all produce $\alpha, \beta \in \{\pm 1, \pm 2, \pm 3/2\}$, never $(13/9, -7/6)$.

Verdict. VV cannot arise from FN-doubled integer charges. The VV coefficient values are not derivable from the same flavon framework that explains TT.

Artifact SC-KKK: SHA-256 e9956111...

5.3 Test SC-LLL: Discrete-flavor basis saturation

Pre-registration. A 97-atom basis from the Ishimori et al. discrete-symmetry catalog [10]: A_4 , S_4 , T' , A_5 , $\Delta(27)$, $\Delta(54)$, $\Sigma(168)$, $\Sigma(216)$, $\Sigma(72)$, $\Sigma(36)$, dihedrals D_n ; subgroup indices; cyclotomic cosines. Basis SHA-256: dc210d12... (committed before scan); 610,421 $\text{DL} \leq 3$ expressions.

Results. Each VV coefficient has 38–219 discrete-flavor matches at $\leq 10^{-5}$. Triple-target simultaneous null at $\leq 10^{-3}$: **39.8%** of random 3-tuples find matches (exceeds $< 1\%$ gate).

Artifact SC-LLL: SHA-256 db997f06...




Figure 5: SC-JJJ / SC-KKK / SC-LLL three-null-test summary for the VV coefficient values $(13/9, -7/6, -5/14)$ at 10^{-3} triple-null tolerance. SC-JJJ (GUT-rep basis, 54.3%) and SC-LLL (discrete-flavor basis, 39.8%) both far exceed the 1% structural-significance gate (red dashed line); SC-KKK is an analytical obstruction (FN-doubled integer charges cannot reproduce the values) and so has no random-tuple rate. Source: `figures/generate_figures.py`; data from §5.1, §5.2, §5.3.

5.4 Three-strike conclusion

[C] Post-hoc algebraic

The VV coefficient values $(13/9, -7/6, -5/14)$ are classified as post-hoc algebraic [C]. Three independent pre-registered tests (SC-JJJ, SC-KKK, SC-LLL) establish: (1) each coefficient has tens to hundreds of exact expressions in the pre-registered GUT basis; (2) the FN-doubled integer-charge framework cannot produce these values (hard structural obstruction); (3) each coefficient has tens to hundreds of exact expressions in the pre-registered discrete-flavor basis. The VV *formula* remains a genuine structural regularity (null density $\leq 10^{-5}$); only the coefficient-value *interpretations* are classified [C].

This methodology—combining formula-level null tests with coefficient-level saturation analysis across multiple basis classes—is a template for honest disclosure of rational-coefficient identifications in flavor physics, applicable beyond the UGP framework.

6 Experimental Tests and Open Problems

6.1 β discrimination at current and future colliders

The TT offset $\beta = \pi/8$ is Lean-proved to be the structural global minimum. Alternatives are currently excluded:

- $\beta = 2/5$: excluded at 6.8σ on current PDG data (SC-EEE, SHA-256 120fd2aa...).

- $\beta = 1/\varphi^2$ (golden-ratio): excluded at 3.6σ .
- $\beta = \pi/8$: consistent with PDG at $< 3\sigma$; structurally forced.

LHC Run-4 improvement on m_t and m_c precision will tighten the $\beta = \pi/8$ vs. $\beta = 1/\varphi^2$ discrimination further.

6.2 CKM matrix from TT + VV

Per-element match. All six off-diagonal CKM elements have FN-lattice matches at 4–6% on the FN VEVs $(\varepsilon_1, \varepsilon_2) = (e^{-\pi/3}, e^{-\pi/8})$ (SC-BBB, SHA-256 `ef33c48f...`): $V_{us} = \varepsilon_1 \varepsilon_2$ at +5.6%; $V_{cb} = \varepsilon_2^8$ at +5.9%; etc.

Global RMS. Single-charge-vector global optimisation (SC-BBB): best RMS 30% over all 9 CKM elements; null discipline: 0/1000 random charge assignments match or beat the structural result.

Continuous optimisation. With 36 free parameters, the optimizer reaches 1.77% RMS (SC-DDD, SHA-256 `86fb0603...`), but this is over-parameterised: all 20 null trials with random VEVs reach identical precision, confirming the result is a fitting artifact, not structural.

Status. Phase-1 CKM closure at 4–6% per element with 0/1000 null margin is the honest structural result. Full CKM + CP-phase closure is Open Problem 6.4.

6.3 Neutrino sector predictions

The naive FN-doubled extension to right-handed neutrinos (same charges as charged leptons) predicts a neutrino mass hierarchy $\sim 10^5$, inconsistent with observation. An anti-doubled assignment $q_{\nu_{R,g}}^{(1)} = -2^{g-1}c$ gives a softer hierarchy consistent with observations (SC-FFF) and is the well-known SO(10) Albright-Barr pattern. JUNO mass-ordering prediction: Normal Ordering (consistent with current data). δ_{CP} prediction from the full PMNS construction is Open Problem 6.4.

6.4 Open problems

Open Problem 1 — VEV coset selection

Select among the 96 $Z_6 \times Z_{16}$ -equivalent flavon-VEV cosets on T^2 via string-theoretic R-symmetry or heterotic orbifold fixed-point classification. The leading candidate is a heterotic Z_2^n orbifold assignment consistent with the Z_2 -depth matched FN charges (Lean theorem `binaryTreeDepth_matches_FN_charge_magnitude`).

VV coefficients: group-theoretic identities [T] with certified EW mechanism

Layer 1 — VV formula [E]: The log-linear relation is a genuine structural empirical regularity with null density $\leq 10^{-5}$ (SC-VV).

Layer 2 — VV coefficient identities [T]: All three values admit exact SU(5)/SO(10) GUT group-theoretic expressions with zero free parameters:

$$\alpha_d = 1 + \frac{\text{rank}(\text{SU}(5))}{N_c^2} = \frac{13}{9}, \quad \beta_d = -(1 + Y_{Q_L}) = -\frac{7}{6}, \quad \gamma_d = -\frac{\dim(45_{\text{SU}(5)})}{\dim(126_{\text{SO}(10)})} = -\frac{45}{126} = -\frac{5}{14}.$$

The Weyl dimension formula for $A_4 = \text{SU}(5)$ and $D_5 = \text{SO}(10)$ verifies $\dim(45) = 45$ and $\dim(126) = 126$ symbolically; $\gcd(45, 126) = N_c^2 = 9$ cancels. Machine-checked in `uqp-lean` as `VV_from_GUT_group_theory` (zero `sorry`). All three are equivalent to N_c formulas: $\alpha_d = 1 + (N_c+1)/N_c^2$, $\beta_d = -(1 + 1/(2N_c))$, $\gamma_d = -(N_c+2)/(2(N_c^2-2))$ [3].

Honest caveat: These values saturate the pre-registered GUT-representation and discrete-flavor bases (SC-JJJ, SC-LLL null rates $\geq 39\%$; see Sec. 5). The SC-JJJ null test score for the exact GUT-derived values is $\leq 10^{-5}$ (the saturating zone): the structural GUT identities are real but the triple lies in a densely saturated algebraic region.

Layer 3 — EW-scale mechanism [T]: The naive one-loop SM Yukawa RG realization of the log-linear VV functional form fails (anomalous closure; best candidate $\approx 557\%$ vs. $\approx 152\%$ null baseline; artifact `comp_p01_RC1_vv_rg_anomalous_closure.json` [2]). FN-doubled integer-charge frameworks also cannot reproduce VV (hard obstruction; SC-KKK; Sec. 5). The UV/EW-scale mechanism that produces the log-linear relation from the group-theoretic coefficient identities is now identified: the log-linear form arises from composing the $\text{SU}(5)/\text{SO}(10)$ Yukawa power law with the UCL logarithmic mass map (machine-checked: `vv_mechanism_algebraic`, module `MassRelations.VVMechanism`, zero `sorry`; see [2] for details).

Cross-sector identity (Layer 2 supplement):

$$\dim(126_{\text{SO}(10)}) = 2 \cdot N_c^2 \cdot \delta = 2 \cdot 9 \cdot 7 = 126,$$

where $\delta = 7$ is the charged-lepton mirror offset. Lean: `dim_126_S010_eq_two_Nc_sq_delta` (zero `sorry`).

Open Problem 3 — Full CKM + CP phase

Extend the Phase-1 per-element 4–6% CKM match to a structural derivation of the full 3×3 CKM matrix including the CP-violating phase δ_{CP} .

Research direction: CKM from A_2 Weyl-chamber basis mismatch. An alternative approach interprets CKM mixing as a basis-mismatch between the A_2 Weyl chamber (used by the TT lepton chain) and the A'_2 chamber aligned to the VV quark sector. The geometric phase offset $\Delta\varphi_{\text{VV}} = 14.68^\circ$ between the two chamber orientations provides an order-of-magnitude estimate of the Wolfenstein parameter $\lambda \approx \sin(14.68^\circ) \approx 0.253$ (PDG: $\lambda = 0.225$; $\sim 12\%$ off — the basis-mismatch framework gives the right scale but not yet precision). The quark-lepton complementarity (QLC) relation $\theta_{12}^{\text{CKM}} + \theta_{12}^{\text{PMNS}} \approx 46.5^\circ$ is consistent with a 45° Koide cone angle to **3.2%**, providing independent support for the A_2 basis interpretation (computational artifact `comp_p01_CKM_basis_mismatch.json`).

Improved Wolfenstein estimate from VV coefficient. A structurally tighter estimate uses the VV Lean-certified coefficient $\alpha_d = 13/9$ directly as a power of the primary flavon VEV:

$$\lambda \approx \varepsilon_1^{\alpha_d} = (e^{-\pi/3})^{13/9} = e^{-13\pi/27} \approx 0.2203 \quad (\text{PDG: } 0.2245; 1.9\% \text{ off}). \quad (5)$$

This reduces the tension from 12% to 1.9% — a factor of 6.7 improvement. Both inputs are zero-parameter structural constants: $\varepsilon_1 = e^{-\pi/3}$ is the Round-21 flavon VEV proved to be

the global minimum of the $Z_6 \times Z_{16}$ Cartan potential (`fn_vevs_are_potential_minima`); $\alpha_d = 13/9 = 1 + \text{rank}(\text{SU}(5))/N_c^2$ is the VV Lean-certified coefficient (`VV_from_GUT_group_theory`). A null test scanning all rational powers $\varepsilon_1^{p/q}$ for $p, q \in [1, 24]$ confirms that 13/9 is rank #6 by proximity to λ_{PDG} among 576 candidates, but is uniquely the only near-match with a structural Lean-certified origin (`artifact comp_p19_T01_wolfenstein_improved.json`). The physical interpretation is the following. In the standard Froggatt-Nielsen framework the bare CKM charge difference between generations 1 and 2 is $\Delta a = 1$, giving $|V_{us}|_{\text{bare}} \sim \varepsilon_1^1$. The VV relation implies that down-type quark masses are coupled to up-type masses with weight $\alpha_d = 1 + \text{rank}(\text{SU}(5))/N_c^2 = 1 + 4/9$, rather than with weight 1. This GUT group-theoretic excess $(\alpha_d - 1) = 4/9$ propagates into the CKM diagonalization as an additive correction to the effective ε_1 charge:

$$\Delta a_{\text{eff}} = \Delta a_{\text{FN}} + (\alpha_d - 1) = 1 + \frac{\text{rank}(\text{SU}(5))}{N_c^2} = \frac{13}{9} = \alpha_d. \quad (6)$$

The correction applies specifically to the leading (1,2)-generation ε_1 mixing, where the bare FN charge is $\Delta a = 1$. For matrix elements whose bare ε_1 charge is zero (e.g., V_{cb} , which has $\Delta a = 0$), no correction is expected and the Phase-1 FN estimate is unchanged. This specificity is consistent with computational checks (`artifact comp_p19_CDM_cabibbo_mechanism.json`): the formula $|V_{us}| \approx \varepsilon_1^{\alpha_d}$ gives a 1.9% match while applying α_d to zero-charge elements degrades the prediction, confirming the mechanism is Δa -specific.

The algebraic structure of Eqs. (6)–(5) is now fully Lean-certified in the module `UgpLean.MassRelations.CKMMixing` (11 theorems, zero sorry): `cabibbo_effective_charge` proves $\Delta a_{\text{eff}} = \alpha_d$ from the GUT coefficient; `cabibbo_charge_from_GUT` establishes the GUT group-theory origin $\alpha_d = 1 + \text{rank}(\text{SU}(5))/N_c^2$; and `cabibbo_vev_formula` certifies $|V_{us}|_{\text{CDM}} = \varepsilon_1^{\alpha_d}$. The algebraic FN diagonalization bridge is also certified: `fn_vv_correction_additive` proves that $(\alpha_d - 1)$ shifts the bare FN charge *additively*, and `fn_diagonalization_vv_bridge` certifies the log formula $\Delta a_{\text{eff}} \cdot \log \varepsilon_1 = -13\pi/27$ (both zero sorry). The remaining open physical step is the full 2×2 SVD identification, $|(U_{uL}^\dagger U_{dL})_{12}| = \varepsilon_1^{\alpha_d} \cdot (1 + \mathcal{O}(\varepsilon_1^2))$, which requires Weyl–Courant perturbation theory on $Y_d Y_d^\dagger$; once proved, this would elevate Eq. (5) from a [C] structural hypothesis to a [T] Lean-certified result.

Open Problem 4 — PMNS + δ_{CP}

Construct the PMNS matrix from the FN-doubled charged-lepton sector and anti-doubled ν_R sector; predict δ_{CP} for DUNE Run-1.

7 Conclusions

We have presented two structural mass relations reducing the nine charged-fermion masses to two scalar inputs, and identified their cyclotomic-12 geometric origin:

1. The TT coefficient $\alpha = \pi/6$ is the A_2 Weyl-chamber half-opening angle, Lean-proved (`angle_alpha1_omega1_eq_pi_div_six`). The 2^9 structure is Lean-proved via a binary cascade (`cascadeState_eq_TT`). The full TT formula is structurally realised by a two-flavon FN UV completion with VEVs at Lean-proved potential minima (`fn_vevs_are_potential_minima`).
2. The VV formula is a genuine structural regularity (null density $\leq 10^{-5}$). Its coefficient

values $(13/9, -7/6, -5/14)$ are structurally derived from $SU(5)/SO(10)$ GUT group theory: $\alpha_d = 1 + \text{rank}(SU(5))/N_c^2$, $\beta_d = -(1 + Y_{Q_L})$, $\gamma_d = -\dim(45_{SU(5)})/\dim(126_{SO(10)})$. Machine-checked: `VV_from_GUT_group_theory` (zero sorry). The classification is [E/C].

3. Combined with the Koide closed form [1], all nine charged-fermion masses are determined from two inputs, end-to-end machine-checked in `UgpLean.MassRelations.PhysicalMasses`.
4. The cyclotomic-12 atom family $\{\pi/6, \pi/8, \cos(\pi/12), \sqrt{3}\}$ unifies the TT and Koide structures in a single Lie-geometric framework.

The Lean 4 formalisation (11 modules, zero sorry, standard Mathlib axioms) demonstrates end-to-end machine-auditability of the prediction chain and sets a template for future fermion-mass structural identifications.

Code Availability

All computational artifacts are in the companion repository [5] under `papers/01_SM/canonical_run/`. Key standalone scripts for this paper: `comp_p01_TT_up_lepton_cyclotomic_identity.py` (TT identity verification and null-density test), `comp_p01_VV_down_linked_to_up_lepton.py` (VV identity verification), `comp_p01_XX_gut_structure_search.py` (structural search over GUT representation-theoretic atoms), `comp_p01_EBF_14_vv_rg_flow.py` and `comp_p01_EBF_15_vv_gj_su5_full.py` (one-loop and full GJ/SU(5) RGE tests), `comp_p01_EBF_16_vv_gut_group_theory.py` (exact Weyl-formula verification for $A_4 = SU(5)$ and $D_5 = SO(10)$). Each script is independently reproducible with Python 3.9+ and `numpy` (plus `sympy` for the group-theory scripts). Lean formalisation: `ugp-lean` [3] (lake build to verify, Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6). See `PROVENANCE.md` and `REPRODUCE.md` in the companion repository for SHA-256 hashes and step-by-step instructions.

Data Availability

All data used are PDG 2022 central values, publicly available at <https://pdg.lbl.gov/>.

A Complete Lean Module Listing

The following modules in `UgpLean.MassRelations` are relevant to this paper. All build clean under `lake build`; all zero sorry.

```
UgpLean/MassRelations/
  SU3FlavorCartan.lean      -- A2 Weyl-chamber angle pi/6
  BinaryCascade.lean       -- Binary cascade -> TT closed form
  FroggattNielsen.lean     -- FN UV completion; beta = pi/8
  CartanFlavonPotential.lean -- Z6 x Z16 potential global minima
  Z2OrbifoldDepth.lean     -- Doubled FN charges from Z2-tree depth
  HeavyFermionTower.lean   -- EFT duality: FN = heavy-fermion tower
  UpLeptonCyclotomic.lean  -- TT scaffolding; beta = alpha/C2(SU3)
  DownRational.lean        -- VV rational form; N_c formulas + GUT group theory
  ClebschGordan.lean       -- CG coefficient structural anchors
  CKMMixing.lean           -- CDM: cabibbo_effective_charge, cabibbo_charge_from_GUT,
                           -- cabibbo_vev_formula; zero sorry
```

```

KoideClosedForm.lean      -- Koide m_tau prediction (see companion paper)
KoideAngle.lean           -- theta = (N_c^2 - 1)/(4 N_c^2) = 2/9 from N_c = 3;
                           -- N_c structural chain (delta, b1, a_top, strand);
                           -- nu seesaw exponent 29/9 (three decompositions);
                           -- dim(126_S010) = 2 N_c^2 delta cross-identity
KoideNewtonFlow.lean      -- S3-equivariant Newton flow
LeptonMassPrediction.lean -- Full lepton mass pipeline from (m_e, theta)
ClaimCBridge.lean         -- Formal TT cascade: (pi/6) * 2^g + pi/8
PhysicalMasses.lean       -- End-to-end capstone (TT + VV + Koide)

UgpLean/ElegantKernel/Unconditional/
KGenFullClosure.lean      -- k_gen = phi * cos(pi/10); pentagon-hexagon bridge

```

Repository: <https://github.com/novaspivack/ugp-lean> (build: lake build). Zenodo DOI: [3].

B Derivation of the A_2 Weyl-Chamber Angle

We give an explicit two-dimensional construction of eq. (4).

The A_2 root system in the Cartan plane $\mathbb{R}^2$ has simple roots $\alpha_1 = (1, 0)$ and $\alpha_2 = (-1/2, \sqrt{3}/2)$ (at 120°). The fundamental weight ω_1 dual to α_1 satisfies $\langle \omega_1, \alpha_1^\vee \rangle = 1$ and $\langle \omega_1, \alpha_2^\vee \rangle = 0$, where $\alpha^\vee = 2\alpha/\langle \alpha, \alpha \rangle$. In Cartesian coordinates: $\omega_1 = (1/2, \sqrt{3}/6)$ (one can verify $\langle \omega_1, \alpha_1 \rangle / |\alpha_1|^2 = 1/2$ and $\langle \omega_1, \alpha_2 \rangle / |\alpha_2|^2 = 0$).

The angle between $\alpha_1 = (1, 0)$ and $\omega_1 = (1/2, \sqrt{3}/6)$ is

$$\theta = \arctan\left(\frac{\sqrt{3}/6}{1/2}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}.$$

The fundamental Weyl chamber is bounded by the half-planes $\langle v, \alpha_i \rangle \geq 0$ for $i = 1, 2$, which in polar coordinates corresponds to angles $\theta \in [0, \pi/3]$. The interior bisector is at $\theta = \pi/6$, i.e., ω_1 lies on the chamber bisector.

This is the content of `angle_alpha1_omega1_eq_pi_div_six` and `omega1_in_weyl_chamber_interior` (both in `UgpLean.MassRelations.SU3FlavorCartan`).

C Pre-Registered Basis Definitions for SC-JJJ, SC-KKK, SC-LLL

SC-JJJ: GUT-representation basis (96 atoms)

Basis SHA-256 (committed before scan):

644a7f90bc9682db672a8d8920df80a2c481e232c10f44c316a1321b626e8899.

Atom classes: classical Lie-group ranks $\{\text{rank}(\text{SU}(n))\}_{n=2}^8$, $\{\text{rank}(\text{SO}(n))\}_{n=4}^{10}$, $\{\text{rank}(E_n)\}_{n=6}^8$, $\text{rank}(F_4)$, $\text{rank}(G_2)$; representation dimensions of $\text{SU}(3)$, $\text{SU}(5)$, $\text{SO}(10)$ including $\dim(45_{\text{SU}(5)}) = 45$, $\dim(126_{\text{SO}(10)}) = 126$; SM hypercharges $Y_Q = 1/6$, $Y_u = 2/3$, $Y_d = -1/3$, $Y_L = -1/2$, $Y_e = -1$, $Y_H = 1/2$; gauge-boson counts $\{1, 3, 8, 4, 12\}$; small rationals $\{1/n\}_{n=1}^{14}$. DL ≤ 3 : 603,840 expressions.

SC-KKK: FN integer-charge derivation (analytical)

Pre-commit SHA-256: `pre_commit_sha256` in `artifact comp_p01_KKK_vv_rg_derivation.json`.

Derived analytically: the FN-doubled framework predicts down-sector charges $q_{d,g}^{(1)} = -(7/6) \cdot 2^{g-1}$ and $q_{d,g}^{(2)} = -13/9$, both non-integer. Seven integer-charge candidates tested computationally; all fail.

SC-LLL: Discrete-flavor basis (97 atoms)

Basis SHA-256 (committed before scan):

dc210d1240f9d8f8d3438fdf0e2443016f2e698f3d4d857a516360fd9861f180.

Discrete-symmetry groups from [10]: A_4 , S_4 , T' , A_5 , $\Delta(27)$, $\Delta(54)$, $\Sigma(168)$, $\Sigma(216)$, $\Sigma(72)$, $\Sigma(36)$, dihedrals D_3 – D_8 ; subgroup indices and representation dimensions; cyclotomic cosines $\cos(2\pi k/n)$ for $n \in \{3, 4, 5, 6, 7, 8\}$. DL ≤ 3 : 610,421 expressions.

D SHA-256 Artifact Table

Table 5: SHA-256 hashes of all computational artifacts used in this paper. Artifacts are in `papers/01_SM/canonical_run/`.

Artifact	File	SHA-256 (first 16 hex chars)
SC-TT	comp_p01_TT_....json	d8227f9d651b3d95
SC-VV	comp_p01_VV_....json	10f1bb1e099d444d
SC-EEE	comp_p01_EEE_....json	120fd2aa2d2270f1
SC-JJJ	comp_p01_JJJ_....json	f82288b4e6816cfe
SC-KKK	comp_p01_KKK_....json	e9956111050db7b6
SC-LLL	comp_p01_LLL_....json	db997f06ff06160f
SC-BBB	comp_p01_BBB_....json	ef33c48fa938e3fa

E Six β -Free Inter-Generational Identities

Table 6: The six β -free inter-generational identities derived from the TT formula by taking differences $R_g - R_{g'}$. No free parameters.

Identity	Predicted	PDG	Residual (%)	Note
$\log(m_c/m_e) - \log(m_u/m_e) = (\pi/6)(2^2 - 2^1)$	5.890	5.868	+0.37	
$\log(m_t/m_\tau) - \log(m_c/m_\mu) = (\pi/6)(2^3 - 2^2)$	5.243	5.250	−0.13	
$\log(m_t/m_\tau) - \log(m_u/m_e) = (\pi/6)(2^3 - 2^1)$	11.133	11.118	+0.14	
$\log(m_c m_e / m_u m_\mu)$	$\pi/3$	1.044	−0.03	
$\log(m_t m_\mu / m_c m_\tau)$	$\pi/3$	1.046	−0.16	
$\log(m_t m_e / m_u m_\tau)$	π	3.134	−0.23	Gelfond-type

References

- [1] N. Spivack. The Koide relation as a cyclotomic-12 closed form: A Lean-certified prediction of m_τ from (m_e, m_μ) at 61 ppm. Zenodo, 2026. Zenodo DOI placeholder — to be updated before submission.

- [2] N. Spivack. A deterministic number-theoretic framework for the standard model parameter spectrum, 2026. Zenodo DOI placeholder — to be updated before submission. Source: <https://github.com/novaspivack/ugp-physics>.
- [3] Nova Spivack. `ugp-lean`: Lean 4 formalization of UGP and GTE. <https://github.com/novaspivack/ugp-lean>, 2026. 117-module Lean 4 library (module count updated as of 2026-05-09). `lake build` to verify (zero sorry, zero custom axioms, standard Mathlib axiom signature). Includes `MassRelations`, `ElegantKernel` unconditional closure, `BraidAtlas` composites and charge derivation, `PSC RCC` infinite-family layer, `TE2.2` scan certification. New in `GTE.GeneralTheorems`: `E8` cyclotomic universality theorems (all 8 Zamolodchikov `E8` mass ratios lie in $\mathbb{Q}(\zeta_{120})$, `e8_all_masses_divisibility`, `e8_cyclotomic_divisibility`). Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [4] Nova Spivack. `ugp-lean`: A machine-checked formalization of the universal generative principle in Lean 4. Zenodo, 2026. Formalization paper (latest version): <https://doi.org/10.5281/zenodo.19554688>. Concept DOI (always-latest): <https://doi.org/10.5281/zenodo.19433538>. 117 modules across twelve architectural layers including `GTE`, `Universality`, `ElegantKernel` (UCL unconditional closure), `MassRelations` (Koide closed form, S_3 -Newton flow, binary cascade, physical mass spectrum), `BraidAtlas` (`ChargeTheorem`, `CompositeTriples`, `ChiralitySquaring`, `ChargeDerivation`, `CoxeterConductor`, `CoxeterConductorTowerLaw`, `EW-Bosons`), `GaloisStructure`, `PSC RCC` infinite-family closure, and `TE2.2` scan certification. Zero sorry, one disclosed axiom (`mirror_winding_number_zero` for `GTE-P7` dark matter, contained to mirror sector only). Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [5] Nova Spivack. `ugp-physics`: Companion code and papers for the UGP physics series. <https://github.com/novaspivack/ugp-physics>, 2026. Includes `computational_concordance/` tables, basin JSON exports, and capstone $\text{\LaTeX}$.
- [6] Howard Georgi and Cecilia Jarlskog. A new lepton-quark mass relation in a unified theory. *Physics Letters B*, 86:297–300, 1979.
- [7] Harald Fritzsch. Calculating the Cabibbo angle. *Physics Letters B*, 70:436–440, 1977.
- [8] C. D. Froggatt and H. B. Nielsen. Hierarchy of quark masses, Cabibbo angles and CP violation. *Nuclear Physics B*, 147:277–298, 1979.
- [9] Guido Altarelli and Ferruccio Feruglio. Discrete flavor symmetries and models of neutrino mixing. *Reviews of Modern Physics*, 82:2701, 2010.
- [10] H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, and M. Tanimoto. Non-Abelian discrete symmetries in particle physics. *Progress of Theoretical Physics Supplement*, 183:1–163, 2010.
- [11] Yoshio Koide. Fermion-boson two-body model of quarks and leptons and cabibbo mixing. *Lettere al Nuovo Cimento*, 34:201, 1982. See also: Y. Koide, “A new relation among lepton masses,” *Mod. Phys. Lett. A* **5**, 2319 (1990); and Y. Koide, arXiv:hep-ph/0506247 (2005) for a review of the trace identity $Q = (m_e + m_\mu + m_\tau)/(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 2/3$.
- [12] Alain Connes and Matilde Marcolli. *Noncommutative Geometry, Quantum Fields and Motives*, volume 55 of *Colloquium Publications*. American Mathematical Society, 2008. Spectral-triple derivation of the Standard Model Lagrangian; arithmetic motives in QFT.

- [13] Yuri I. Manin and Matilde Marcolli. Quantum statistical mechanics of the absolute Galois group. *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*, 16:038, 2020. Galois group action on arithmetic structures; dessins d’enfant; Grothendieck-Teichmüller group.

Acknowledgments

AI assistants were used for adversarial review, coding assistance, and debugging during manuscript preparation and code development. All scientific claims, derivations, and numerical results were independently verified by deterministic scripts and Lean proofs; no AI-generated content appears as unverified scientific output.