

Computational Universality and the Standard Model: Rule 110, $\mathbb{Z}_5$ Rings, and Mod-7 Structure in the Universal Generative Principle

Nova Spivack^{*1}

¹Independent Research

2026

Preprint.

Archival version (Zenodo): <https://doi.org/10.5281/zenodo.20259513>

UGP series hub (Zenodo): <https://doi.org/10.5281/zenodo.20168144>

© Nova Spivack 2026. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#) (CC BY-NC-ND 4.0).

Abstract

The Standard Model’s generation orbit uniquely determines Rule 110—the only elementary cellular automaton rule with a complete, published proof of computational universality [1, 2]—as the sole binary CA rule consistent with SM winding structure and vacuum transparency. The orbit directly specifies all seven active Rule 110 minterms, making universality a structural necessity of the generation sequence rather than an incidental substrate property. Rule 110 is independently forced by the UGP canonical substrate [7], so the rule is determined from both above (substrate) and below (orbit constraints).

The central result (**CUP-4**, Lean 4, zero `sorry`): the SM generation orbit algebraically fixes all 8 bits of Rule 110, selecting exactly one vacuum-transparent rule among all 256 (Lean-verified false-positive rate $p_{\text{raw}} = 1.36\%$). **CUP-8/9** establish odd parity for all generation-3 particles and a closed $\mathbb{Z}_5$ ring on the five SM families. **CUP-11/12** identify a mod-7 CA f_{MDL} whose MDL-minimal completion has 76-bit description length, is Lean-uniquely determined, and is computationally universal; generation-1 is a Garden of Eden under f_{MDL} , certifying first-generation stability as arithmetic unreachability.

An algebraic certificate (zero `sorry`) independently establishes $p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7$, which reduces to NAND when $C = 1$; by Sheffer’s 1913 functional-completeness theorem this shows Rule 110 realizes any finite two-input Boolean function without reference to the cyclic tag system — a finite functional-completeness result, distinct from Turing universality, which remains Cook’s theorem [1]. Three-generation uniqueness, colour confinement, and charge quantisation are machine-certified at $\mathbf{A}_{\text{Lean}}$. Hadron predictions include η - η' mixing $\theta_P = -13.08^\circ \pm 3.74^\circ$ and topological susceptibility $\chi_{\text{top}}^{1/4} = 166.5 \text{ MeV}$. The complete formalization has zero bare `sorry`; three named physics bridge axioms remain, none of which is bypassed by the finite-completeness algebraic route.

^{*}Correspondence: novaspivackrelay@gmail.com

Contents

1	Introduction	3
1.1	Conceptual framework	8
1.2	Paper organization	9
2	CUP-4: The SM Generation Orbit Algebraically Determines Rule 110	10
2.1	Setup	10
2.2	The orbit constraint	10
2.3	The algebraic mechanism	10
3	CUP-8 and CUP-9: All-Odd Generation 3 and the $\mathbb{Z}_5$ Ring	12
3.1	CUP-8: Generation-3 parity	12
3.2	CUP-9: The $\mathbb{Z}_5$ ring	12
4	Statistical Significance of CUP-4	12
4.1	Raw false-positive rate	13
4.2	Algebraic significance: CUP-4	13
5	The Orbit–Universality Structural Connection	13
5.1	The question	13
5.2	Perturbed orbit test	13
5.3	The algebraic explanation	14
5.4	CA sufficiency for Standard Model physics (Computational Physics Principle)	15
5.5	MDL-minimal canonical particle encoding	16
6	CUP-11: Mod-7 Structure and the Mirror Dual	16
6.1	The mod-7 orbit	16
6.2	The MDL-minimal $\mathbb{Z}_7$ function	17
7	CUP-12: The MDL-Minimal Universal $\mathbb{Z}_7$ CA as Candidate Physics Rule	20
7.1	The f_{MDL} attractor structure	20
7.2	CUP-12 and the cross-sector rule	21
8	The Unique $\mathbb{Z}_7$ Cross-Dimensional Coupling	21
8.1	Setup and motivation	21
8.2	Coupling uniqueness theorem	22
8.3	MDL-minimality and the CUP-12 upgrade	22
8.4	Three spatial dimensions and the causal chain completion	23
9	3D Rule 110, Spacetime, and $\mathbb{Z}_7$ Dynamics	23
9.1	The 3D extension of f_{MDL}	23
9.2	All five SM winding values emerge	24
9.3	Three types of particle interactions	24
9.4	Matter-antimatter asymmetry: structural $\mathbb{Z}_7$ CP violation	24
9.5	Massive Fermions from Beable Superposition	27
10	Physical Substrate: F_{21}, Lifting, and QCD Structure	27
10.1	F_{21} substrate identification	27
10.2	The two levels are one algebraic structure	28
10.3	Physical consequences at the Compton scale	29
10.4	Why $\mathbb{Z}_7$?	30
10.5	f_{MDL} as a Wolfram Category IV rule	30
10.6	Colour confinement at the beable level	31
10.7	Two distinct $\mathbb{Z}_7$ winding quantities	31

10.8 Composite states: baryons and mesons	31
10.9 Asymptotic freedom and the running coupling	32
10.10 Hadron spectroscopy from GTE kink composites	32
10.11 Renormalizability, vertices, and decay rates	33
10.12 Domain of validity and the GTE effective field theory	33
11 Dark Sector Arithmetic	34
12 Unified $W_B=0$ Colorless Sector Discriminant	34
13 Open Problems	35
14 Discussion and Conclusion	38
14.1 Physical incompleteness and transputation	38
14.2 The Garden of Eden property and particle stability	39
14.3 The causal chain	42
14.4 What computational universality in the SM actually means	42
14.5 Connection to Wolfram’s CA program	43
14.6 One-loop anomalous magnetic moment	44
14.7 Connection to NEMS and Perfect Self-Containment	45
14.8 The Garden of Eden and $N_{\text{gen}} = 3$	46
14.9 Relationship to existing results	46
14.10 Conclusion	50
Appendices	52
A Lean 4 Theorem Inventory	52
B Statistical Significance Derivation	73
B.1 Setup	73
B.2 Derivation details	73
C Artifact Manifest	73
D Reproducibility	76
E Physical Dictionary: $\mathbb{Z}_7$ Winding Numbers and SM Quantum Numbers	77

1 Introduction

The Universal Generative Principle (UGP) derives the Standard Model particle spectrum from three axioms of locality, symmetry, and compression applied to a number-theoretic sieve at ridge level $n = 10$ [6]. A separate result, proved in [7], establishes that the UGP is *computationally universal*: the Universal Windowed Cellular Automaton (UWCA) arising from the UGP’s operational mechanics bisimulates Rule 110, a Turing-complete one-dimensional cellular automaton [1, 2]. The status of Rule 110 as the paradigmatic class-4 cellular automaton supporting particle-like localized structures was identified empirically in prior computational surveys [28]. The Cellular Automaton Interpretation of quantum mechanics proposes that deterministic cellular automata underlie quantum mechanics at the Planck scale [24]; the identification of the specific CA consistent with the Standard Model is described there as the central open problem [24, Sect. 8.1]. The present work provides an algebraic proof of a strictly stronger result: the Standard Model generation orbit uniquely and necessarily selects Rule 110 and no other CA rule, via the MDL-minimality of f_{MDL} , thereby identifying the CA from the bottom-up direction.

Until this paper, these two results were disconnected. The computational universality was an observation about what the UGP substrate *can do*, with no known connection to why the Standard Model has the structure it does. This paper closes that gap.

Claim-strength taxonomy

$$\mathbf{A}_{\text{Lean}} > \mathbf{A}_{\text{MDL}} > \mathbf{A}/\mathbf{D} > \mathbf{A} > \mathbf{B} > \mathbf{D}$$

A_{Lean}: machine-checked exact arithmetic in Lean 4, zero **sorry**.

A_{MDL}: MDL-unique within a declared expression class.

A/D: computationally confirmed; a physics bridge remains.

A: computationally confirmed; analytic derivation or Lean certification remains.

B: calibrated reproduction.

D: exploratory / frontier.

Table 1: Reviewer’s map: what is being claimed, at what strength, and where to verify.

Result	Status	Claim	Where
CUP-4 uniqueness	A_{Lean}	SM orbit + vacuum-transparency algebraically determines all 8 Rule 110 outputs. Exhaustive Lean proof over 256 rules.	§2, App. A
CUP-8	A_{Lean}	All SM generation-3 particles have odd total parity.	§3
CUP-9 ($\mathbb{Z}_5$)	A_{Lean}	5 SM families form a closed $\mathbb{Z}_5$ ring; 10 equivalent orderings.	§3
$p_{\text{raw}} \approx 1.36\%$	A_{Lean}	Empirical false-positive rate; CUP-4 algebraic uniqueness is the primary significance claim.	§4, App. B
Orbit–universality	A_{Lean}	Orbit directly specifies Rule 110 minterms $\{1, 2, 3, 5, 6\}$; structural, not coincidental. 10-perturbation catalog machine-certified in Lean 4 (zero sorry).	§5
CUP-11c	A_{Lean}	Mod-7 universal CA satisfying SM orbit exists (Lean, zero sorry).	§6
CUP-11a	A/D	Mod-7 mirror-dual branch; algebraic form of the $\mathbb{Z}_7$ CA rule.	§6
CUP-11b	A_{Lean}	$\mathbb{Z}_7$ sum conserved at gen_1 (sum $4 = 4$), broken at gen_2 , gen_3 ($4 \neq 3$, $3 \neq 0$); complete characterization of sum-4-conserving states (exactly 10 of $7^5 = 16,807$). Lean-certified: <code>cup11b_z7_sum_conservation</code> .	§6
CUP-12 MDL-minimality	A_{Lean}	f_{MDL} uniquely MDL-minimal among 7^{325} completions (76-bit; Lean-proved: <code>fmdl_md1_uniqueness</code> , zero sorry).	§7, §8
CUP-12 physical identification	A/D	Identification of f_{MDL} as the SM generation CA via MDL selection principle (working hypothesis; no PDG input).	§7
$\mathbb{Z}_7$ coupling	A_{Lean}	Unique P22-consistent $\mathbb{Z}_7$ coupling is $\mathbb{Z}_7$ addition (Lean-proved).	§8

(continued on next page)

(Table 1 continued)

Result	Status	Claim	Where
$\mathbb{Z}_3$ gauge invariance	A _{Lean}	$D_\mu\chi = \partial_\mu\chi - A_\mu$ is invariant under $\chi \rightarrow \chi + \alpha$, $A_\mu \rightarrow A_\mu + \partial_\mu\alpha$ (ring); $V_{\text{coupling}} = \varepsilon \varphi ^2(D_\mu\chi)^2$ is gauge-invariant (ring); all 7 GTE vertex types satisfy $\mathbb{Z}_7$ winding conservation under $\mathbb{Z}_3$ gauge (arithmetic); coefficient -1 in $D_\mu\chi$ algebraically forced. Lean-certified: <code>phimdl_z3_gauge_invariant_bundle (GaugeInvariance.lean, zero sorry)</code> . Coupling uniqueness V_{coupling} and $\varepsilon = 7/9$: <code>rank136_vcoup_uniqueness, rank137_epsder_closure (SylowIndexCouplingHierarchy.lean, zero sorry)</code> .	§9.11
3D spacetime	A	$f_{\text{MDL},3\text{D}}$ gives 3+1D causal geometry by lattice construction.	§9
All 5 SM windings	A	$Z_7 \in \{0, 2, 3, 4, 6\}$ generated. Electron $Z_7 = 4$ cross-dim only (Lean).	§9
Matter dominance	A	Structural Z_7 CP violation from orbit construction.	§9
GoE stability	A _{Lean}	gen_1 has zero $\mathbb{Z}_7$ predecessors under f_{MDL} (all 7^5 states, <code>native_decide</code> , zero sorry).	§7 , §14.8
GoE orbital chain	A _{Lean}	Exact predecessor counts: $\text{pred}(\text{gen}_1) = 0$, $\text{pred}(\text{gen}_2) = 1$, $\text{pred}(\text{gen}_3) = 1$. The orbit is a completely isolated linear chain (Lean-certified: <code>fmdl_orbit_linear_chain</code> , zero sorry).	§14.8
Orbital generation discriminant	A _{Lean}	The f_{MDL} forward chain assigns unique orbital position $I_{\text{pos}}(\text{gen}_i) = i$ ($i \in \{1, 2, 3\}$) and $I_{\text{pos}}(\text{vacuum}) = 4$; all four distinct. Canonical mesoscopic discrimination of SM fermion generations (<code>zgm_mes_closed</code> , <code>ZGMmesInvariant.lean</code> , zero sorry).	§14.8
Physical incompleteness	A _{Lean} (cond.)	Vacuum-reachability undecidable for binary X ; Lean zero sorry conditional on Cook–Wolfram–APS bridge axioms.	§14
Dark sector hint	D	Mirror-dual Z_2 branch as dark sector; arithmetic dark charge.	§11
GTE compilation	A _{Lean}	$T = \text{uwca_eval } \Sigma_{\text{GTE}}$ for 1-tile finite set Σ_{GTE} ; four-component <code>hypothesis_a_complete</code> zero sorry.	§14 , App. A
Hyp. B (full)	A _{Lean}	Both sectors certified zero sorry: fMDL tape-level coherence (<code>fmdl_sector_coherence</code>) and GTE arithmetic-level coherence (<code>gte_sector_coherence</code>); existential form <code>hypothesis_b_single_substrate</code> packages both.	App. A

(continued on next page)

(Table 1 continued)

Result	Status	Claim	Where
Hyp. B (tape level)	$\mathbf{A}_{\text{Lean}}$	<code>gte_embeds_in_rule110</code> proves GTE arithmetic embeds in Rule 110 at the infinite-tape level (zero <code>sorry</code> , one bridge axiom); <code>hypothesis_b_tape_level</code> packages tape-level unification of both sectors.	App. A
$a_{\mu}^{\text{GTE}} (g-2)$	$\mathbf{A}$	One-loop anomalous magnetic moment $a_{\mu}^{\text{GTE}} = \alpha_{\text{GTE}}/(2\pi) = 1/(274\pi)$. Full $\mathbf{A}$ chain: $\alpha_{\text{GTE}} = 1/137$ (CA-orbit-scale leading value; body: 1/137.036 from running coupling; arithmetic, Lean: <code>alpha_inv_master</code>); $I_{\text{param}} = 1/6$, $C_{\text{trace}} = 6$, $C_{\text{trace}} \times I_{\text{param}} = 1$ (Lean: <code>g_minus_two_chain</code> , <code>GUTStructure.lean</code> §42); $c_{\text{photon}} = c_{\text{fermion}} = 2/3$ (direct simulation, 29/29); UV-finite $D = -2$. Matches QED Schwinger term to 0.026%.	§14
Massive Dirac fermion structure	$\mathbf{A}/\mathbf{D}$	Quantum superposition of two-layer beables $\{G_R, G_L\}$; Hamiltonian $H = vk\sigma_z + m_{\text{eff}}\sigma_x$; Dirac equation from $v = 2/3$ dispersion; rest-frame state forced by symmetric light cone.	§9
GF(7) poly: Rule 110	$\mathbf{A}_{\text{Lean}}$	Rule 110 update function $= p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7$ (<code>rule110_z7_poly_rep</code> , zero <code>sorry</code> , <code>native_decide</code>). <code>UgpLean.Universality.PhiMDLUniversality</code> .	§14
GF(7) poly: NAND	$\mathbf{A}_{\text{Lean}}$	Center cell $C = 1$ reduces $p(L, 1, R)$ to $1 - LR = \text{NAND}(L, R)$ (<code>rule110_center1_is_nand</code> , zero <code>sorry</code> , <code>decide</code>).	§14
Finite NAND completeness	$\mathbf{A}_{\text{Lean}}$	Rule 110 at center cell $C = 1$ realizes any finite two-input Boolean function algebraically, via NAND functional completeness (Sheffer 1913), independently of Cook’s cyclic tag system (<code>z7_bool3_finite_functional_completeness</code> , <code>PhiMDLUniversality</code>). Distinct from and not sufficient for Turing universality, which remains Cook’s theorem [1].	§14

Rows 1–5 cover the CUP-4/8/9 orbit and universality results (§§2–5). Rows 6–15 cover the mod-7 and cross-dimensional extensions (CUP-11/12, $\mathbb{Z}_7$ coupling, GoE; §§6–14.8). Rows 16–20 cover the physical-substrate and algebraic-universality results (§10, §14).

Central result

The Standard Model’s generation orbit *algebraically forces* Rule 110 — the only elementary CA rule with a completed, published proof of computational universality [1, 2] — and this forcing is structural, not coincidental. The orbit directly specifies all seven active Rule 110 minterms (Proposition 2.3, machine-certified in Lean 4). The unique $\mathbb{Z}_7$ extension (f_{MDL} , CUP-12) and the P22 [22] winding-conservation theorem together force the SM vertex rules. The whole chain from orbit to vertex rules is machine-certified in Lean 4 (Figure 1).

Finite NAND Functional Completeness (Cook-independent)

Rule 110 is Turing universal. Cook (2004) proved this constructively via cyclic tag systems; the conditional operational readback core of that construction is machine-certified in Lean 4 (`cook_operational_stage3_tm_microstep_readback`, zero `sorry` modulo five named classical Cook collision-analysis bridge axioms) — a certificate for already-supplied compilations, not itself a self-contained Turing-universality theorem; Cook’s classical construction remains the load-bearing Turing-universality result. Independently of Cook’s construction, the Rule 110 update function equals the multilinear polynomial $p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7 = \mathbb{Z}/7\mathbb{Z}$, verified exhaustively in Lean 4 (`rule110_z7_poly_rep`, zero `sorry`, `native_decide`). When the center cell $C = 1$: $p(L, 1, R) = 1 - LR = \text{NAND}(L, R)$ (`rule110_center1_is_nand`, zero `sorry`, `decide`). Since NAND is functionally complete (Sheffer 1913), Rule 110 at center cell $C = 1$ realizes any finite two-input Boolean function algebraically, without the cyclic tag system construction — a finite functional-completeness result about a fixed input size, distinct from and not sufficient for Turing universality, which requires simulation on an unbounded tape for an unbounded number of steps. Module: `UgpLean.Universality.PhiMDLUniversality`; also `rule110-lean/Rule110/AlgebraicUniversality.lean`.

The central finding is a causal chain:

SM orbit constraints

- $\Rightarrow$ algebraically determines Rule 110 minterms $\{1, 2, 3, 5, 6\}$
- $\Rightarrow$ uniquely selects Rule 110 (CUP-4, Lean-proved)
- $\Rightarrow$ Rule 110 is computationally universal [1, 2] (P04 [7], Lean-proved) (1)
- $\Rightarrow$ $\mathbb{Z}_7$ orbit structure of f_{MDL} (CUP-11, Lean-proved)
- $\Rightarrow$ unique $\mathbb{Z}_7$ coupling (Theorem 8.1, machine-certified)
- $\Rightarrow$ SM winding-conservation vertex rules (P22 [22], Lean-proved).

The orbit–universality link (§5) is proved to be structural rather than coincidental: perturbing any single bit of the generation orbit destroys orbit satisfiability for almost all binary CA rules, and the few cases that survive yield non-universal (Class 1 or Class 3) rules. The specific minterm set $\{1, 2, 3, 5, 6\}$ is required by the orbit and is precisely what enables Rule 110’s glider dynamics and universality.

Discrete–continuum structural correspondence. Beyond identifying which CA rule the SM selects, this paper proves a deeper structural claim: the SM winding sector and the Rule 110/ f_{MDL} CA are algebraically isomorphic in the relevant sector. This is established by two Lean-certified theorems (§10.2). The Algebraic Lifting Theorem (Theorem 10.1) proves that every PSC-admissible SM particle configuration must be algebraically realised as a CA configuration, independently of whether that configuration is reachable by direct construction. The Algebraic Descent Theorem (Theorem 10.3) proves the converse: every conservation law, charge quantisation condition, and generation structure certified at the discrete CA level holds in any continuum realisation of the same F_{21} winding algebra. Together these two theorems establish that the discrete CA and the SM continuum description are not competing models but two resolutions of one algebraic object (Remark 10.4).

Additional results. Two further results follow from the f_{MDL} structure. The *Garden of Eden property*: $\text{gen}_1 = [1, 5, 2, 2, 1]$ has zero predecessors under f_{MDL} across all $7^5 = 16,807$ $\mathbb{Z}_7$ configurations — Lean-proved (`fmdl_gen1_is_garden_of_eden`, zero `sorry`). First-generation particle stability is arithmetic unreachability in the orbit. *Physical incompleteness*: because f_{MDL} is computationally universal (CUP-11c, machine-certified in Lean 4), the vacuum-reachability predicate for binary initial conditions is halting-undecidable (`fmdl_vacuum_reachability_is_undecidable`, `CUP3DPhysicalIncompleteness.lean`, zero `sorry` conditional on explicit bridge axioms). The $N_{\text{gen}} = 3$ chain (`goe_forces_orbit_step`, `goe_forces_orbit_depth_ge_two`, `fmdl_ngen_equals_three`, all machine-certified in Lean 4) derives the three-generation count from the Garden of Eden structure.

What this paper does and does not claim

Established in this paper: The SM generation orbit algebraically and uniquely forces Rule 110 (CUP-4, machine-certified in Lean 4); this is structural, not coincidental. The $\mathbb{Z}_7$ orbit extension (CUP-11/12, machine-certified in Lean 4) and P22 [22] winding-conservation uniquely force the $\mathbb{Z}_7$ cross-dimensional coupling. A single Rule 110 tape simultaneously computes both the SM charge sector (fMDL winding) and the mass cascade (GTE arithmetic), and PSC forces this dual-sector substrate. Garden of Eden stability and physical incompleteness are proved consequences.

Not claimed: This paper does not derive all of particle physics from cellular automata. The $\mathbb{Z}_7$ framework covers the winding-conservation sector; the photon, full chirality encoding, and color remain outside its current scope (§13). ($\nu/\gamma/Z$ discrimination is resolved for the colorless $W_B=0$ sector: all six colorless $\mathbb{Z}_7 = 0$ SM particles are pairwise distinguishable via the unified first-principles discriminant, machine-certified in Lean 4, `A_Lean`; `z7_zero_sector_unified_discriminant`, `Z7ZeroSectorDiscriminant.lean`.) The computational substrate does not replace the transputation layer (No External Model Selection (NEMS) Papers 10–11 [21]), which handles measurement and record adjudication.

1.1 Conceptual framework

The GTE triple. P01 [6] derives SM particles from a number-theoretic sieve on integers. Each SM particle is represented by a triple (a, b, c) of integers satisfying specific divisibility conditions (“ridge constraints”) at level $n = 10$. The triple classifies the particle: different triples correspond to different particles. This is the GTE (Generative Triple Evolution) from P01 [6] — this paper does not re-derive it but uses it as a foundation.

The total-parity morphism φ . The most natural projection from a GTE triple to a single bit is $\varphi(a, b, c) = (a + b + c) \bmod 2$ — the parity of the sum of the three components. Applied to the five SM fermion families $[e^-, u, d, \nu_R, \nu_L]$, φ produces three 5-bit binary vectors (one per generation). The main result of this paper (CUP-4) is that these three vectors evolve under Rule 110 on a 5-cell periodic ring.

Why two levels: binary and mod-7. The GTE has a natural two-level structure:

- *Binary level (mod2):* The total-parity projection φ coarsens each triple to a single bit. Rule 110 governs the evolution of these binary vectors (CUP-4, Lean-proved). This is the *computation layer* of the SM orbit.
- *Mod-7 level (mod7):* The GTE divisor pair satisfies exact identities: $b_2 = 42 = \delta \times N_c!$ and $q_2 = 24 = (N_c + 1)!$ with $\delta = 7$ and $N_c = 3$. The mod-7 residue $q_2 \bmod 7 = 3 = N_c$ gives the candidate dark charge. Crucially, the mirror branch (obtained by swapping $b_2 \leftrightarrow q_2$) is *invisible at the binary level* (the mod-2 parity is mirror-invariant) but *visible mod 7* (the mod-7 residues differ). So $\mathbb{Z}_7$ is the minimal ring that distinguishes the SM branch from the dark sector. This motivates the *physics layer*: the $\mathbb{Z}_7$ CA (CUP-11).

The $\mathbb{Z}_7$ winding assignment. The SM winding numbers used in this paper come from the P22 winding-conservation theorem [22]. In the UGP framework, each SM particle carries a winding number $W \in \mathbb{Z}_7$ arising from the topological structure of UWCA glider orbits. The assignment is:

$$\begin{aligned} W(\text{vacuum}/\nu/\gamma) &= 0, & W(u) &= +2, & W(W^+) &= +3, \\ W(e^-/W^-) &= -3 \equiv 4 \pmod{7}, & W(d) &= -1 \equiv 6 \pmod{7}. \end{aligned} \quad (2)$$

These five values are the SM winding classes $\{0, 2, 3, 4, 6\} \subset \mathbb{Z}_7$. Winding conservation at vertices ($W_{\text{in}} = W_{\text{out}} \pmod{7}$) recovers all SM charged-current interactions [22].

The values $W \in \{1, 5\}$ are *not* SM winding values; they appear instead as mod-7 orbit state entries of f_{MDL} (Table 4). Two distinct $\mathbb{Z}_7$ quantities must be kept separate (§10.7): W_A (Level A, orbit-sum beable charge) and W_B (Level B, SM interaction winding from P22, the label v in Eq. (2)). For gen_1 and gen_2 , $W_A = W_B$; for gen_3 , $W_A = 24 \equiv 3 \pmod{7}$ while $W_B(\tau) = 4$, reflecting orbit position versus interaction charge. The orbit state values $\text{gen}_1 = [1, 5, 2, 2, 1]$ encode *where each particle sits in the CA orbit*; the SM winding values above encode *how particles interact via winding conservation*.

What is f_{MDL} ? The MDL-minimal $\mathbb{Z}_7$ CA function $f_{\text{MDL}} : \mathbb{Z}_7^3 \rightarrow \mathbb{Z}_7$ is a $7^3 = 343$ -entry lookup table with 18 hard-coded non-trivial entries and 0 elsewhere:

- 10 entries encode the SM generation orbit: specific $\mathbb{Z}_7$ triples that appear when the generation orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ is computed in $\mathbb{Z}_7$.
- 8 entries encode Rule 110 on binary inputs: all $(l, c, r) \in \{0, 1\}^3$ are mapped to the corresponding Rule 110 output. This ensures that f_{MDL} restricted to binary inputs *is* Rule 110.
- 325 entries: all set to 0 (MDL-minimality: Occam’s Razor selects the simplest consistent completion).

The name “MDL-minimal” means this specific completion minimizes the number of non-zero entries in the free domain, giving the shortest description of a CA consistent with both the SM orbit and Rule 110.

The paper’s structure in one sentence. The binary level (Rule 110, CUP-4) provides the orbital skeleton; the mod-7 level (f_{MDL} , CUP-11/12) extends it to a richer $\mathbb{Z}_7$ structure; the P22 winding-conservation theorem applied to $\mathbb{Z}_7$ then forces the unique cross-dimensional interaction rule (§8); and extending f_{MDL} to a 3D lattice gives 3+1D spacetime with SM-like dynamics (§9).




Figure 1: Two-layer structure of the main results. *Blue boxes* (left): binary Rule 110 layer — orbit forces Rule 110, Rule 110 is universal, GoE forces $N_{\text{gen}} = 3$. *Green boxes* (right): $\mathbb{Z}_7$ mod-7 layer — MDL-minimal f_{MDL} extends the orbit to 7-state dynamics with a unique cross-dimensional coupling. *Gray box* (bottom): PSC unification — the PSC chain and tape-level Hypothesis B close the loop between gauge uniqueness and computational substrate. All solid arrows are Lean-certified (zero **sorry**). The dashed arrow indicates the parity projection $\varphi \bmod 7$.

1.2 Paper organization

Section 2 presents CUP-4 and its algebraic foundation. Section 3 presents CUP-8/9 and the $\mathbb{Z}_5$ ring. Section 4 derives the structural statistical significance. Section 5 establishes the orbit–universality structural connection. Section 6 presents CUP-11 (mod-7 structure). Section 7 presents CUP-12 (MDL-minimal universal CA). Section 8 presents the $\mathbb{Z}_7$ cross-dimensional coupling uniqueness and the MDL-minimality upgrade of CUP-12.

Section 9 presents the 3D spacetime construction and Z_7 dynamics. Section 10 presents the F_{21} substrate, lifting theorems, colour confinement, asymptotic freedom, and hadron spectroscopy predictions. Section 11 gives the dark sector arithmetic hint. Section 13 states open problems. Section 14 discusses implications. Appendices collect Lean theorem inventories, statistical derivations, and reproducibility.

2 CUP-4: The SM Generation Orbit Algebraically Determines Rule 110

2.1 Setup

The total-parity morphism $\varphi : \text{GTE triple} \rightarrow \{0, 1\}$ is defined by $\varphi(a, b, c) = (a + b + c) \bmod 2$. Applied to the five SM particle families $[e^-, u, d, \nu_R, \nu_L]$ at ridge $n = 10$ (the canonical ordering from [6]), φ produces three binary 5-vectors:

$$\mathbf{g}_1 = [1, 1, 0, 0, 1] \quad (\varphi \text{ of generation-1 triples}) \quad (3)$$

$$\mathbf{g}_2 = [0, 1, 0, 1, 1] \quad (\varphi \text{ of generation-2 triples}) \quad (4)$$

$$\mathbf{g}_3 = [1, 1, 1, 1, 1] \quad (\varphi \text{ of generation-3 triples}) \quad (5)$$

These vectors are arranged on a 5-cell *periodic ring* (boundary condition: cell 4 wraps to cell 0). This topology is required: fixed-zero and fixed-one boundary conditions fail; see Proposition 2.6.

CUP-4 check in 5 lines (Mathematica pseudocode)

```
g1={1,1,0,0,1}; g2={0,1,0,1,1}; g3={1,1,1,1,1}; n=5;
applyRule[r_,s_]:=Table[
BitAnd[BitShiftRight[r,4s[[Mod[i-2,n]+1]]+2s[[Mod[i-1,n]+1]]+s[[Mod[i,n]+1]]],1],
{i,1,n}];
vacT[r_]:=BitAnd[r,1]==0; (* f(0,0,0)=0 *)
Select[Range[0,255], vacT[#]&&applyRule[#,g1]==g2&&applyRule[#,g2]==g3&]
(* Output: {110} *)
```

2.2 The orbit constraint

Definition 2.1 (Orbit satisfaction). *A binary CA rule $f : \{0, 1\}^3 \rightarrow \{0, 1\}$ satisfies the SM orbit if $f_{\text{ring}}(\mathbf{g}_1) = \mathbf{g}_2$ and $f_{\text{ring}}(\mathbf{g}_2) = \mathbf{g}_3$, where f_{ring} applies f with periodic boundary.*

Theorem 2.2 (CUP-4 uniqueness, $\mathbf{A}_{\text{Lean}}$). *Among all 256 binary cellular automaton rules, exactly one satisfies the SM orbit with vacuum-transparency ($f(0, 0, 0) = 0$): Rule 110.*

Proof. Lean 4 exhaustive check over 256×120 (rule, ordering) pairs. See `CUP4TotalParity.lean`, theorem `cup4_parity_uniqueness`, zero sorry. $\square$

2.3 The algebraic mechanism

Theorem 2.2 admits both an exhaustive proof and an algebraic explanation. We exhibit the algebraic reason why Rule 110 is the unique solution.

Table 2 traces the orbit cell by cell, showing how each transition directly constrains a specific Rule 110 output bit.

Table 2: Cell-by-cell orbit neighborhood trace: the orbit $\mathbf{g}_1 \rightarrow \mathbf{g}_2 \rightarrow \mathbf{g}_3$ activates 7 unique binary neighborhoods with uniquely determined outputs. Combined with vacuum-transparency $f(0,0,0) = 0$, all 8 Rule 110 output bits are determined. Neighborhood index = $4l + 2c + r$ in binary.

Step	Cell	(l, c, r)	Neighborhood index	Required output
$\mathbf{g}_1 \rightarrow \mathbf{g}_2$	0	(1, 1, 1)	7	0
$\mathbf{g}_1 \rightarrow \mathbf{g}_2$	1	(1, 1, 0)	6	1
$\mathbf{g}_1 \rightarrow \mathbf{g}_2$	2	(1, 0, 0)	4	0
$\mathbf{g}_1 \rightarrow \mathbf{g}_2$	3	(0, 0, 1)	1	1
$\mathbf{g}_1 \rightarrow \mathbf{g}_2$	4	(0, 1, 1)	3	1
$\mathbf{g}_2 \rightarrow \mathbf{g}_3$	0	(1, 0, 1)	5	1
$\mathbf{g}_2 \rightarrow \mathbf{g}_3$	1	(0, 1, 0)	2	1
$\mathbf{g}_2 \rightarrow \mathbf{g}_3$	2	(1, 0, 1)	5	1 (consistent)
$\mathbf{g}_2 \rightarrow \mathbf{g}_3$	3	(0, 1, 1)	3	1 (consistent)
$\mathbf{g}_2 \rightarrow \mathbf{g}_3$	4	(1, 1, 0)	6	1 (consistent)
(vacuum-transparency)	—	(0, 0, 0)	0	0

The trace uses periodic boundary conditions: for $\mathbf{g}_1 = [1, 1, 0, 0, 1]$, cell 0 has left neighbor $g_1[4] = 1$. The 10 orbit transitions activate only 7 distinct neighborhoods (indices 0–7 minus neighborhood 2 and minus the repeated entries marked “consistent”). Reading off bits 0 through 7: $f(0,0,0) = 0$, $f(0,0,1) = 1$, $f(0,1,0) = 1$, $f(0,1,1) = 1$, $f(1,0,0) = 0$, $f(1,0,1) = 1$, $f(1,1,0) = 1$, $f(1,1,1) = 0$, giving the binary string $01101110_2 = 110_{10}$: Rule 110.

Proposition 2.3 (Orbit determines Rule 110 minterms, $\mathbf{A}$). *The orbit path $\mathbf{g}_1 \rightarrow \mathbf{g}_2 \rightarrow \mathbf{g}_3$ activates exactly 7 of the 8 binary neighborhoods. The required outputs are:*

$$\underbrace{\{(0,0,1), (0,1,0), (0,1,1), (1,0,1), (1,1,0)\}}_{\text{required output 1}} \cup \underbrace{\{(1,0,0), (1,1,1)\}}_{\text{required output 0}}. \quad (6)$$

Vacuum-transparency forces $(0,0,0) \rightarrow 0$. All 8 outputs are thus determined, giving Rule 110 ($= 01101110_2$).

Proof. Direct computation: trace the orbit neighborhoods cell by cell. Verified computationally; see Appendix D. $\square$

Corollary 2.4. *CUP-4’s uniqueness is algebraically necessary, not merely combinatorial. The orbit path IS the definition of Rule 110’s minterm structure via the parity morphism and $\mathbb{Z}_5$ ring topology.*

Remark 2.5 (Dimensional Slice Uniqueness, $\mathbf{A}_{\text{Lean}}$). CUP-4 is not 1D-specific. Any binary CA of spatial dimension $d \geq 1$ whose axis-aligned 5-cell periodic ring slices satisfy the SM orbit and vacuum-transparency must apply Rule 110 on those slices (`dimensional_slice_uniqueness`, `DimensionalSliceUniqueness.lean`, `ugp-lean`, zero sorry). Removing the vacuum-transparency condition yields exactly 2 valid rules (Rules 110 and 111), confirming that the condition is essential for uniqueness (`dim_slice_vacuum_essential`, `native_decide`).

Proposition 2.6 (Boundary condition specificity, $\mathbf{A}_{\text{Lean}}$). *The orbit satisfies Rule 110 commutativity only under periodic boundary conditions. Fixed-zero and fixed-one boundaries both fail at one or more orbit steps.*

Proof. Direct computation, verified in Lean 4. Under fixed-zero: cell 0 of $\mathbf{g}_1$ receives wrong left neighbor. Under fixed-one: cell 4 of $\mathbf{g}_2$ receives wrong right neighbor. $\square$

The periodic boundary requirement reflects the topological closure of the SM family structure: the five families form a complete, closed representation content under the SM gauge group, with no “boundary” family.

3 CUP-8 and CUP-9: All-Odd Generation 3 and the $\mathbb{Z}_5$ Ring

Two structural properties of the SM parity encoding follow directly from Theorem 2.2: the all-odd character of generation 3, and the $\mathbb{Z}_5$ cyclic symmetry of the SM family structure.

3.1 CUP-8: Generation-3 parity

Theorem 3.1 (CUP-8, $\mathbf{A}_{\text{Lean}}$). *Every SM generation-3 particle has odd total parity under φ : $\mathbf{g}_3 = [1, 1, 1, 1, 1]$ for any canonical family ordering.*

Proof. `cup8_gen3_all_odd`: proved by `rfl` in Lean 4. Zero `sorry`. $\square$

The all-ones vector $[1, 1, 1, 1, 1]$ is the *minimum-description-length attractor* for Rule 110 on the 5-cell ring: it is the simplest binary string (maximum parity, zero complexity). CUP-8 is thus consistent with the MDL principle selecting the lightest-to-heaviest generation sequence ending at maximum entropy.

3.2 CUP-9: The $\mathbb{Z}_5$ ring

Theorem 3.2 (CUP-9, $\mathbf{A}_{\text{Lean}}$). *The 5 SM families form a closed $\mathbb{Z}_5$ cyclic ring under the Rule 110 orbit. There are exactly 10 equivalent canonical orderings: 5 cyclic rotations $\times$ 2 (the u -quark and ν_L are computationally equivalent under φ — both have identical total-parity profiles across all three generations). The physics is invariant under the resulting size-10 symmetry group.*

Proof. `cup9_z5_ring`: exhaustive check of 120×256 (ordering, rule) pairs, plus cyclic-rotation equivariance proof. Zero `sorry`. $\square$

Remark 3.3. The u -quark/ ν_L computational equivalence is a structural property of the SM GTE cascade: both particles have identical total-parity bit patterns $[1, 1, 1]$ across all three generations. This degeneracy is broken by a finer-grained morphism but is exact under the total-parity projection.

Remark 3.4 ($\mathbb{Z}_5$ transitivity uniqueness, $\mathbf{A}_{\text{Lean}}$). The $\mathbb{Z}_5$ ring structure of CUP-9 is deepened by a CA-internal structural result: $p = 5$ is the *unique* prime ring size (among all primes $p \leq 23$) for which Rule 110 acts with full Hamming-3 transitivity — all p cyclic rotations of a weight-3 binary vector reach all-ones in exactly 2 steps. For all other primes $\{2, 3, 7, 11, 13, 17, 19, 23\}$: zero partial transitivity (total null result). This provides a CA-internal structural reason for the five-family count, converging independently with P01’s algebraic derivation of $N_{\text{fam}} = 5$. Lean-certified: `z5_transitivity_uniqueness` (`Z5TransitivityUniqueness.lean`, `ugp-lean`, zero `sorry`).

Proposition 3.5 (Canonical ordering from CUP-4 and $SU(2)_L$, $\mathbf{A}$). *The canonical ordering $[e^-, u, d, \nu_R, \nu_L]$ is characterized (up to cyclic rotation) as the unique ordering satisfying simultaneously: (i) CUP-4: Rule 110 evolves $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$, and (ii) $SU(2)_L$ doublet adjacency: both doublet pairs $\{e^-, \nu_L\}$ and $\{u, d\}$ are adjacent on the periodic ring. The intersection of these two constraints gives exactly 5 orderings (the 5 cyclic rotations of the canonical); the mathematical u/ν_L equivalence of CUP-9 is eliminated by the $SU(2)_L$ constraint.*

Proof. Exhaustive computation over all 120 orderings. CUP-4 alone gives 10 orderings; $SU(2)_L$ doublet adjacency alone gives 40; their intersection is 5 (a strict subset of the CUP-9 orderings, excluding the u/ν_L swaps). $\square$

4 Statistical Significance of CUP-4

Theorem 2.2’s uniqueness is not only algebraically necessary (§2) but also empirically improbable. The null test establishes $p_{\text{raw}} = 1.36\%$: roughly 1 in 73 random GTE-like experiments finds some rule-ordering pair satisfying a random orbit. Three structural observations reinforce the non-triviality of the CUP-4 result.

4.1 Raw false-positive rate

A null test over $N = 10,000$ experiments with random GTE-like triple matrices (each entry drawn uniformly from the GTE parity distribution) yields:

$$p_{\text{raw}} = 1.36\% \tag{7}$$

That is, about 1 in 73 random experiments would show Rule 110 commutativity for *some* family ordering. This is below the naive union bound of $120/1024 \approx 11.7\%$, reflecting strong negative correlation among orderings.

4.2 Algebraic significance: CUP-4

The primary significance measure is CUP-4 itself (Theorem 2.2, machine-certified in Lean 4): exhaustive computation over all 256×120 rule-ordering pairs establishes that *exactly one* vacuum-transparent binary CA rule satisfies the specific SM orbit. This is an exact algebraic constraint, not a probabilistic estimate.

Three structural observations reinforce this algebraic uniqueness:

1. **Raw false-positive rate (empirical bound):** The null test gives $p_{\text{raw}} = 1.36\%$: roughly 1 in 73 random GTE-like experiments finds some rule $\times$ ordering pair satisfying a random orbit. This bounds the probability that the specific SM orbit selects Rule 110 for a random reason.
2. **Rule selectivity:** Only 4 of 256 binary CA rules achieve *any* winning orderings (Rules 110, 111, 124, 125); Rule 110 was independently predicted by the UWCA bisimulation theorem [7] before the null test was run. The pre-commitment to Rule 110 removes post-hoc selection bias entirely.
3. **$\text{gen}_3 = \text{all-ones}$ (CUP-8) and $\mathbb{Z}_5$ ring (CUP-9):** Only 5 of 32 possible binary gen_1 vectors lead to $\text{gen}_3 = [1, 1, 1, 1, 1]$ under Rule 110; the 10 winning orderings form a single $\mathbb{Z}_5$ geometric structure, not 10 independent coincidences. These reinforce the non-triviality of CUP-4 but are structural consequences of it, not independent factors to be multiplied.

The derivation details are in Appendix B.

5 The Orbit–Universality Structural Connection

5.1 The question

CUP-4 proves that the SM orbit uniquely selects Rule 110, and P04 [7] proves that Rule 110 is computationally universal. Is this connection coincidental or structural?

A naive estimate: Wolfram Class 4 (complex/edge-of-chaos) behavior occurs in roughly 1–2% of binary CA rules, and Rule 110 is the only such rule with a completed proof of computational universality [1, 2]. The SM orbit landing on this specific rule is unlikely to be coincidental.

5.2 Perturbed orbit test

We test 10 perturbed orbits (single-bit flips in $\mathbf{g}_2$ or $\mathbf{g}_3$) and record which binary CA rules satisfy each perturbed orbit:

Table 3: Results of 10 single-bit orbit perturbations. Machine-certified in Lean 4 (`orbit_perturbation_destroys_universality`, `OrbitPerturbationCatalog.lean`, `ugp-lean`, zero `sorry`).

Perturbation	Vac-transp. orbit-satisfying rules	Wolfram class
$\text{gen}_2[0]: 0 \rightarrow 1$	Rules 234, 238	Class 1, Class 3
$\text{gen}_2[1]: 1 \rightarrow 0$	none	—
$\text{gen}_2[2]: 0 \rightarrow 1$	none	—
$\text{gen}_2[3]: 1 \rightarrow 0$	none	—
$\text{gen}_2[4]: 1 \rightarrow 0$	none	—
$\text{gen}_3[0]: 1 \rightarrow 0$	none	—
$\text{gen}_3[1]: 1 \rightarrow 0$	Rule 106	Class 1
$\text{gen}_3[2]: 1 \rightarrow 0$	none	—
$\text{gen}_3[3]: 1 \rightarrow 0$	none	—
$\text{gen}_3[4]: 1 \rightarrow 0$	none	—

Findings: 8 of 10 perturbations yield *no* orbit-satisfying rule. The 2 that do yield simple (Class 1 or Class 3) non-universal rules. *Zero Class 4/universal rules* appear among any perturbed orbit. This 10-perturbation catalog is machine-certified in Lean 4 (zero `sorry`): theorem `orbit_perturbation_destroys_universality`, module `OrbitPerturbationCatalog.lean` (`ugp-lean`). Furthermore, `rule110_orbit_complete_isolation` certifies that $(\text{smGen}_2, \text{smGen}_3)$ is the unique 2-step orbit output of Rule 110 from smGen_1 over all 1024 possible orbit pairs, confirming that the SM orbit is isolated beyond single-bit perturbations.

Note on gen_1 perturbations. Perturbations of gen_1 are not tested in the same way as $\text{gen}_2/\text{gen}_3$ perturbations: gen_1 is the *initial condition* of the orbit, not a CA output, so perturbing it changes the initial data and defines a new orbit problem rather than a perturbation of the current one. The relevant question — whether perturbed initial conditions yield universal rules — is answered by the 10-perturbation sample above, which shows that specific perturbations of the *output* generations gen_2 and gen_3 yield no universal CA.

5.3 The algebraic explanation

Proposition 2.3 provides the algebraic explanation: the orbit path directly specifies the required minterm set as $\{1, 2, 3, 5, 6\}$. This is *exactly* Rule 110’s minterm set. Rules with different minterm sets (the perturbed-orbit selections) are adjacent sets $\{1, 3, 5, 6\}$, $\{1, 2, 5, 6\}$, etc., none of which produce Class 4 behavior.

Theorem 5.1 (Orbit–universality structural connection, $\mathbf{A}_{\text{Lean}}$). *The SM orbit constraints directly require the Rule 110 minterm set $\{1, 2, 3, 5, 6\}$. This minterm set uniquely enables Rule 110’s glider dynamics and computational universality. The orbit–universality connection is structural (via shared minterm structure), not coincidental. Among all weight-5 binary CA rules ($\binom{8}{5} = 56$ rules), exactly one satisfies the SM orbit with vacuum-transparency: Rule 110 [Lean-certified, `rule110_unique_weight5_orbit_satisfier`, `weight5_orbit_satisfier_count = 1`, `CUP4TotalParity.lean`]. The minterm set $\{1, 2, 3, 5, 6\}$ is Lean-proved to be exactly the set produced by the orbit [`minterm_set_as_finset`, zero `sorry`]. A physical-combinatorial equivalence follows (orbit_weight_dichotomy, machine-certified in Lean 4): among orbit-satisfying rules, vacuum-transparency is equivalent to Hamming weight 5.*

Remark 5.2. The three non-minterm neighborhoods $\{0, 4, 7\}$ — the binary triples not activated by the orbit — have direct physical interpretations: neighborhood 0 is the vacuum triple (all-zero, fixed point by vacuum-transparency), neighborhood 4 is the left-only excitation (contributes to orbit termination), and neighborhood 7 is the all-ones triple (maximal excitation). That these three neighborhoods are structurally excluded from the minterm set is Lean-certified: `rule110_non_minterm_set` (`CUP4TotalParity.lean`, zero `sorry`).

Remark 5.3 (Orbit-admissible function count and MDL uniqueness, $\mathbf{A}$). The SM orbit and the Rule 110 binary sublayer together impose 23 constraints on a $\mathbb{Z}_7$ CA function $f : \mathbb{Z}_7^3 \rightarrow \mathbb{Z}_7$ (15 from the 3-step

orbit, 8 from the binary sublayer; no overlaps). The remaining 320 neighborhoods are unconstrained. The orbit-admissible class therefore has $7^{320} \approx 10^{270}$ elements — a vast equivalence class of CA rules consistent with the SM particle spectrum. Among these, MDL-minimality (all 320 free neighborhoods $\rightarrow 0$) selects a unique element: f_{MDL} . This is the CA analogue of the PSC universe selection: the SM orbit creates an astronomically large admissible class, but the MDL principle uniquely identifies the physically realized CA rule. (Note: CUP-11c counts 7^{325} free completions rather than 7^{320} , because that theorem constrains only the 18 explicitly non-zero orbit+binary neighborhoods and does not impose the 5 gen_3 -to-vacuum zero outputs as separate constraints — those are orbit-required to output 0 but already match the MDL default, so they do not reduce the number of universal completions. The 7^{320} count here is the correct size of the fully orbit-admissible class.) (computationally verified, Python-verified: `orbit_admissible_count.py`.)

Remark 5.4 (CUP-4 as the binary anchor of the direct-interpolation lift, $\mathbf{A}_{\text{Lean}}$). CUP-4 is the certified binary anchor of a larger selection chain. The Direct-Interpolation Lift Theorem (machine-certified in Lean 4, zero `sorry`: `ugp_orbit_interpolation_lift`, with orbit provenance `gte_orbit_parity_provenance` and binary corollary `interpolation_lift_binary_corollary`, `ugp-lean`) shows that the same orbit data, placed as total-parity constraints on the $\mathbb{Z}_5$ family ring with vacuum transparency — itself MDL-equivalent among orbit satisfiers (`orbit_weight_dichotomy`, Theorem 5.1) — is full-rank over $\text{GF}(7)$ and forces exactly one multilinear rule in the full 7^8 multilinear class: the GTE polynomial $p(L, C, R) = C + R - CR - LCR$, with Rule 110 as its binary restriction. The accompanying sparsity floor (`rule110_lift_sparsity_floor`, `gf7_rule110_sparsity_floor`: any $\text{GF}(7)$ rule whose binary restriction is Rule 110 flattens to p and has at least 4 monomials, a floor that p itself realizes with exactly 4) makes multilinearity MDL-forced rather than declared. The ordering census (`orbit_chirality_census`) refines the rule-selectivity observation of §4: over all 120 family orderings, the vacuum-transparent survivor union is exactly the chiral pair {Rule 110, Rule 124}, with ordering reversal bijecting the Rule 110 orderings onto the Rule 124 orderings, so the canonical ordering of CUP-4 is the chirality gauge choice — consistent with, and grounding, the sublayer-consistency selection of Rule 110 over Rule 124 (§13). These statements are arithmetic-certificate results; no claim of the present paper changes — Theorem 2.2, the rule-selectivity count (Rules 110/111/124/125), Proposition 3.5, and the CUP-9 $\mathbb{Z}_5$ ring all stand verbatim. The full lift chain is developed in P49 [16] and P48 [10].

Remark 5.5 (Self-simulating embeddings). Rule 110 Turing-universality [1, 2] implies that a Rule 110 simulator can be encoded as a Rule 110 initial condition, realizing the rule as a self-simulating cellular automaton embedding (Rule 110-in-Rule 110). This is the Rule 110 analogue of Rendell’s Game-of-Life-in-Game-of-Life universal Turing machine construction [4]. In the UGP multi-level CA hierarchy, each computational level is precisely such a self-embedded copy, grounding the hierarchical level structure in the constructive mechanism provided by Cook universality. The ether temporal period 7 on aligned rings is a verified building block ($\mathbf{A}$). Naive single-tape macro encodings fail systematic simulation tests; a full constructive Rule 110-in-Rule 110 embedding awaits Cook (2009) block assembly. A phased CTS milestone reproduces the Lean-certified init tape with ossifier support; one-step `cook_min_len6` post-decode readback at the origin cell is confirmed, but central-region data-cone readback (blocks C–G) remains negative (0/6 slots), consistent with Lean witnesses—full FCA substrate not claimed until Cook §4 collision certificates close the gap.

5.4 CA sufficiency for Standard Model physics (Computational Physics Principle)

Sufficiency is $\mathbf{A}_{\text{Lean}}$ partial: (i) the Algebraic Lifting Theorem (`LiftingTheorem.lean`) promotes beable-level certified properties to $[\mathbf{D}]$ -weighted observables; (ii) Φ_{MDL} Turing universality is certified via the Cook route composed with the proved stepwise Φ_{MDL} -Rule 110 simulation (`phimdl_turing_universal`, $\mathbf{A}_{\text{Lean}}$ conditional; one named axiom: `cook_rule110_simulates_computable`); the $\text{GF}(7)$ polynomial identity (`rule110_z7_poly_rep`, zero `sorry`) independently certifies finite NAND functional completeness, a distinct and weaker result; (iii) computation/transputation division follows the OIR framework. **Necessity** (discrete spacetime $\Rightarrow$ CA) remains conjectural. Cook’s theorem [1] remains the load-bearing Turing-universality result; it is not a corollary of the finite-completeness algebraic identity.

5.5 MDL-minimal canonical particle encoding

MDL selects a *unique canonical encoding* at each certified level: rule (`mdl_robustness_z7`, `fmdl_mdl_uniqueness`); winding class (orbit closure $\{0, 1, 2, 5\}$, photon as unique ether fixed point); beable ($\text{gen}_1/2/3$ ring plus GTE triple discrimination, `GTPNeutralDiscrimination.lean`). Turing-equivalent encodings may exist; MDL picks the canonical one. Part B selection (non-minimal encodings $\Rightarrow$ higher effective mass) remains **A/D** prose. Explicit spatial Rule 110 glider species per particle are not claimed (2-ZGM hard form deferred: distinguishing massless gliders from massive orbit beables at the Rule 110 spatial layer requires additional CA-dynamics input beyond current scope). The mesoscopic form (2-ZGM-MES) is **closed** ($\mathbf{A}_{\text{Lean}}$): the f_{MDL} forward chain $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3 \rightarrow \text{vacuum}$ assigns a unique orbital position $I_{\text{pos}}(\text{gen}_i) = i$ ($i \in \{1, 2, 3\}$) and $I_{\text{pos}}(\text{vacuum}) = 4$, with all four values distinct; machine-certified in `ZGMMesInvariant.lean` (ugp-lean, zero sorry).

6 CUP-11: Mod-7 Structure and the Mirror Dual

The binary Rule 110 structure (CUP-4) operates at the parity-projection layer of the SM generation orbit.

A deeper layer emerges when the GTE triples are examined modulo 7: the number $\delta = 7$ that characterizes the UGP ridge structure naturally lifts the binary orbit to a $\mathbb{Z}_7$ orbit, distinguishes the SM branch from a mirror-dual dark-sector branch, and admits a computationally universal $\mathbb{Z}_7$ CA.

6.1 The mod-7 orbit

Theorem 6.1 (CUP-11a, **A/D**). *The mirror-dual $\mathbb{Z}_2$ branch (defined by swapping the divisor pair $b_2 = 42 \leftrightarrow q_2 = 24$) is invisible to mod-2 parity but visible mod 7. The algebraic identities $b_2 = \delta \times N_c! = 42$ and $q_2 = (N_c + 1)! = 24$ (where $\delta = 7$ and $N_c = 3$) hold exactly. The mod-7 residue $b_2 \bmod 7 = 0$, $q_2 \bmod 7 = 3 = N_c$, giving the candidate dark charge $Q_{\text{dark}} = N_c \bmod 7$.*

Theorem 6.2 (CUP-11b, $\mathbf{A}_{\text{Lean}}$). *A $\mathbb{Z}_7$ CA rule $f : \mathbb{Z}_7^3 \rightarrow \mathbb{Z}_7$ realizing the $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ orbit exists with 1730 of 1920 possible orderings conflict-free.¹ The totalistic special case has rule $g = [6, 5, 6, 3, 3, 6, 3]$ (Wolfram Class 2); g is a structurally related totalistic CA over $\mathbb{Z}_7$ but does not realize the SM orbit. The $\mathbb{Z}_7$ sum conservation result (Lean-certified, zero sorry) holds for the MDL-minimal rule f_{MDL} : the $\mathbb{Z}_7$ sum is conserved at gen_1 (sum 4 = 4) and broken at gen_2 and gen_3 ($4 \neq 3, 3 \neq 0$). [`cup11b_z7_sum_conservation`, `CUP3DUniqueness.lean`]*

Remark 6.3 (Complete characterization of $\mathbb{Z}_7$ -sum-4-conserving states, $\mathbf{A}_{\text{Lean}}$). Among all $7^5 = 16,807$ states on the $\mathbb{Z}_5$ ring, exactly 10 conserve the $\mathbb{Z}_7$ sum at value 4: the 5 cyclic rotations of $\text{gen}_1 = [1, 5, 2, 2, 1]$ and the 5 cyclic rotations of $[0, 2, 5, 2, 2]$. Gen_1 therefore belongs to the rarest conservation class: only $10/16,807 \approx 0.06\%$ of all states conserve the $\mathbb{Z}_7$ sum at the gen_1 value. This complete characterization is Lean-certified: `cup11b_z7_sum4_conserving_count` (count = 10, `native_decide`) and `cup11b_z7_sum4_conserving_characterization` (explicit membership characterization), both in `CUP3DUniqueness.lean`, zero sorry.

Remark 6.4 (Physical coherence of $\mathbb{Z}_7$ sum conservation, $\mathbf{A}_{\text{Lean}}$). The conservation-breaking pattern is coherent with the generation stability hierarchy: the stable first generation is simultaneously a Garden of Eden state under f_{MDL} and the unique $\mathbb{Z}_7$ -sum-conserving generation. $\mathbb{Z}_7$ sum conservation is a property of the arithmetic ground state; first-generation particles, being stable (infinite lifetime), satisfy it; heavier unstable generations do not.

Theorem 6.5 (CUP-11c, $\mathbf{A}_{\text{Lean}}$). *The mod-7 CA rule can be made computationally universal (Turing-complete simulation of Rule 110 on binary initial conditions) while respecting the SM generation orbit. The MDL-minimal universal completion has 18 fixed neighborhoods and 7^{325} free completions, all of which simulate Rule 110 on binary inputs.*

Proof. `cup11c_universal_mod7_CA_exists`: proved by `decide`; see `CUP11ModSeven.lean`. Zero sorry. $\square$

¹The figure 1920 counts candidate neighborhood-to-output assignment orderings in the $\mathbb{Z}_7$ orbit analysis of the 5-cell periodic ring; the exact combinatorial breakdown is implemented in `CUP11ModSeven.lean`.

6.2 The MDL-minimal $\mathbb{Z}_7$ function

The mod-7 generation orbit assigns a $\mathbb{Z}_7$ state value to each SM particle family in each generation; these values are the GTE triple components modulo 7. Table 4 records the orbit state values, which serve as the input/output entries for the 10 fixed orbit neighborhoods of f_{MDL} . The SM winding numbers W (from P22 [22], also shown) are a distinct $\mathbb{Z}_7$ quantity encoding interaction charge, not orbit position; see (2) and §8 for the coupling that connects them.

Table 4: The mod-7 generation orbit: $\mathbb{Z}_7$ state values for each SM family across three generations. The orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ under f_{MDL} gives the 10 fixed $\mathbb{Z}_7$ neighborhoods that define the orbit constraints. The SM winding W (from P22 [22]) encodes interaction charge and is a *distinct* $\mathbb{Z}_7$ quantity from the orbit state values.

Family	$\text{gen}_1 (\mathbb{Z}_7)$	$\text{gen}_2 (\mathbb{Z}_7)$	$\text{gen}_3 (\mathbb{Z}_7)$	SM winding W
e^-	1	2	5	$-3 \equiv 4 \pmod{7}$
u	5	5	6	+2
d	2	2	5	$-1 \equiv 6 \pmod{7}$
ν_R	2	0	3	0
ν_L	1	2	5	0
Vector	[1, 5, 2, 2, 1]	[2, 5, 2, 0, 2]	[5, 6, 5, 3, 5]	—

The MDL-minimal completion $f_{\text{MDL}} : \mathbb{Z}_7^3 \rightarrow \mathbb{Z}_7$ is defined by 18 fixed neighborhoods (10 orbit + 8 Rule 110 binary) and 325 free neighborhoods set to 0. Table 5 lists the 10 orbit neighborhood entries cell by cell.

Table 5: The 10 $\mathbb{Z}_7$ orbit neighborhood entries of f_{MDL} : the cell-by-cell transitions $\text{gen}_1 \rightarrow \text{gen}_2$ and $\text{gen}_2 \rightarrow \text{gen}_3$ on the periodic $\mathbb{Z}_5$ ring. These 10 entries, combined with 8 Rule 110 binary entries (one for each binary triple), constitute the 18 fixed neighborhoods of f_{MDL} ; all remaining 325 neighborhoods map to 0 by MDL-minimality.

Step	Cell	(l, c, r)	f_{MDL} output	Note
$\text{gen}_1 \rightarrow \text{gen}_2$	0	(1, 1, 5)	2	
$\text{gen}_1 \rightarrow \text{gen}_2$	1	(1, 5, 2)	5	
$\text{gen}_1 \rightarrow \text{gen}_2$	2	(5, 2, 2)	2	
$\text{gen}_1 \rightarrow \text{gen}_2$	3	(2, 2, 1)	0	
$\text{gen}_1 \rightarrow \text{gen}_2$	4	(2, 1, 1)	2	
$\text{gen}_2 \rightarrow \text{gen}_3$	0	(2, 2, 5)	5	
$\text{gen}_2 \rightarrow \text{gen}_3$	1	(2, 5, 2)	6	quark flavor-change neighborhood
$\text{gen}_2 \rightarrow \text{gen}_3$	2	(5, 2, 0)	5	
$\text{gen}_2 \rightarrow \text{gen}_3$	3	(2, 0, 2)	3	W^+ emission neighborhood
$\text{gen}_2 \rightarrow \text{gen}_3$	4	(0, 2, 2)	5	

The neighborhood $(2, 5, 2) \rightarrow 6$ ($\text{gen}_2 \rightarrow \text{gen}_3$, cell 1) is the same neighborhood responsible for quark flavor change ($u + \bar{u} \rightarrow d$) in the interaction table (§9), and $(2, 0, 2) \rightarrow 3$ implements W^+ emission. The orbit neighborhoods are thus not only generation-transition data but also encode the fundamental interaction vertices: the same CA table entries that propagate the generation orbit also implement the SM charged-current interactions.

The Lean theorems `fmdl_quark_flavor_change` ($f_{\text{MDL}}(2,5,2) = 6$) and `fmdl_w_emission` ($f_{\text{MDL}}(2,0,2) = 3$) are proved by `decide` in `CUP3DUniqueness.lean` (zero sorry).

Theorem 6.6 (d-Quark Production Uniqueness, $\mathbf{A}_{\text{Lean}}$). *Among all $343 = 7^3$ possible input neighborhoods, exactly one produces the d-quark winding ($\mathbb{Z}_7 = 6$) under f_{MDL} : the triple $(2,5,2)$ (u-quark center flanked by u-quarks). Formally, for all $l, c, r \in \mathbb{Z}_7$:*

$$f_{\text{MDL}}(l, c, r) = 6 \iff l = 2 \wedge c = 5 \wedge r = 2.$$

Lean-certified: `dquark_unique_production_vertex`, `dquark_unique_preimage`, `dquark_preimage_count` (`GUTStructure.lean` §44, `native_decide`, zero sorry). This uniqueness is the CA-level structural analog of CKM suppression: quark flavor change $u \rightarrow d$ occurs at exactly 1 of 343 neighborhoods at each single CA evaluation step, corresponding to the single orbit vertex $(2,5,2) \rightarrow 6$ that implements the $\text{gen}_2 \rightarrow \text{gen}_3$ generation transition.

It has been proved in Lean 4 that f_{MDL} never outputs the value 4 (electron winding number $W = -3 \equiv 4 \pmod{7}$):

Theorem 6.7 (Electron exclusion, $\mathbf{A}_{\text{Lean}}$). $\forall l, c, r \in \mathbb{Z}_7 : f_{\text{MDL}}(l, c, r) \neq 4$.

Proof. `fmdl_never_outputs_4`: proved by `decide` over all 343 neighborhoods. See `CUP3DUniqueness.lean`. Zero sorry. $\square$

Corollary 6.8. *The electron winding value $Z_7 = 4$ can only arise from cross-dimensional $\mathbb{Z}_7$ -addition interactions ($v_1 + v_2 \pmod{7} = 4$), never from any single-axis f_{MDL} evaluation. The electron is structurally a cross-dimensional particle in the f_{MDL} framework.*

Corollary 6.9 (Structural Lepton Number Conservation, $\mathbf{A}_{\text{Lean}}$). *Since $4 \notin \text{range}(f_{\text{MDL}})$ (Theorem 6.7), the CA substrate cannot produce winding-4 excitations from any input triple. As a consequence, any tape that contains no winding-4 cells cannot develop any under f_{MDL} dynamics for any number of evolution steps. Combined with the center-decay result (`winding_4_maps_to_vacuum`), winding 4 is both uncreatable as a direct output and unstable as a center: it is a structural boundary condition of the substrate, not a dynamical product. This is a structural derivation of lepton non-creation at the CA level — the charge conservation of winding 4 is not a separately postulated symmetry but follows directly from the definition of f_{MDL} and MDL minimality. Lean-certified: `lepton_production_impossible`, `lepton_number_conservation`, `lepton_number_preserved_from_L0_initial`, `lepton_number_conservation_summary` (`GUTStructure.lean` §46, zero sorry).*

Remark 6.10 ($\mathbb{Z}_7$ conservation landscape, $\mathbf{A}_{\text{Lean}}$). The complete $\mathbb{Z}_7$ sum conservation landscape of f_{MDL} over all $7^5 = 16,807$ states is:

Sum value v	States with sum v conserved	Fraction
$v = 0$	2026	12.05%
$v = 1$	100	0.595%
$v = 2$	80	0.476%
$v = 3$	45	0.268%
$v = 4$	10	0.0595% (SM orbit; rarest non-zero class)
$v = 5$	80	0.476%
$v = 6$	30	0.179%

The SM orbit sum ($v = 4$) is the uniquely sparsest non-zero conservation class: only 10 of 16,807 states are sum-4 conserving: the 5 cyclic rotations of gen_1 and 5 rotations of $[0, 2, 5, 2, 2]$ (Lean-certified: `cup11b_z7_sum4_conserving_characterization`, `CUP3DUniqueness.lean`, zero sorry). Lean certs: `z7_conservation_count_table`, `z7_sum4_uniquely_sparse` (`CUP3DUniqueness.lean` §8, zero sorry). Note that gen_3 does not conserve its $\mathbb{Z}_7$ sum ($3 \rightarrow 0$ upon cascade): the orbit sum trajectory is $4 \rightarrow 4 \rightarrow 3 \rightarrow 0$, monotonically decreasing after gen_2 . Lean cert: `z7_gen3_not_sum_conserving` (`CUP3DUniqueness.lean` §8).

Remark 6.11 (f_{MDL} nonzero count and Higgs cascade depth, $\mathbf{A}_{\text{Lean}}$). The MDL-minimal function f_{MDL} produces nonzero output on exactly 14 of the 343 possible (l, c, r) input neighborhoods: this equals $c_H + 1 = 13 + 1$, where $c_H = 13$ is the GTE Higgs cascade depth. Equivalently, the count decomposes as $N_{\text{gen}} + 2N_{\text{fam}} + 1 = 3 + 10 + 1 = 14$. Lean-certified: `GUTStructure.fmdl_count_eq_chiggs_plus_one` (`GUTStructure.lean` §9, zero sorry).

Remark 6.12 (f_{MDL} vertex table partition, $\mathbf{A}_{\text{Lean}}$). The 343 neighborhoods of f_{MDL} partition into 329 transparent entries (output 0) and 14 non-zero interaction entries. The 14 non-zero neighborhoods further partition as: 9 orbit-sourced (encoding the $\text{gen}_1 \rightarrow \text{gen}_2$ and $\text{gen}_2 \rightarrow \text{gen}_3$ orbital transitions) and 5 Rule 110-sourced (anti- d production from the binary sublayer). All 14 outputs lie in $\{0, 1, 2, 3, 5, 6\} \subset \mathbb{Z}_7$; the value $Z_7 = 4$ (W^-/e^- winding) is arithmetically excluded from all 343 outputs (Theorem 6.7; Lean-certified: `fmdl_never_outputs_4`, zero sorry).

Remark 6.13 (f_{MDL} as a perfect code, $\mathbf{A}$). The function f_{MDL} is a *perfect code*: its 14 non-zero outputs achieve the minimum count consistent with orbit-consistency and the Rule 110 binary sublayer. The 14 non-zero neighborhoods decompose as 9 orbit-forced (neighborhoods in the SM orbit with required non-zero output) and 5 binary-forced (Rule 110 binary neighborhoods with non-zero output). These two sets are necessarily disjoint — orbit neighborhoods carry $\mathbb{Z}_7$ values ≥ 2 while binary neighborhoods carry values in $\{0, 1\}$ — so the lower bound is additive: at least $9 + 5 = 14$ non-zero outputs are required. f_{MDL} , as the MDL-minimal orbit-consistent function, sets all 320 free neighborhoods to 0 and achieves exactly this lower bound. The count 14 is machine-certified by `fmdl_nonzero_count_14` (`GUTStructure.lean` §9, `native_decide`, zero sorry).

Remark 6.14 ($b_{\text{sum}} = 2 \times 3 \times 5 \times 13$ — all SM structural numbers, $\mathbf{A}_{\text{Lean}}$). The sum of SM generation N -values $b_1 + b_2 + b_3 = 73 + 42 + 275 = 390$ factorizes as $390 = 2 \times 3 \times 5 \times 13$, encoding all four key structural parameters of the SM in f_{MDL} : the binary Rule 110 basis (2), the generation count $N_{\text{gen}} = 3$, the family ring size $N_{\text{fam}} = 5$, and the Higgs cascade depth $c_H = 13$. The Weinberg numerator ($N_{\text{gen}} = 3$) and denominator ($c_H = 13$) are both co-determined prime factors of b_{sum} . Additional structural identities: $N_{\text{gen}} + c_H = 3 + 13 = 16 = 2^4$ (the GUT-to-EW ridge subtraction constant). Lean-certified: `b_sum_is_product` ($b_{\text{sum}} = 2N_{\text{gen}}N_{\text{fam}}c_H$), `weinberg_numerator_in_bsum`, `weinberg_denominator_in_bsum`, `weinberg_ratio_from_bsum`, `nw_plus_chiggs_eq_pow2` (`GUTStructure.lean` §16, zero sorry).

Remark 6.15 (Goldstone cascade formula, $\mathbf{A}_{\text{Lean}}$). The EW boson GTE cascade depths obey a Goldstone counting formula: $c_P = c_H - d_P$, where $c_H = 13$ is the Higgs cascade depth and $d_P \in \{0, 1, 2\}$ is the number of Goldstone degrees of freedom absorbed by particle P . Explicitly: $c_{H^0} = 13$ ($d = 0$: scalar, no Goldstone absorbed); $c_Z = 12$ ($d = 1$: one longitudinal mode absorbed); $c_{W^\pm} = 11$ ($d = 2$: two transverse modes absorbed). Each unit c -step encodes one absorbed Goldstone degree of freedom. Lean-certified: `goldstone_cascade_higgs` ($c_{H^0} = 13$), `goldstone_cascade_z` ($c_Z = 12$), `goldstone_cascade_w` ($c_{W^\pm} = 11$), `goldstone_cascade_formula` ($c_P = c_H - d_P$) (`EWBosonStructure.lean` §5, zero sorry).

Remark 6.16 (f_{MDL} uniform fixed point: photon as CA vacuum, $\mathbf{A}_{\text{Lean}}$). Among all $\mathbb{Z}_7$ winding values $k \in \{0, \dots, 6\}$, the vacuum ($k = 0$) is the unique value satisfying $f_{\text{MDL}}(k, k, k) = k$. For $k = 1$: the Rule 110 binary constraint forces $f_{\text{MDL}}(1, 1, 1) = 0$. For $k \geq 2$: MDL-minimality assigns 0 to all free diagonal neighborhoods. The photon ($Z_7 = 0$) is thus the unique winding value that a uniform homogeneous background reproduces unchanged — the CA’s quiescent ether. This closes the structural “why” of the photon’s GTE-triple absence: γ has $K_{\text{MDL}} = 0$ because it *is* the CA vacuum. Lean-certified: `fmdl_unique_uniform_fixed_point`, `photon_is_ca_ether`, `fmdl_uniform_fp_uniqueness_count` (`CUP3DUniqueness.lean` §7d, zero sorry).

Theorem 6.17 ($\mathbb{Z}_7$ charge conjugation structure, $\mathbf{A}_{\text{Lean}}$). Define the $\mathbb{Z}_7$ charge conjugation map $C(v) = (7 - v) \bmod 7$. The following hold, all Lean-certified in `Z7ChargeConjugation.lean` (zero sorry):

1. $C(v) = v \iff v = 0$: the vacuum is the unique self-conjugate value [`z7_conj_self_conjugate_iff_zero`].
2. $C(C(v)) = v$ for all $v \in \mathbb{Z}_7$: C is an involution [`z7_conj_involution`].
3. $v + C(v) \equiv 0 \pmod{7}$ for all v : conjugate pairs sum to zero [`z7_conj_sum_zero`].

4. The three non-trivial conjugate pairs are: (1, 6) (anti-d/d), (2, 5) (u/anti-u), (3, 4) ($W^+/W^-/e^-$) [`z7_conj_pairs`].

The f_{MDL} output range $\{0, 1, 2, 3, 5, 6\}$ is not closed under C : the value $Z_7 = 3$ (W^+) is in the range, but $C(3) = 4$ (W^-/e^-) is not [`fmdl_output_range_not_conj_closed`]. The pair ($W^+, W^-/e^-$) is therefore the unique maximally CP-asymmetric pair: f_{MDL} directly produces W^+ (exactly one neighborhood (2, 0, 2); `fmdl_w_plus_unique_neighborhood`) and never produces W^-/e^- [`fmdl_conj_pair_asymmetry_unique`, machine-certified in Lean 4]. Preimage counts: $0 \rightarrow 329$, $1 \rightarrow 5$, $2 \rightarrow 3$, $3 \rightarrow 1$, $4 \rightarrow 0$, $5 \rightarrow 4$, $6 \rightarrow 1$ [`fmdl_preimage_counts`, `native_decide`].

Remark 6.18 (W^+/W^- beable asymmetry and amplitude conjugation; $\mathbf{A}_{\text{Lean}}/\mathbf{A}/\mathbf{D}$). At the CA level, f_{MDL} never outputs $w = 4$ (`fmdl_never_outputs_4`, zero sorry), making W^- (winding 4) unproducible from any CA configuration. The W^+ -only producibility of f_{MDL} — W^+ appears in $\text{range}(f_{\text{MDL}})$ while W^- does not—constitutes a CA-level CP asymmetry: the CA beable dynamics are maximally W^+ -biased. The W^- boson and all W^- -mediated processes arise as quantum Hilbert-space constructions from the W^+ sector via amplitude conjugation (computationally confirmed), not as independent CA beables. The physical consequences for matter-antimatter asymmetry are analyzed in detail in [11].

7 CUP-12: The MDL-Minimal Universal $\mathbb{Z}_7$ CA as Candidate Physics Rule

CUP-11c establishes that a computationally universal $\mathbb{Z}_7$ CA satisfying the SM orbit exists, with 325 free neighborhoods. Among all valid completions, the MDL principle (Occam’s Razor applied to CA description length) selects a unique one: f_{MDL} , with all 325 free neighborhoods set to zero.

This section establishes CUP-12 — the MDL-minimal $\mathbb{Z}_7$ CA — and proves its key structural properties in Lean.

7.1 The f_{MDL} attractor structure

The totalistic rule $g = [6, 5, 6, 3, 3, 6, 3]$ has a period-2 attractor ($[3, 3, 3, 3, 3] \leftrightarrow [6, 6, 6, 6, 6]$). Generations gen_2 and gen_3 are in the basin of attraction of this attractor ($\text{gen}_2 = [2, 5, 2, 0, 2]$ reaches $[3, 3, 3, 3, 3]$ in 5 steps under g); gen_1 is not in the attractor basin. Under g , the heavy generations lie in the basin of the arithmetic oscillator; the lightest generation, as a Garden of Eden state under f_{MDL} , stands outside this basin, consistent with the picture that stability corresponds to arithmetic isolation from dynamical attractors.

Theorem 7.1 (Garden of Eden stability, $\mathbf{A}_{\text{Lean}}$). *Under the MDL-minimal CA f_{MDL} :*

1. $\text{gen}_1 = [1, 5, 2, 2, 1]$ is a Garden of Eden state: it has zero predecessors across all $7^5 = 16,807$ possible $\mathbb{Z}_7$ configurations. No $\mathbb{Z}_7$ state evolves into gen_1 ; gen_1 can only exist as an initial condition. [`fmdl_gen1_is_garden_of_eden`, Lean-proved via `native_decide`, zero sorry, `CUP3DUniqueness.lean`]
2. $\text{gen}_3 = [5, 6, 5, 3, 5]$ has predecessors (the generation orbit is non-GoE above gen_1).

Under the totalistic rule g alone: gen_1 and $\text{gen}_2 = [2, 5, 2, 0, 2]$ are Garden of Eden states; gen_3 has 9 predecessors (computationally verified).

The Garden of Eden structure gives a physical explanation of generation-1 stability: first-generation particles (electron, up quark, down quark, electron neutrino) are stable because the CA dynamics cannot produce them from any heavier state. Their stability is structural (unreachability), not thermodynamic.

Remark 7.2 (All gen_1 rotations are Garden of Eden states, $\mathbf{A}_{\text{Lean}}$). The GoE property extends to all five particle families of the first generation. All five cyclic rotations of $\text{gen}_1 = [1, 5, 2, 2, 1]$ on the $\mathbb{Z}_5$ ring are Garden of Eden states under `fmdl_step5`: each has zero predecessors among all $7^5 = 16,807$ states. The $\mathbb{Z}_5$ ring symmetry of the CA directly manifests as a 5-fold GoE family structure: the first generation is rotationally symmetric in both its particle content ($N_{\text{fam}} = 5$) and its CA stability property. This unifies two independent counting results — ring size and GoE stability — into a single CA-arithmetic theorem. Lean-certified: `fmdl_gen1_all_rotations_are_goe`, `fmdl_gen1_all_rotations_no_predecessor` ($5 \times 16,807$ cases, `native_decide`, zero sorry; `GoEStabilityHierarchy.lean` §6).

Remark 7.3 ($\mathbb{Z}_5$ equivariance of f_{MDL} , $\mathbf{A}_{\text{Lean}}$). The MDL-minimal function f_{MDL} is exactly $\mathbb{Z}_5$ -equivariant: for any state $v \in \mathbb{Z}_7^5$ and any cyclic rotation $k \in \mathbb{Z}_5$, $f_{\text{MDL}}(\sigma_k(v)) = \sigma_k(f_{\text{MDL}}(v))$. The five SM particle families $[e^-, u, d, \nu_R, \nu_L]$ are thus related by exact $\mathbb{Z}_5$ ring symmetry in the CA dynamics. Verified over all 84,035 cases ($7^5 \times 5$, `native_decide`, zero sorry). Lean-certified: `CUP3D.fmdl_z5_equivariant` (`CUP3DUniqueness.lean` §10, `ugp-lean`).

Remark 7.4 (GoE property as Part 1 of a 6-property causal isolation). The Garden of Eden property of gen_1 is Part 1 of a 6-property CA-structural characterization of the SM generation orbit: the orbit is the unique chain in $\mathbb{Z}_7^5$ (and $\mathbb{Z}_2^5$) simultaneously satisfying GoE, unique predecessor chain, $\mathbb{Z}_7$ sum conservation pattern, and global $\mathbb{Z}_2$ isolation. Lean-certified as `sm_orbit_complete_causal_isolation` (`SMOrbitCausalIsolation.lean`, zero sorry). The full 6-part master theorem is developed in a companion paper (see §13 for the forward-reference to the deeper-consequences treatment).

Theorem 7.5 (Orbit-Closure of Self-Propagating Centers, $\mathbf{A}_{\text{Lean}}$). *A winding $w \in \mathbb{Z}_7$ is self-propagating under f_{MDL} if there exist $l, r \in \mathbb{Z}_7$ such that $f_{\text{MDL}}(l, w, r) = w$. The set of self-propagating windings is exactly the orbit-closure of the SM generation orbit:*

$$\{w \in \mathbb{Z}_7 : \exists l, r, f_{\text{MDL}}(l, w, r) = w\} = \{0, 1, 2, 5\}.$$

That is, w is self-propagating if and only if w appears in the SM generation orbit or background ($w \in \{0, 1, 2, 5\}$). The partition $\{0, 1, 2, 5\}$ vs. $\{3, 4, 6\}$ is not arbitrary: it is entirely determined by the orbit structure together with MDL minimality, which sets all free diagonal neighborhoods to zero, eliminating any additional self-propagating witnesses. Lean-certified: `orbit_winding_set`, `self_propagating_iff_orbit_winding`, `orbit_closure_theorem` (`GUTStructure.lean` §44, `native_decide`, zero sorry).

7.2 CUP-12 and the cross-sector rule

Theorem 7.6 (CUP-12, $\mathbf{A}/\mathbf{D}/\mathbf{A}_{\text{Lean}}$). *The MDL-minimal universal $\mathbb{Z}_7$ CA embedding both SM and dark sector generation orbits (f_{CROSS}) has:*

- 27 fixed neighborhoods: SM 10 + dark 9 + Rule 110 8 (zero conflicts)
- Description length: 76 bits (computationally confirmed)
- Wolfram class: Class 3 on random $\mathbb{Z}_7$ initial conditions; Class 4 behavior (Rule 110 simulation) on binary initial conditions

The MDL-minimality content — that f_{MDL} is the unique completion with all free neighborhoods set to 0 — is Lean-certified (zero sorry); see Theorem 8.3.

The 76-bit description length of f_{CROSS} is the first *quantitative* MDL measure of the complexity of the SM + dark sector orbital structure. The 76 bits count the compact binary description of f_{CROSS} 's 27 fixed non-trivial neighborhoods (SM orbit 10 + dark orbit 9 + Rule 110 binary 8): each neighborhood entry requires specifying an index (3 values in $\mathbb{Z}_7$, i.e. $\lceil \log_2 7 \rceil = 3$ bits) and an output value ($\lceil \log_2 7 \rceil = 3$ bits), giving approximately $27 \times (9 + 3)/2 \approx 76$ bits in a compact packing; the exact count is computed by `t_cup12_cross_sector.py` (`canonical_run/`).

8 The Unique $\mathbb{Z}_7$ Cross-Dimensional Coupling

8.1 Setup and motivation

The CUP-4/11 results establish the winding assignment $W(v) = v$ for SM particles in $\mathbb{Z}_7$. P22 [22] (Lean-proved) establishes winding conservation as the vertex law. A natural question: under these two independent results, is the cross-dimensional $\mathbb{Z}_7$ coupling uniquely determined?

8.2 Coupling uniqueness theorem

Theorem 8.1 (P22 coupling uniqueness [22], $\mathbf{A}_{\text{Lean}}$). *The unique function $C : \mathbb{Z}_7 \times \mathbb{Z}_7 \rightarrow \mathbb{Z}_7$ satisfying P22 winding conservation [22] (Theorem 6.5, §6) with the CUP-4/11 winding assignment $W(v) = v$ is $\mathbb{Z}_7$ addition: $C(v_1, v_2) = (v_1 + v_2) \bmod 7$.*

Proof. P22’s biconditional forces $W(C(v_1, v_2)) = W(v_1) + W(v_2)$. Substituting $W = \text{id}$: $C(v_1, v_2) = (v_1 + v_2) \bmod 7$. Each of the 49 entries is uniquely determined. `p22_z7_coupling_unique`: proved by function extensionality in `CUP3DUniqueness.lean`. Zero sorry. $\square$

Corollary 8.2 (SM winding-conservation vertex rules recovered). *Under P22 + CUP-4/11, the complete SM winding-conservation vertex table is recovered: $u + W^- \rightarrow d$ ($\mathbb{Z}_7$: $2 + 4 = 6$), $d + W^+ \rightarrow u$ ($6 + 3 = 2$), $e^- + W^+ \rightarrow \nu$ ($4 + 3 = 0$), $e^- + \gamma \rightarrow e^-$ ($4 + 0 = 4$), etc. All 49 cross-dimensional pairs produce correct winding-conserving outputs.*

8.3 MDL-minimality and the CUP-12 upgrade

The mathematical content of CUP-12 (Theorem 7.6) is Lean-certified (zero sorry):

Theorem 8.3 (CUP-12 MDL-minimality, $\mathbf{A}_{\text{Lean}}$). *f_{MDL} assigns output 0 to all 325 free (non-fixed) neighborhoods, making it the MDL-minimal completion of the orbit+Rule 110 constraints.*

Proof. `fmdl_zero_on_free_neighborhoods`: $\forall l, c, r : \mathbb{Z}_7$, if (l, c, r) is not one of the 18 fixed neighborhoods, then $f_{\text{MDL}}(l, c, r) = 0$. Proved by `decide`. Combined with `fmdl_nonzero_only_at_fixed` (by `decide`), this establishes MDL-minimality. Zero sorry. $\square$

Remark 8.4 (Upgrading CUP-12). CUP-12 is a mathematical claim (the MDL-minimality of f_{MDL} , Lean-certified by Theorem 8.3) combined with the scientific methodology principle that Nature selects the simplest consistent rule (Occam’s Razor). The mathematical content is Lean-proved: f_{MDL} assigns output 0 to all 325 free neighborhoods and is the unique CA rule satisfying the orbit+Rule 110 constraints with the minimum number of non-zero entries. The physical identification of f_{MDL} as the CA governing the SM generation sector follows from the MDL principle as a working hypothesis, standard in information-theoretic physics. Whether Nature in fact selects the MDL-minimal rule, rather than one of the 7^{325} other valid completions, is a physical question that goes beyond the mathematical statement of CUP-11c and MDL-minimality alone.

Remark 8.5 (MDL selection stability at all orbit depths, $\mathbf{A}$). Exhaustive computation on the $\mathbb{Z}_2^5$ ring (5-cell binary, 32 states) confirms that the MDL identification of Rule 110 holds at orbit depths $k \in \{3, 4, 5\}$. At every tested depth, every Rule 110-consistent orbit leaves $(0, 0, 0)$ as the unique free neighborhood; MDL sets $(0, 0, 0) \rightarrow 0$, uniquely selecting Rule 110. The forced minterms $\{1, 2, 3, 5, 6\}$ and the single MDL-free minterm $(0, 0, 0)$ are structural properties of the Rule 110 rule table, not artifacts of the depth-3 orbit length. Depth-4 and depth-5 orbits exist; accordingly, the derivation of $N_{\text{gen}} = 3$ rests on $\mathbb{Z}_7$ arithmetic (`fmdl_ngen_equals_three`, `CUP3DPSCUnification.lean`, zero sorry), not on counting binary free neighborhoods as a function of depth. Script: `orbit_depth_free_minterms.py` (computationally verified, exhaustive on $\mathbb{Z}_2^5$).

Remark 8.6 ($\mathbb{Z}_7$ neighborhood count and MDL robustness, $\mathbf{A}_{\text{Lean}}$). The $\mathbb{Z}_7$ neighborhood structure of f_{MDL} is Lean-certified in `GUTStructure.lean` §28 (zero sorry):

- `z7_fixed_neighborhood_count`: exactly 18 of the $343 = 7^3$ possible neighborhoods satisfy `isFixedNeighborhood` — 10 from the SM generation orbit transitions and 8 from the binary Rule 110 sublayer.
- `z7_free_neighborhood_count`: exactly 325 neighborhoods are unconstrained. (The orbit also constrains 5 $\text{gen}_3 \rightarrow \text{vacuum}$ neighborhoods whose required output is 0, matching the MDL default; the fully orbit-admissible class is therefore 7^{320} , as noted in Remark 5.3.)

- `mdl_robustness_z7`: any $\mathbb{Z}_7$ CA function satisfying both the orbit constraints (`isFixedNeighborhood`) and MDL-minimality (all free neighborhoods $\rightarrow 0$) must equal f_{MDL} .

The `mdl_robustness_z7` theorem is the machine-certified form of MDL uniqueness: f_{MDL} is not merely the simplest valid completion but provably the *only* MDL-minimal orbit-admissible $\mathbb{Z}_7$ CA rule. This upgrades the Python-verified count of Remark 5.3 to a Lean-certified result (zero `sorry`). Total non-zero f_{MDL} outputs: 14 (Remark 6.11), decomposing as 9 from orbit transitions (4 from $\text{gen}_1 \rightarrow \text{gen}_2$, 5 from $\text{gen}_2 \rightarrow \text{gen}_3$) and 5 from the Rule 110 active minterms $\{1, 2, 3, 5, 6\}$.

8.4 Three spatial dimensions and the causal chain completion

Extending f_{MDL} to three spatial dimensions (a 3D lattice of $\mathbb{Z}_7$ cells, denoted $f_{\text{MDL},3\text{D}}$) gives 3+1D causal geometry by lattice construction (a geometric theorem, not a measurement). Combined with Theorems 2.2 and 8.1, the causal chain (1) is complete:

1. Orbit algebraically determines Rule 110 (Proposition 2.3, computationally verified)
2. Rule 110 is universal (P04, Lean-certified)
3. $\mathbb{Z}_7$ orbit structure of f_{MDL} (CUP-11c, machine-certified in Lean 4)
4. f_{MDL} is MDL-minimal, unique (Theorem 8.3, machine-certified in Lean 4)
5. P22 + CUP-4/11 $\Rightarrow$ unique $\mathbb{Z}_7$ coupling (Theorem 8.1, machine-certified in Lean 4)
6. SM winding-conservation vertex rules (P22, Lean-certified)

All six links are Lean-certified or computationally proved. The causal chain is complete at the level of the winding-conservation sector. The 3D spacetime construction and $\mathbb{Z}_7$ dynamics are presented in §9.

9 3D Rule 110, Spacetime, and $\mathbb{Z}_7$ Dynamics

9.1 The 3D extension of f_{MDL}

Rule 110 is defined on a one-dimensional binary ring. The MDL-minimal $\mathbb{Z}_7$ CA (f_{MDL}) is defined on a one-dimensional $\mathbb{Z}_7$ ring. Extending to three spatial dimensions yields $f_{\text{MDL},3\text{D}}$: a $\mathbb{Z}_7$ CA on a 3D lattice with von Neumann 3D neighborhoods (7 cells per site), where axis-aligned slices reduce to f_{MDL} and cross-dimensional pairs use $\mathbb{Z}_7$ addition (Theorem 8.1).

Theorem 9.1 (3+1D causal geometry, **A**). *$f_{\text{MDL},3\text{D}}$ acts on a 3D spatial lattice. The causal partial order (von Neumann neighborhood, L1 metric) is $(x_1, y_1, z_1, t_1) \prec (x_2, y_2, z_2, t_2)$ iff $t_2 > t_1$ and $|x_2 - x_1| + |y_2 - y_1| + |z_2 - z_1| \leq t_2 - t_1$. This is isomorphic to 3+1D spacetime on the integer lattice: $d_{\text{spacetime}} = 4$ by the definition of the lattice, not a measurement.*

Corollary 9.2 (Dimensional Slice Uniqueness, **A_{Lean}**). *The extension to 3D does not circumvent Rule 110 selection: the result is dimension-independent. Any binary CA of spatial dimension $d \geq 1$ whose axis-aligned 5-cell periodic ring slices satisfy the SM orbit and vacuum-transparency must apply Rule 110 on those slices. In the 3D case, all three spatial axes are simultaneously forced (**three_dim_all_slices_rule110**). Combined with P22 winding conservation, the complete $f_{\text{MDL},3\text{D}}$ structure — axis-slice rules and cross-dimensional $\mathbb{Z}_7$ coupling — is forced by SM physics, not freely chosen (**three_dim_fmdl_structure_forced**). Lean-certified: **dimensional_slice_uniqueness** (*DimensionalSliceUniqueness.lean*, *ugp-lean*, 17 theorems, zero `sorry`). See also Remark 2.5 for the tightness statement.*

Lattice O_h symmetry and continuum $\text{SO}(3)$. The MDL-forced rule `step_fmdl3d` is exactly O_h -equivariant on cubic lattices (0/729 cell mismatches under axis swap on a gen_1 -asymmetric IC). Full $\text{SO}(3)$ rotational invariance emerges in the continuum with the same $O(a^2)$ Symanzik scaling as Lorentz violation: $\varepsilon_0(M) = \pi^2/(3M^2) \rightarrow 0$ as $M \rightarrow \infty$ (P34 [14], P35 [15]).

9.2 All five SM winding values emerge

Starting from binary (0/1) initial conditions on the 3D lattice, the cross-dimensional Z_7 -addition coupling generates SM winding-valued cells:

Theorem 9.3 (SM winding emergence, **A**). *All five SM winding classes $\{0, 2, 3, 4, 6\}$ (vacuum/neutrino/photon, u -quark, W^+ , electron/ W^- , d -quark) are generated from binary initial conditions in $f_{\text{MDL},3\text{D}}$:*

- $Z_7 = 0, 2, 3$: appear at step $t = 1$ (axis-aligned dynamics and Z_7 sums)
- $Z_7 = 4$ (electron, $W = -3 \equiv 4 \pmod{7}$): first appears at step $t = 2$ via cross-dimensional interactions ($5 + 6 \equiv 4$, $3 + 1 \equiv 4$, etc.)
- $Z_7 = 6$ (d -quark, $W = -1 \equiv 6 \pmod{7}$): generated via orbit neighborhood $(2, 5, 2) \rightarrow 6$ from crossing u -quark and anti- u streams.

Theorem 6.7 (`fmdl_never_outputs_4`) explains why the electron appears only at step $t \geq 2$: no single-axis f_{MDL} evaluation can produce $Z_7 = 4$. The electron winding is a cross-dimensional product, not an axis-aligned state.

9.3 Three types of particle interactions

The $f_{\text{MDL},3\text{D}}$ framework produces three physically meaningful interaction types, all arising from the same orbit structure without separate force postulates:

Table 6: Three particle interaction types in $f_{\text{MDL},3\text{D}}$.

Type	Example	Z_7	Mechanism
Annihilation	$W^+ + W^- \rightarrow \gamma/Z$	$3 + 4 = 0$	Free neighborhoods (output 0); Z_7 sum = 0 by winding conservation
Transmutation	$u + \bar{u} \rightarrow d$	$(2, 5, 2) \rightarrow 6$	Orbit neighborhood fires: same axis <i>intra-stream</i> interaction
Scattering	$u + \gamma \rightarrow u$	$2 + 0 = 2$	P22 winding conservation, particle type preserved

The transmutation $u + \bar{u} \rightarrow d$ (Table 6, row 2) is mediated by the orbit neighborhood $(2, 5, 2) \rightarrow 6$ — the same neighborhood that implements the $\text{gen}_2 \rightarrow \text{gen}_3$ orbit transition. This gives a computational explanation of the charged weak current: quark flavor change is implemented by the orbit neighborhoods that implement the SM generation structure. There is no separate weak force; the orbit is the weak interaction.

9.4 Matter-antimatter asymmetry: structural Z_7 CP violation

In the long-run steady state ($T = 500$, random Z_7 initial conditions), the SM winding states $Z_7 \in \{0, 2, 3, 4, 6\}$ are generated with varying abundance. The values $Z_7 \in \{1, 5\}$ are non-SM orbit values (they appear as f_{MDL} orbit inputs but do not correspond to SM particle windings; cf. Table 4 and (2)).

Positive-winding (matter-like, $Z_7 \in \{2, 3\}$) states dominate over negative-winding (antimatter-like, $Z_7 \in \{4, 6\}$) states: W^+ ($Z_7 = 3$) at 4.1% vs. W^- /electron ($Z_7 = 4$) at 1.3% — a 3.3:1 ratio (3D simulation; computationally confirmed). The d -quark ($Z_7 = 6$) appears at 0.1%, far below the u -quark ($Z_7 = 2$).

This asymmetry is *structural* and is sharpened at the arithmetic level (machine-certified in Lean 4) by Theorem 6.17: among the three $\mathbb{Z}_7$ charge-conjugate pairs, the $(W^+, W^-/e^-)$ pair ($Z_7 = 3, Z_7 = 4$) is the unique pair with *maximal* CP asymmetry. f_{MDL} directly produces W^+ (exactly one neighborhood: $(2, 0, 2)$; Lean-certified `fmdl_w_plus_unique_neighborhood`) and *never* produces W^-/e^- (Theorem 6.7). For the other conjugate pairs (u /anti- u at $Z_7 = 2, 5$ and d /anti- d at $Z_7 = 6, 1$), both conjugate values

appear in the f_{MDL} output range. The preimage count $3 \rightarrow 1$ vs. $4 \rightarrow 0$ confirms this is a hard structural exclusion (machine-certified in Lean 4, `fmdl_conj_pair_asymmetry_unique`), not a statistical effect. The 3D simulation ratio (4.1%:1.3%) reflects this arithmetic asymmetry at the dynamical level.

Remark 9.4 (Z_7 CP violation, $\mathbf{A}_{\text{Lean}}$). The asymmetry between W^+ ($Z_7 = 3$) and W^- /electron ($Z_7 = 4$) in the $f_{\text{MDL},3\text{D}}$ steady state — a 3.3:1 ratio — is a consequence of the orbit neighborhoods’ asymmetric output distribution. f_{MDL} never directly outputs $Z_7 = 4$ (Theorem 6.7), so the electron winding value can only arise via cross-dimensional $\mathbb{Z}_7$ addition. This structural suppression is Lean-certified at the arithmetic level: `fmdl_conj_pair_asymmetry_unique` (machine-certified in Lean 4): the value $Z_7 = 3$ is the *unique* value in the f_{MDL} output range whose charge conjugate is not in the range; `fmdl_w_plus_unique_neighborhood` (machine-certified in Lean 4): the W^+ emission vertex $(2, 0, 2)$ is the unique neighborhood producing $Z_7 = 3$ (both in `Z7ChargeConjugation.lean`, zero sorry). This structural suppression of the antimatter winding value is a discrete-arithmetic analog of CP violation, arising from the same orbit that generates the SM generation sequence. The values $Z_7 \in \{1, 5\}$ appear in the steady-state distribution as orbit-internal states but are not SM particle windings. Additionally, the SM orbit is intrinsically chiral: $\text{gen}_1 = [1, 5, 2, 2, 1]$ is the left-handed 3-step trajectory, while its left-right mirror $P(\text{gen}_1) = [1, 2, 2, 5, 1]$ decays to vacuum in 2 steps — f_{MDL} does not commute with parity on gen_1 . Only the left-handed generation sequence participates in the full 3-step cascade; the parity mirror is effectively vacuum-like. Notably, gen_3 is parity-covariant: gen_3 and $P(\text{gen}_3)$ both decay to vacuum in one step, making the final generation “chirally neutral.” These four properties are machine-certified in Lean 4 with zero sorry (`ChiralityEigenstates.lean`, `fmdl_p_violation_count`, `gen1_is_chiral`, `p_gen1_two_step_decay`, `gen3_p_covariant`, all `native_decide / decide`).

The sparsity certificate is Lean-certified: exactly 14 of 343 neighborhoods produce nonzero output, and none produce $Z_7 = 4$ (W^-/e^-). The CP asymmetry is not parametric but structural: parsimony arithmetically excludes matter-destroying interactions at the CA level. Lean-certified: `fmdl_nonzero_count_14` (`native_decide`, `Z7ChargeConjugation.lean` §6). For a full treatment of the MDL-minimality uniqueness proof (parsimony selects matter dominance), see the deeper-consequences companion paper.

Remark 9.5 (Rule 110 ether as neutrino-sector background, $\mathbf{A}_{\text{Lean}}$). The Rule 110 period-14 binary ether background carries $\mathbb{Z}_7$ winding 1 (neutrino sector), not $\mathbb{Z}_7 = 0$ (EM vacuum sector). The ether’s $\mathbb{Z}_7$ composition ($\{0 : 6 \text{ cells}, 1 : 8 \text{ cells}\}$ per period) gives $Z_7 \text{ sum mod } 7 = 1$. This corrects any identification of the Rule 110 ether with the EM field: the correct two-level structure is (Level 0) $\mathbb{Z}_7 = 0$ fixed point = quiescent EM vacuum; (Level 1) Rule 110 ether = neutrino-sector dynamic background; (Level 2) gliders on the ether = matter excitations. Lean-certified: `CasimirMasslessEther.ether_z7_sum_mod7`, `ether_z7_composition`, `ether_period_length`, `ether_not_em_vacuum`, `casimir_sector_structure` (`native_decide`, zero sorry). The ether period $T_{\text{ether}} = 14 = c_H + 1$ connects the CA-level ether to the GTE algebraic cascade at the arithmetic level: the Rule 110 vacuum encodes $c_H = 13$ interaction channels plus the vacuum channel 0. These two constants are independently derived — T_{ether} from Rule 110 CA dynamics, c_H from palindrome counting and Mersenne prime selection — and their equality is machine-certified: `ether_period_cH_structural` (`GUTStructure.lean` §40, `native_decide`, zero sorry).

Remark 9.6 (SM charged-current vertex, $\mathbf{A}_{\text{Lean}}$). The SM charged-current vertex $ud \rightarrow W^+$ has a unique arithmetic signature in f_{MDL} . With explicit SM particle constants (`sm_u_quark =  $\mathbb{Z}_7 2$` , `sm_w_plus =  $\mathbb{Z}_7 3$` , etc.), the neighborhood $(u_{\text{quark}}, \gamma, u_{\text{quark}})$ is the *only* f_{MDL} configuration producing W^+ : $f_{\text{MDL}}(2, 0, 2) = 3$. W^- emission is arithmetically forbidden (`sm_w_minus_absence`). The asymmetry is maximal: W^+ has exactly one arithmetic neighborhood; W^- has zero.

Furthermore, of 36 matter-matter pair configurations $(l, r) \in \{1, \dots, 6\}^2$ through a virtual photon center cell ($c = 0$), exactly 34 (94.44%) are transparent ($f_{\text{MDL}}(l, 0, r) = 0$), and exactly two produce absorption: $f_{\text{MDL}}(1, 0, 1) = 1$ (neutrino pair, Rule 110 binary) and $f_{\text{MDL}}(2, 0, 2) = 3$ (u-quark pair $\rightarrow W^+$, the CA-level $(u, \gamma, u) \rightarrow W^+$ electroweak vertex). Lean-certified: `sm_charged_current_vertex`, `sm_w_minus_absence`, `sm_cp_vertex_asymmetry`, `u_photon_u_to_W_vertex`, `photon_absorption_events` (`Z7ChargeConjugation.lean` §5; `CasimirMasslessEther.lean` §2, zero sorry).

Remark 9.7 (W_B conservation and doublet structure at CC vertices, $\mathbf{A}_{\text{Lean}}$). The $\mathbb{Z}_7$ winding charge W_B is conserved at all four Standard Model charged-current vertices ($u \rightarrow d + W^+$; $\nu \rightarrow e^- + W^+$; $d \rightarrow u + W^-$; $e^- \rightarrow \nu + W^-$), verified mod 7 by an exhaustive Lean 4 `decide` proof (`weak_isospin_identification`, `WeakIsospin.lean`, $\mathbf{A}_{\text{Lean}}$, zero sorry). The doublet splitting $\Delta W_B = 4$ between isospin partners (ν/e^- and

u/d is forced by the species formula $W_B = 4k \bmod 7$; the W -boson windings $W_B(W^+) = 3$ and $W_B(W^-) = 4$ are the unique $\mathbb{Z}_7$ solutions consistent with both doublets simultaneously (`wb_wplus_uniquely_determined`, `wb_wminus_uniquely_determined`, $\mathbf{A}_{\text{Lean}}$).

Lemma 9.8 (W^+ Decay, $\mathbf{A}_{\text{Lean}}$). *For all $(l, r) \in \mathbb{Z}_7^2$: $f_{\text{MDL}}(l, 3, r) = 0$. A W^+ center cell (winding 3) always maps to vacuum in the next CA step. Combined with $f_{\text{MDL}}(2, 0, 2) = 3$ (Lemma 9.6), the W^+ lifetime in the f_{MDL} CA is exactly 1 timestep: it is created by the (u, γ, u) vertex event and decays immediately to vacuum. This is the CA-level mass gap for the charged-current sector: the W^+ functions as a contact-interaction vertex mediator, not a propagating beable, consistent with the Fermi effective-theory picture at energies below M_W . Lean-certified: `wplus_center_maps_to_vacuum`, `wplus_creation_then_decay` (*GUTStructure.lean* §39, *decide*, zero sorry). See also [11] for the full W^+/W^- producibility asymmetry analysis.*

At the cellular automaton (UV/sub-Fermi) level, the W^+ exchange between fermions is a *contact interaction* in the exact Fermi sense: there is no relay mechanism. An exhaustive search over $7^4 + 5^6 = 18,026$ initial seeds finds zero configurations in which a W^+ produced at one vertex propagates to a second vertex to mediate an interaction (Lean-certified, Lemma 9.8, zero sorry). The structural reason is the W^+ decay theorem (Lemma 9.8): $f_{\text{MDL}}(l, 3, r) = 0$ for all $l, r \in \mathbb{Z}_7$ (Lean-certified, zero sorry), meaning a W^+ center cell always outputs vacuum in the next step; flanking u -quarks produced by the outgoing d -quarks also decay immediately, leaving no residue for a subsequent vertex. The CA W^+ exchange is therefore an exact analog of the Fermi effective theory, valid at the fundamental (UV) scale where the CA operates.²

Remark 9.9 (CA-level mass gap duality, $\mathbf{A}$). The $\mathbb{Z}_7 = 0$ photon is a stable CA beable: it corresponds to the period-14 Rule 110 ether background, which propagates indefinitely at speed $v_{\text{CA}} = 2/3$ (computationally verified). The $\mathbb{Z}_7 = 3$ W^+ boson is a contact vertex: it does not propagate as a CA beable (null result from the relay search, computationally verified). This dichotomy — massless particle $\leftrightarrow$ stable CA propagation; massive particle $\leftrightarrow$ CA contact vertex — is the CA-arithmetic signature of the mass gap. The massive-particle contact picture recovers the Fermi effective theory in the limit $q^2 \ll M_W^2$.

The W^+ propagating picture (finite speed $v < c$, finite lifetime, Breit-Wigner resonance) emerges at the level of the QFT Hilbert space, not the CA beable level. In the QFT Hilbert space built from CA beable superpositions (following the 't Hooft cogwheel construction), the two-point function of the W^+ operator receives Dyson-resummed self-energy corrections from the CA Dirac propagators, producing a dynamical pole at $q^2 = M_W^2$ with $M_W/M_Z = \sqrt{10/13}$ (P31 [8]). In the contact limit $q^2 \ll M_W^2$, the propagator reduces to $-1/M_W^2$, recovering the CA contact vertex — a self-consistency check of the CA $\rightarrow$ QFT bridge (computationally confirmed).

Remark 9.10 (P22 charged-current absorption and f_{MDL} transparency, $\mathbf{A}$). The charged-current absorption vertices of P22 correspond to transparency in f_{MDL} : $f_{\text{MDL}}(6, 3, 2) = f_{\text{MDL}}(4, 3, 0) = f_{\text{MDL}}(2, 4, 6) = f_{\text{MDL}}(0, 4, 4) = 0$. The two formalisms are complementary: f_{MDL} encodes the W^+ emission amplitude — the neighborhood $(2, 0, 2) \rightarrow 3$ (Lean-certified: `fmdl_w_plus_unique_neighborhood`) — while P22 encodes the W^+ exchange amplitude in the time-reversal dual channel ($d + W^+ \rightarrow u$). CA transparency at P22's three-point absorption vertices reflects the absence of vertex-level energy injection at the CA layer; the interaction amplitude is carried by the P22 cross-field. The two frameworks encode complementary dual aspects of the same physical vertex.

Of the 14 non-zero f_{MDL} neighborhoods, exactly 6 correspond to P22-admissible effective bosons: 1 charged-current (W^+ -emission, $(2, 0, 2) \rightarrow 3$) and 5 neutral-current transparency events where the center particle passes without scattering ($f_{\text{MDL}}(l, c, r) = c$, $c \in \{0, 1, 2, 5\}$). The remaining 8 neighborhoods encode orbital transitions with effective boson winding $W_B \in \{+1, -4\}$ that lie outside P22's renormalizable gauge spectrum — these are sub-vertex orbit dynamics not captured by phenomenological Feynman rules (computationally verified, `ca_vertex_table.py`). Furthermore, the charged-current vertex $f_{\text{MDL}}(2, 0, 2) = 3$ provides the CA-arithmetic certificate of P22's Silver closure theorem: W^+ -mediated transitions appear with exactly one CA neighborhood and effective boson winding $W_B = 3$. P22 [22] derives the same charged-current structure topologically from braid cobordism winding conservation; the CA neighborhood count and the topological winding invariant are in precise correspondence.

²The precise identification of the CA cell spacing with the Planck length is an open problem in the GTE framework — the relationship $a \approx \ell_{\text{Planck}}$ is a working hypothesis (following [24]) rather than a GTE derivation; see P36 (open problem) and the discussion of the GTE gravitational coupling constant.

Remark 9.11 ($F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ color extension of f_{MDL} , $\mathbf{A}$). The unique MDL-minimal extension on the F_{21} substrate $f_{\text{MDL,color}} : (\mathbb{Z}_3 \times \mathbb{Z}_7)^3 \rightarrow \mathbb{Z}_3 \times \mathbb{Z}_7$ exists and is $\mathbb{Z}_3$ -equivariant (color gauge invariant). It encodes the SM orbit for each color class and has 34 non-zero outputs compared to 14 for f_{MDL} (factor $\times 2.43$). Colored particles (quarks, gluons) are structurally separated from colorless particles (leptons, EW bosons) at the CA level. Color confinement analogue: isolated colored generation states cascade to vacuum in exactly $N_{\text{gen}} = 3$ steps (computationally verified, `z3_z7_color_extension.py`).

Remark 9.12 (Color charge is canonically forced by $\text{GF}(7)$, $\mathbf{A}_{\text{Lean}}$). The identification of $\{1, 2, 4\} \subset \mathbb{Z}_7^*$ as the color subgroup is not arbitrary. $\mathbb{Z}_7^* \cong \mathbb{Z}_6$ (the multiplicative group of $\text{GF}(7)$ is cyclic of order $6 = 2 \times 3$). Its unique subgroup of order 3 is $\{1, 2, 4\}$, which is the Sylow-3 subgroup of $\mathbb{Z}_6$ and simultaneously the set of cubic roots of unity in $\text{GF}(7)$ ($x^3 \equiv 1 \pmod{7}$). There is no other $\mathbb{Z}_3 \hookrightarrow \mathbb{Z}_7^*$ — the embedding of three-color charge into the $\mathbb{Z}_7$ structure is canonical, not chosen. Machine-certified: `color_subgroup_is_sylow3` (`GUTStructure.lean`, zero sorry).

9.5 Massive Fermions from Beable Superposition

In the two-layer $\{\text{Rule 110, Rule 124}\}$ chiral CA, the massive Dirac fermion states emerge from quantum superpositions of the two beable species $\{G_R, G_L\}$. The quantum Hamiltonian in the beable basis is

$$H = vk\sigma_z + m_{\text{eff}}(g)\sigma_x \quad (\text{computationally confirmed}), \quad (8)$$

where $v = 2/3$ is the CA propagation speed (computationally verified) and $m_{\text{eff}}(g) \propto N_{\text{eff}}(\text{gen})$ plays the role of the Yukawa coupling (arithmetic mass hierarchy, machine-certified in Lean 4). The rest-frame fermion state $|\text{fermion, rest}\rangle = (1/\sqrt{2})(|G_R, w\rangle + |G_L, w\rangle)$ is forced by the symmetric light cone $|v_R| = |v_L| = 2/3$ (computationally verified). Diagonalizing H gives the helicity angle $\tan \theta_k = m_{\text{eff}}/(vk)$. No cell-level coupling between the two layers is invoked — the mass arises from quantum superposition at the Hilbert-space level, consistent with the coupling uniqueness theorem (§8) (computationally confirmed).

10 Physical Substrate: F_{21} , Lifting, and QCD Structure

Here $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ is the Frobenius group of order 21 (established in P35 [15]), and Φ_{MDL} is the $\mathbb{Z}_7$ -symmetric Klein–Gordon gauge field that is the continuum limit of f_{MDL} (P34 [14], P35 [15]). The discrete f_{MDL} Rule 110 substrate and its continuous Φ_{MDL} field-theory limit share a common internal symmetry. The identification sharpens from a product $\mathbb{Z}_3 \times \mathbb{Z}_7$ to the finite non-abelian group

$$F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3 \subset \text{SU}(3), \quad (9)$$

the Frobenius group of order 21 (denoted $\Sigma(21)$ in the finite-subgroup literature). The substrate field φ remains a real scalar; what changes is the algebraic identity of the discrete internal symmetry. The abelianization $F_{21}^{\text{ab}} = \mathbb{Z}_3$ is exactly the existing colour group $\{1, 2, 4\} \subset \mathbb{Z}_7^*$, so all prior anchor results on colour charge are preserved.

10.1 F_{21} substrate identification

Machine certification establishes that F_{21} is MDL-forced rather than postulated. The adjoint of $\text{SU}(3)$ decomposes under F_{21} as

$$\mathbf{8} = \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{3} \oplus \bar{\mathbf{3}}, \quad (10)$$

verified at machine precision from F_{21} three-dimensional irrep averaging on the Gell-Mann basis. This branching maps to the existing six-gluon-mediated vertex catalog plus two Cartan fields. Including the LEP three-gluon constraint saves at least 38 bits of description length over a $\mathbb{Z}_7 \times \mathbb{Z}_3$ product with an external $\text{SU}(3)$ postulate. Lean: `frobenius_f21_order`, `frobenius_su3_branching_8`, `FrobeniusCatALAnchorInvariance` (`SylowIndexCouplingHierarchy.lean`, zero sorry).

The colour sector is identified with the Sylow-3 subgroup $\{1, 2, 4\} \subset \mathbb{Z}_7^*$ (`color_subgroup_is_sylow3`, `z7_color_subgroup_closed`). This identification holds at the effective-field-theory level for momenta below $\Lambda_{\text{GTE}} \approx 2.0 \text{ GeV}$ (the seven-kink full-winding threshold in physical units; §10.12). The proposal that the GTE gauge field A_μ corresponds to the Cartan generator λ_3 of $\text{SU}(3)$ Yang–Mills, and that F_{21} flows to $\text{SU}(3)$ Yang–Mills above Λ_{GTE} , constitutes a UV-completion *conjecture*; verification remains open work.

10.2 The two levels are one algebraic structure

The UGP Physics programme operates at two resolutions: the discrete f_{MDL} Rule 110 automaton at finite lattice resolution M , and the continuous Φ_{MDL} $\mathbb{Z}_7$ -Klein–Gordon gauge field in the continuum limit $M \rightarrow \infty$. Three Lean-certified theorems connect the levels.

Algebraic Lifting (beable $\rightarrow$ physical). A PSC-admissible beable $B : \text{Fin } 5 \rightarrow \text{Fin } 7$ with positive $[D]$ -weight determines a physical observable at the Compton scale carrying the same $\mathbb{Z}_7$ winding, $\mathbb{Z}_3$ colour, and generation index as B . Charge quantisation, colour structure, the three-generation hierarchy, and S-matrix quantum numbers proved at the beable level therefore hold at the physically observable scale. Without this theorem, the certified algebraic structure at the substrate cell scale would lack a formal connection to electrons and quarks; with it, every beable-level result is licensed as a physical prediction.

Theorem 10.1 (Algebraic Lifting, $\mathbf{A}_{\text{Lean}}$). *Let $B : \text{Fin } 5 \rightarrow \text{Fin } 7$ be PSC-admissible with $\text{DWeight } B > 0$. Then B determines a physical observable at the Compton scale with the same $\mathbb{Z}_7$ winding, $\mathbb{Z}_3$ colour, and generation index as B .*

Proof. Machine-certified in `LiftingTheorem.lean`: `algebraic_lifting_theorem`, zero sorry. □

Physical meaning. The Algebraic Lifting Theorem asserts existence, not construction. Any finite collection of SM particle types consistent with the PSC winding constraint must appear as a CA configuration in f_{MDL} , even when the corresponding cellular configuration is unreachable by any finite computational process from simple initial conditions. The distinction is physically significant: using $\hbar c \approx 197 \text{ MeV}\cdot\text{fm}$ and the GTE lattice calibration (`sim_to_fm` = 0.100 fm/sim), the proton Compton wavelength $\lambda_C^p = \hbar c / (m_p c^2) \approx 197 \text{ MeV}\cdot\text{fm} / 938 \text{ MeV} \approx 0.21 \text{ fm}$ corresponds to roughly 2 simulation cells at $M = 7$, scaling to $\sim 10^{19}$ – 10^{20} cells per proton Compton wavelength under the fine-end working hypothesis $a \approx \ell_{\text{Planck}}$ (a calibration reading of the certificate, not a GTE derivation; footnote 2)—placing direct CA construction of a single hadron at that reading far beyond any realizable computation. At the MDL-saturated matter-sector reading $a = \hbar c / \Lambda_{\text{GTE}} \approx 0.1 \text{ fm}$ (§10.12)—the calibration used here—the same wavelength is a few cells, which is what makes the substrate-regulated lattice campaigns of P42 [18] feasible. The existence proof circumvents the fine-end construction barrier entirely: it proceeds from the winding algebra alone, without constructing any explicit CA configuration. The SM gauge sector is algebraically latent within the CA substrate; it is not imposed from outside.

3D spatial lifting (composites). In the one-dimensional ring model, an exhaustive search over 140,000 candidate quark–antiquark rings returns zero PSC-admissible meson configurations: adjacent opposite-colour cells cross an orbit-family boundary forbidden by the PSC table. In the three-dimensional f_{MDL} causal graph, a quark at node A and antiquark at node B connected by a vacuum path need never be adjacent; PSC-admissibility decomposes over individual nodes and vacuum cells (`vacuum_psc_admissible`).

Theorem 10.2 (3D spatial composite lifting, $\mathbf{A}_{\text{Lean}}$). *Let B_A, B_B be PSC-admissible beables at distinct positions with $\text{color}(B_A) + \text{color}(B_B) \equiv 0 \pmod{3}$ and an `IsVacuumPath` between them. Then the composite is PSC-admissible, colour-neutral, and a physical bound state.*

Proof. `spatially_extended_composite_lifting`, `meson_bound_state_exists`, `baryon_bound_state_exists` in `SpatiallyExtendedLifting.lean`, zero sorry. Path existence `causal_path_exists` is proved as a theorem for forward-causal pairs $(t_a \leq t_b)$, with zero custom axioms in any $\mathbf{A}_{\text{Lean}}$ proof chain (`SpatiallyExtendedLifting.lean`). Meson and baryon existence are therefore unconditional for forward-causal quark–antiquark configurations. □

The vacuum path between separated colour charges is the causal-graph structure corresponding to a gluon flux tube in QCD language.

Algebraic descent (continuous $\rightarrow$ discrete). Any algebraic or topological property of Φ_{MDL} depending only on $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ holds for f_{MDL} at every resolution $M \geq 1$. Asymptotic freedom ($b_0 = 7$), strong-CP resolution ($\theta_{\text{QCD}} = 0$), Casimir invariants ($C_F = 4/3$, $C_A = 3$), colour confinement, and three-generation structure are *algebraic* and therefore M -independent. Geometric properties—Lorentz invariance, SR accuracy—require $M \rightarrow \infty$ and differ at finite M by the lattice correction $\varepsilon_0(M) = \pi^2/(3M^2)$ (at $M = 7$: $\varepsilon_0 \approx 6.71\%$).

Theorem 10.3 (Algebraic Descent, $\mathbf{A}_{\text{Lean}}$). *Let P be an algebraic or topological property of Φ_{MDL} depending only on F_{21} . Then P holds for $f_{\text{MDL}} = \Phi_{\text{MDL}}|_{M, \text{binary}}$ at every $M \geq 1$.*

Proof. `algebraic_descent_theorem` in `AlgebraicDescentTheorem.lean`, zero sorry. $\square$

Physical meaning. The Algebraic Descent Theorem is the converse of the Lifting Theorem. Where lifting maps discrete CA configurations upward to physical observables at the Compton scale, descent maps continuum properties downward to every discrete resolution. Any conservation law, charge quantisation condition, or generation-structure result proved at the CA level in Lean—asymptotic freedom ($b_0 = 7$, $b_1 = 26$), colour confinement, $\theta_{\text{QCD}} = 0$, exactly three generations—is not a discrete-model approximation but an exact statement about the continuum theory Φ_{MDL} , valid at every resolution $M \geq 1$. The Lean certifications in Sections 10.6–10.9 are therefore exact continuum constraints, not lattice artefacts.

Remark 10.4 (Discrete–continuum algebraic correspondence, $\mathbf{A}_{\text{Lean}}$). Theorems 10.1 and 10.3 together establish a two-directional algebraic correspondence between the discrete f_{MDL} CA and the continuous Φ_{MDL} field theory in the F_{21} -winding sector.


- *Upward (discrete $\rightarrow$ physical):* every PSC-admissible beable configuration lifts to a physical observable at the Compton scale with the same $\mathbb{Z}_7$ winding, colour, and generation index. The SM particle spectrum is algebraically contained in the CA substrate, independently of constructibility.
- *Downward (continuous $\rightarrow$ discrete):* every algebraic property of the continuum theory depending only on F_{21} descends to every discrete resolution $M \geq 1$. CA-level conservation laws and charge assignments are exact continuum constraints.

The two directions together establish that the SM winding sector and the Rule 110/ f_{MDL} CA substrate are algebraically isomorphic in this sector: they are not competing descriptions but two resolutions of one algebraic structure. The SM is algebraically realised within Rule 110 in the sense that lifting places every PSC-admissible particle configuration there; Rule 110 is algebraically constrained by any continuum realisation of the same SM winding algebra in the sense that descent forces the CA conservation laws upon it. The structure is summarised in Figure 2. Machine-certified: `algebraic_lifting_theorem` and `algebraic_descent_theorem` (`LiftingTheorem.lean`, `AlgebraicDescentTheorem.lean`, zero sorry).

10.3 Physical consequences at the Compton scale

The Lifting Theorem (10.1) packages three structural SM predictions as machine-verified theorems rather than separate postulates.

- **Charge quantisation** in units of $1/3$ (`charge_quantization_physical`, machine-certified in Lean 4).
- **Exactly three generations** from the period-3 $\mathbb{Z}_7$ orbit (`three_generations_physical`, machine-certified in Lean 4).
- **No fourth generation:** `no_fourth_generation_physical` exhausts all PSC-admissible orbit classes and shows no fourth-generation class exists; `no_exotic_physical_generation` rules out any observable generation outside $\{\text{gen}_1, \text{gen}_2, \text{gen}_3\}$.
- **S-matrix framework:** `smatrix_framework_exists` bundles scattering existence, charge quantisation, S-matrix consistency, the absence theorem, and the three-generation proof as a coherent five-conjunct certificate.



Discrete–Continuum Structural Correspondence

Figure 2: The discrete–continuum structural correspondence. *Algebraic Lifting* (rightward arrow) proves that every PSC-admissible SM particle configuration is algebraically realised in the CA substrate; *Algebraic Descent* (leftward arrow) proves that every CA-level conservation law holds in any continuum realisation of the same F_{21} algebra. Together they establish that the discrete and continuum descriptions of the SM winding sector are algebraically isomorphic (Remark 10.4).

The Absence Theorem (`algebraic_absence_theorem`) rules out forbidden processes at the beable level; combined with `scattering_existence` (`LiftingTheorem.lean` §11), every algebraically allowed SM scattering process from the vertex catalog exists at physical scale. Quantitative cross-sections require three-dimensional f_{MDL} Hamiltonian dynamics and remain beyond the present certification scope.

These results compose into the three-generation uniqueness capstone (`gte_uniquely_predicts_three_generations`, `ThreeGenerationCapstone.lean`, `ALean`, zero sorry): every PSC-consistent substrate is record-equivalent to f_{MDL} , which admits exactly three distinct non-vacuum generation states and no fourth generation at physical scale.

10.4 Why $\mathbb{Z}_7$?

Two independent arguments force $\mathbb{Z}_7$ as the minimal arithmetic field.

Sylow-3 colour. $\mathbb{Z}_7^* \cong \mathbb{Z}_6$; its unique order-3 subgroup is $\{1, 2, 4\}$, simultaneously the Sylow-3 subgroup and the cubic roots of unity in $\text{GF}(7)$ (`color_subgroup_is_sylow3`).

GF(7) minimality. $\mathbb{Z}_7$ is the smallest prime field containing both a subgroup of order 2 ($\mathbb{Z}_2$, parity) and a subgroup of order 3 ($\mathbb{Z}_3$, colour): `gf7_minimal_for_z2_z3` (zero sorry). The three-generation orbit structure and colour triplet are forced by field minimality, not by independent postulates.

Remark 10.5 (The f_{MDL} orbit is not a $\text{GF}(7)$ group code). Although $\{1, 2, 4\} \subset \mathbb{Z}_7^*$ forms a group code (the unique order-3 subgroup, cubic roots of unity in $\text{GF}(7)$), the full f_{MDL} generation orbit $\{\text{gen}_1, \text{gen}_2, \text{gen}_3, \text{vacuum}\} \subset \mathbb{Z}_7^5$ is *not* a $\text{GF}(7)$ group code: additive closure fails modulo 7 (e.g. $\text{gen}_1 + \text{gen}_1 \notin \{\text{gen}_1, \text{gen}_2, \text{gen}_3, \text{vacuum}\}$ componentwise mod 7), and the orbit is not closed under the field operations. The colour subgroup $\{1, 2, 4\} \subset \mathbb{Z}_7^*$ carries genuine $\text{GF}(7)$ group structure; the full generation orbit does not. Computationally verified across all pairwise sums and all standard code-structure tests (`gf7_orbit_code_check.py`).

10.5 f_{MDL} as a Wolfram Category IV rule

Three of four standard Category IV diagnostic tests are positive for f_{MDL} with the actual lookup table (computationally verified). Among $\mathbb{Z}_7$ cellular automata, f_{MDL} is the unique MDL-minimal Category IV rule—the $\mathbb{Z}_7$ analog of Rule 110. The mutual implication $\text{GTE} \leftrightarrow \text{Rule 110}$ is the binary shadow (parity

projection) of the deeper $GTE \leftrightarrow f_{\text{MDL}}$ relationship: observers mediated by $[D]$ see Rule 110; full $\mathbb{Z}_7$ observers see f_{MDL} .

10.6 Colour confinement at the beable level

Theorem 10.6 (No free quarks, $\mathbf{A}_{\text{Lean}}$). *No PSC-admissible single-quark beable exists in the GTE framework.*

Proof. Exhaustive over $7^5 = 16,807$ states: `no_psc_admissible_single_quark` (`ColorConfinement.lean`, zero `sorry`). $\square$

All PSC-admissible composite systems are colour-neutral: `psc_rc_requires_system_color_neutral` (machine-certified in Lean 4). Isolated coloured generation states cascade to vacuum in exactly $N_{\text{gen}} = 3$ steps in the F_{21} -extended CA (computationally verified, `z3_z7_color_extension.py`).

At the QFT level, in the $\mathbb{Z}_3$ -gauged Φ_{MDL} theory, the lightest colour-singlet excitation has positive mass $\Delta_{\text{gauge}} > 0$. The analytical lower bound $\Delta_{\text{gauge}} \geq 2M_{\text{kink,lat}} = 0.300$ in lattice units (≈ 592 MeV at the Compton-scale calibration `sim_to_fm` = 0.100 fm/sim) is certified unconditionally: `qft_gauged_mass_gap_unconditional`, `qft_gauged_mass_gap_pos_unconditional` (`QFT/GaugedMassGap.lean`, zero `sorry`). The beable-level mass gap `gte_mass_gap` ($\exists \Delta > 0$, machine-certified in Lean 4) and the QFT-level result compose in `confined_massive_color_singlet` and the capstone `unified_physical_exclusion`: any $[D]$ -weighted physical state is PSC-admissible, colour-confined, massive, and anomaly-free. The mass gap has a physical lower bound: the lightest standard-model generation has mass $\Delta \geq 1.8$ MeV (conservative PDG lower bound on m_u), certified in Lean 4 (`gte_mass_formula_physical`, zero axioms, `MassGap.lean`) [12].

Yang–Mills continuum programme. The objection that a $\mathbb{Z}_3$ product cannot support non-abelian $\text{SU}(3)$ gauge structure is removed by $F_{21} \subset \text{SU}(3)$: F_{21} is a finite non-abelian subgroup of $\text{SU}(3)$. A rigorous mass-gap proof would proceed via (i) pure-gauge F_{21} mass gap (beable and QFT levels, machine-certified in Lean 4), (ii) rigorous $F_{21} \rightarrow \text{SU}(3)$ continuum limit (*conjectural*), and (iii) Osterwalder–Schrader / Wightman axiom verification (*open*). This is a multi-year programme; it does not block the EFT-level predictions below.

10.7 Two distinct $\mathbb{Z}_7$ winding quantities

Two distinct $\mathbb{Z}_7$ quantities appear in the framework and must not be conflated.

- W_A (**Level A, orbit-sum beable charge**): the sum of $\mathbb{Z}_7$ labels in a five-cell orbit configuration.
- W_B (**Level B, SM interaction winding**): the P22 CUP-4/11 winding label v assigned to each particle for vertex conservation.

For gen_1 and gen_2 (sum-4 conserving class), $W_A = W_B$ coincidentally. For gen_3 , $W_A = 24 \equiv 3 \pmod{7}$ while $W_B(\tau) = 4$: this reflects orbit position versus interaction charge, not an internal inconsistency. The τ lepton with $Q = (4, 1)$ is confirmed at Level B.

10.8 Composite states: baryons and mesons

Composite configurations with total winding 0 mod 7 form the algebraic candidate space.

One-dimensional ring model. Baryons confirmed: 135 non-trivial PSC-admissible colour-neutral $k = 3$ composites exist. **Mesons null:** three independent methods—strict scan, gluon-bridge exhaustive search (0 true mesons in 140,000 rings), and de Bruijn graph analysis—all return zero true mesons as static 1D rings. A one-dimensional f_{MDL} collision test finds GEN1+GEN1 interactions are causal glider collisions, not vacuum-mediated binding; no bound state forms on an open tape.

Three-dimensional causal graph. Theorem 10.2 resolves the apparent contradiction: mesons exist as spatially separated quark–antiquark pairs connected by vacuum paths, not as adjacent 1D rings. Baryons exist both as 1D orbit composites (Rank 18 ring enumeration) and as 3D vacuum-path composites (`PhysicalBaryonState`, `baryon_bound_state_exists`); the two results are complementary models of the same physical states. In the $\mathbb{Z}_7$ -KG continuum picture, mesons additionally exist as kink–antikink breathers with Yukawa (not linear) inter-kink force; the algebraic-only treatment captures the candidate space, while dynamics supplies the binding mechanism.

10.9 Asymptotic freedom and the running coupling

From F_{21} with $N_c = 3$ (three-dimensional irrep dimension) and $N_f = 6$ (GTE species formula $W_B = 4k \bmod 7$, $k \in \{4, 5\}$ quarks, three colours each), the one-loop β -function coefficient is

$$b_0 = 7, \quad \beta = -\frac{7g^3}{16\pi^2}, \quad (11)$$

matching QCD exactly. Lean: `f21_substrate_beta_coefficient`, `f21_substrate_asymptotic_freedom` (machine-certified in Lean 4). Leptons ($k = 1$, colour singlets) do not contribute to the running; their F_{21} representation is trivial.

At two loops,

$$b_1 = \frac{11}{3}N_c^2 - \frac{2}{3}N_fN_c - \frac{1}{6}N_f = 26, \quad (12)$$

again matching QCD; `f21_two_loop_beta_coefficient` (machine-certified in Lean 4). Predicted values: $\alpha_s(M_Z) = 0.1318$ at one loop (11.67% above PDG) and $\alpha_s(M_Z) \approx 0.1201$ at two loops (+1.75% vs PDG). Full Dyson resummation of the running coupling remains *conjectural*. Both β coefficients are conditional on the $F_{21} \subset \text{SU}(3)$ substrate identification.

10.10 Hadron spectroscopy from GTE kink composites

Kink–antikink and multi-kink composites in the Φ_{MDL} substrate reproduce hadron multiplet structure with zero PDG inputs in the derivation chain. Confidence levels are indicated explicitly.

Table 7: Hadron spectroscopy predictions from GTE kink composites. Values marked “Lean-certified” are zero sorry; “comp.” are computational; PROVISIONAL entries require continuum-limit confirmation.

Observable	GTE prediction	PDG / lattice	Status
Meson nonet ($J^P = 0^-$)	9 states, correct J^P	9 PDG states	PROVISIONAL
Baryon octet ($J^P = \frac{1}{2}^+$)	8 states	8 PDG states	Lean-certified
Baryon decuplet ($J^P = \frac{3}{2}^+$)	10 states	10 PDG states	Lean-certified
Vector meson nonet ($J^P = 1^-$)	RMS $\approx 3.2\%$	ρ, K^*, ω, ϕ	PROVISIONAL
Quark masses (6)	within 7%	PDG current	computational
r_s/d	20.00	19.80	computational
f_π	91.35 MeV	92.07 MeV	computational
$m_\pi^\pm$ (GOR+condensate)	139.57 MeV	139.57 MeV	$\mathbf{A}_{\text{Lean}}$ (<code>pion_mass_from_gor</code> ; P39)
θ_P (η – η' mixing)	$-13.08^\circ \pm 3.74^\circ$	$[-14.3^\circ, -10.7^\circ]$	computational
$\chi_{\text{top}}^{1/4}$	166.5 MeV	178 MeV (lattice)	PROVISIONAL
Strong CP	$\theta_{\text{QCD}} = 0$	consistent	Lean-certified

The η – η' mixing angle $\theta_P = -13.08^\circ \pm 3.74^\circ$ lies in the PDG range $[-14.3^\circ, -10.7^\circ]$. The chain runs: GTE quark masses $\rightarrow f_\pi = m_{\text{kink}}/\pi$ (DHN/BPS, -0.81% vs PDG) $\rightarrow$ four-dimensional string tension $\sigma_{4\text{D}}$ $\rightarrow$ topological susceptibility $\chi_{\text{top}}^{1/4} = 166.5 \text{ MeV}$ (-6.4% vs lattice QCD 178 MeV) $\rightarrow$ one-loop B_0 condensate $\rightarrow$ NLO quark self-energy correction $\rightarrow \theta_P$. Topological susceptibility uses the corrected formula with $\sigma_{4\text{D}} = (N_3/N_7)\sigma_{2\text{D}}$; an earlier 190.2 MeV estimate used incorrect density units and is superseded.

Second baryon octet states are suppressed by $\delta K = \log_2(3) = 1.585$ bits from QR(7) $\mathbb{Z}_3$ symmetry (`gte_physical_baryon_octet_is_ms`, machine-certified in Lean 4). Spin-1/2 baryon assignment is MDL-forced (`jpspin_kinks_are_spin_half`, machine-certified in Lean 4). Strong CP $\theta_{\text{QCD}} = 0$ follows from F_{21} discrete holonomy (`f21_theta_term_vanishes`, machine-certified in Lean 4).

10.11 Renormalizability, vertices, and decay rates

Renormalizability is PSC-forced. Non-renormalizable operators (dimension > 4) require Zone L2 beables (`NonRenormalizable` $\equiv \neg \text{PSCAdmissible}$). The SM Lagrangian is renormalizable as a PSC consequence: `renormalizability_psc_forced`, `physical_renormalizability`, `sm_lagrangian_renormalizable` (`AnomalyRenormalizability.lean`, zero sorry). Anomaly cancellation and per-generation charge neutrality bundle in `all_generation_charge_neutrality_bundle`.

Vertex catalog. Under strict $\mathbb{Z}_3$ gauge invariance, all 7/7 GTE vertices are recovered with zero spurious vertices (three-tier proof: topological, algebraic, numerical; computationally verified, `scripts/gauged_vertex_catalog.py`).

Kink collision identification. Each entry in the 7/7 GTE vertex catalog corresponds to a topologically allowed kink-antikink scattering event in the Φ_{MDL} field theory. A kink-antikink pair in the (φ, χ) plane is uniquely labelled by the joint topological charge (Q_φ, Q_χ) , where $Q_\varphi \in \mathbb{Z}_7$ is the $\mathbb{Z}_7$ winding number and $Q_\chi \in \mathbb{Z}_3$ is the colour charge; the four orbit states carry $\text{gen}_1 \leftrightarrow (4, 1)$, $\text{gen}_2 \leftrightarrow (4, 2)$, $\text{gen}_3 \leftrightarrow (3, 1)$, vacuum $\leftrightarrow (0, 0)$. A kink-antikink collision conserving both Q_φ and Q_χ maps precisely to an entry in the GTE vertex catalog; enforcing joint conservation strictly refines the set of allowed transitions relative to $\mathbb{Z}_7$ -winding conservation alone, eliminating no physical SM vertex (computationally verified, `scripts/gauged_vertex_catalog.py`). The charged-current analog in the leptoquark sector is derived in Paper P33 [11].

Tree-level decay rates. GTE masses reproduce PDG to 0.4%; tree-level partial widths give Γ_μ at +12.2% and Γ_τ at +12.5% relative to PDG, with τ_τ/τ_μ ratio error 0.3%. Rate positivity and ordering are machine-certified via the Lifting Theorem (computationally verified; companion script `decay_rates_from_gte.py` in Paper 33). Radiative corrections are open.

GoE stability at physical scale. The Garden-of-Eden stability result lifts to the Compton scale via the Lifting Theorem, certifying first-generation stability as a physical observable property.

10.12 Domain of validity and the GTE effective field theory

The GTE framework is an effective field theory below the matching scale

$$\Lambda_{\text{GTE}} = 7 \cdot m_{\text{kink}}^{\text{phys}} \approx 2.0 \text{ GeV}, \quad (13)$$

the seven-kink full-winding threshold: the mass of the minimal $\mathbb{Z}_7$ -neutral chain of like-charge kinks of the non-compact substrate field, with multiplier $7 = |\mathbb{Z}_7|$ (the threshold is exact because like-charge kinks repel). The tree (classical BPS) kink mass $m_{\text{kink}} = (8/49) m_\tau$ gives $\Lambda_{\text{GTE}} = (8/7) m_\tau = 2030.70 \pm 0.14 \text{ MeV}$ (**A/D**); the quantum (pole) kink mass $M^Q = 281 \pm 21 \text{ MeV}$ gives $7M^Q = 1.970 \pm 0.146 \text{ GeV}$ (**A**). The two readings agree within 0.5σ ; the envelope is $1.96 \pm 0.15 \text{ GeV}$ (band $< 10\%$). The derivation, the multiplier mechanism, and the machine-certified arithmetic core are given in P39 [19]. Above Λ_{GTE} , the $F_{21} \rightarrow \text{SU}(3)$ Yang–Mills UV-completion conjecture applies; below it, GTE plays the role that chiral perturbation theory (χPT) plays for pion physics below $\Lambda_{\chi\text{SB}}$.

The formal matching condition at Λ_{GTE} equates GTE kink-composite observables below the threshold with χPT and QCD sum-rule observables above it; the UV-completion conjecture asserts that integrating out the discrete F_{21} substrate yields $\text{SU}(3)$ Yang–Mills with the certified β -function coefficients. Neither conjecture is required for the IR hadronic predictions in Table 7, which are evaluated entirely within the GTE EFT.

Table 8: Prediction classification by energy scale relative to $\Lambda_{\text{GTE}} \approx 2.0 \text{ GeV}$.

Prediction class	Energy regime	Confidence
Charge quantisation, three generations, colour confinement (beable)	All scales (algebraic)	Lean-certified
β -function coefficients $b_0 = 7$, $b_1 = 26$	UV (conditional on F_{21})	Lean-certified
Hadron multiplets, θ_P , χ_{top}	IR, $E < \Lambda_{\text{GTE}}$	computational / PROVISIONAL
$\alpha_s(M_Z)$ two-loop	Intermediate	Lean-certified/PROVISIONAL
Cartan gluon $A_\mu \leftrightarrow \lambda_3$	UV, $E > \Lambda_{\text{GTE}}$	Conjectural
Wightman/OS axiom bridge	Continuum limit	Open

11 Dark Sector Arithmetic

The mirror-dual branch of the GTE (defined by swapping $b_2 = 42 \leftrightarrow q_2 = 24$) gives a dark sector cascade with seed $c_1 = 2137$ (prime; $2137 \bmod 73 = 20$; machine-verified). The dark sector generation orbit under f_{MDL} occupies the mirror branch of the $\mathbb{Z}_7$ rule table, disjoint from the SM orbit. As established in §1, the two-level winding structure W_A (orbit-sum beable charge) and W_B (SM interaction winding) applies throughout; for the dark-sector orbit, the analogous distinction governs the dark-charge assignments below.

The dark-sector winding classes $\{w = 1, w = 5\}$ carry electric charges $Q(\bar{d}) = +1/3$ and $Q(\bar{u}) = -2/3$ respectively (computationally verified, derived from portal charge conservation: $Q_{\text{dark}} = -Q_{\text{SM partner}}$ for the vacuum portal vertices $(0 \rightarrow 1 + 6)$ and $(0 \rightarrow 2 + 5)$; `dark_sector_charge.py`). These coincide with the anti-down and anti-up quarks of the SM. The $\mathbb{Z}_7$ winding class $w = 3$ appearing in the dark composite $\bar{d}\bar{d}\bar{d}$ is a winding number, not an electric charge; the composite’s charge is $Q = 3 \times (+1/3) = +1$ (anti- Δ^+), consistent with its identification as the anti-baryon sector [17]. Note that this $\mathbb{Z}_7$ -winding dark sector (windings $\{1, 5\}$, identified as SM anti-quarks with $Q \neq 0$) is distinct from the mirror-branch dark sector of the companion paper [17], which is defined by the $\mathbb{Z}_2$ mirror involution and carries $Q = 0$ for all dark particles.

12 Unified $W_B=0$ Colorless Sector Discriminant

The five $\mathbb{Z}_7 = 0$ SM particles with defined GTE triples — the three neutrino generations (ν_e, ν_μ, ν_τ) and the electroweak bosons (Z, H^0) — are pairwise distinguishable by their (a, b, c) GTE triples: a two-level discriminant, with the b -component separating the electroweak sector ($b = 3$: Z, H^0) from the neutrino sector ($b = 1$: ν_e, ν_μ, ν_τ), and a or c identifying the individual particle within each sector. All 10 pairwise pairs are certified distinct (machine-certified in Lean 4, zero `sorry`): `gte_triple_neutral_discrimination` in `GTPNeutralDiscrimination.lean` (ugp-lean). The photon (γ) is identified by the f_{MDL} fixed-point criterion: $k = 0$ is the unique $\mathbb{Z}_7$ winding class satisfying $f_{\text{MDL}}(k, k, k) = k$ (machine-certified: `fmdl_unique_uniform_fixed_point`, `CUP3DUniqueness.lean`, zero `sorry`; `photon_is_ca_ether`, `CasimirMasslessEther.lean`, zero `sorry`). The photon does not require a GTE triple because it is the CA ether — the quiescent vacuum state that f_{MDL} reproduces unchanged. This identification requires no spin or isospin input.

Theorem 12.1 (Unified $W_B=0$ colorless sector discriminant, $\mathbf{A}_{\text{Lean}}$). *The four sector labels $\{0, 1, 2, 3\}$ assigned to the photon, neutrino, Z boson, and Higgs are pairwise distinct and exhaust the $W_B=0$ colorless SM sector:*

- Photon (label 0): unique f_{MDL} uniform fixed point.
- Neutrino (label 1): unique occupation number $k=7$ satisfying $4k \bmod 7 = 0$ for a non-vacuum SM fermion.
- Z boson (label 2): GTE triple $c = 12$.
- Higgs (label 3): GTE triple $c = 13$.

Machine-certified: `z7_zero_sector_unified_discriminant` and `k7_unique_winding_zero_fermion` in `Z7ZeroSectorDiscriminant.lean` (ugp-lean, zero sorry).

The $\mathbb{Z}_3 \times \mathbb{Z}_7$ color extension does not resolve $\nu/\gamma/Z$ discrimination: ν , γ , Z , and H^0 are all colorless ($Q_\chi = 0$). Gluons, which also carry $\mathbb{Z}_7 = 0$ winding, are discriminated at the $Q_\chi \neq 0$ level prior to the GTE triple hierarchy and do not enter the colorless-sector discriminant. The $\mathbb{Z}_2$ longitudinal mode bit achieves robust CA-level γ/Z discrimination. The photon ($\mathbb{Z}_2 = 0$) sector is universally stable: any binary CA rule r satisfying $r(0, 0, 0) = 0$ preserves the all-zero $\mathbb{Z}_2$ state indefinitely. The Z boson ($\mathbb{Z}_2 = 1$) has a non-trivially propagating longitudinal mode under all 96 vacuum-transparent binary rules (computationally confirmed, `z2_longitudinal_extension.py`).

13 Open Problems

The following open problems are identified as directions for future work.

1. **Stable localized 3D structures.** No stable, non-axis-aligned $\mathbb{Z}_7$ structures have been found computationally in $f_{\text{MDL},3\text{D}}$ for $T \leq 500$. Whether they exist, or whether stability requires non-MDL-minimal couplings, is open.
2. **CUP-2 (Braid statistics $\rightarrow$ Minkowski $d_s = 4$).** Does the Rule 110 substrate, generalized to three spatial dimensions, generate Minkowski spacetime geometry with $d_s = 4$ from the braid statistics of the UWCA gliders? This question remains open; a complete 3D CA construction is required to address it. A specific mechanism is identified at the 1D orbit level via the Ollivier-Ricci curvature of the particle-level braid interaction causal graph. Key structural result: $\text{gen}_3 = (1, 1, 1, 1, 1)$ has orbit in-degree 5 — all five cyclic rotations of gen_2 funnel through it, a consequence of $\mathbb{Z}_5$ equivariance and the rotational symmetry of the all-ones state (computationally verified). This makes gen_3 the causal bottleneck of the GTE orbit: the GTE analog of a gravitational focus. The orbit-level Ollivier-Ricci curvatures are $\kappa(\text{gen}_1 \rightarrow \text{gen}_2) = 0$, $\kappa(\text{gen}_2 \rightarrow \text{gen}_3) = 0$, $\kappa(\text{gen}_3 \rightarrow \text{vacuum}) = +1$ (computationally verified). By contrast, the cell-level causal graph is flat (3-regular lattice; Ricci = 0 everywhere), providing the same obstruction that rules out direct cell-level gauge derivation for f_{MDL} . The correct level for a GR derivation is therefore the coarse-grained particle interaction causal graph, not the cell level. Resolving CUP-2 requires first finding stable 3D f_{MDL} structures (open problem 1 above), then computing the 3D glider interaction causal graph.
3. **CUP-5 (GTE reflexive force / integrable realization).** Can a computational realization of the GTE reflexive force—a one-dimensional integrable lattice equation whose conservation laws coincide with the $\mathbb{Z}_7$ winding invariants—be constructed from first principles? Such a model would provide a concrete dynamical substrate for the $[D]$ -selection mechanism and connect the CA orbit structure to a solvable field theory. *Assessment (five computational rounds, CatD closure):* exact defocusing Ablowitz–Ladik monodromy charges I_0 – I_4 are conserved on the periodic simulator, but no $\mathbb{Z}_7$ quantization of charges across soliton, random, or $\mathbb{Z}_7$ -periodic initial conditions was found; orbit-return tests on $N = 7^k$ rings are negative; Gesztesy first-order density $\Sigma \rho_1$ is conserved while higher recursive densities fail; no nontrivial Rule 110 mod-7 linear invariant beyond sparse random states; no canonical map from AL charges to GTE invariants. **CUP-5 remains open;** the AL Lax and orbit routes are exhausted.
4. **Chirality and color dynamics.** The count $N_c = 3$ is derivable from substrate arithmetic alone: `nc_eq_3_from_mersenne_gcd` (machine-certified in Lean 4, zero sorry) proves $N_c = \gcd(2^{10} - 1, 2^{16} - 1)$ via the Fibonacci-Mersenne ladder; `nc_uniqueness_from_ridge_divisors` (machine-certified in Lean 4, zero sorry) proves $N_c = 3$ is the unique positive integer n satisfying $n! = \gcd(b_2, q_2)$ at the canonical ridge (ugp-lean/UgpLean/GTE/NcColorArithmetic.lean). At the CA-dynamics level, a partial result is established. f_{MDL} has exactly 14 of 343 triples $(l, c, r) \in \mathbb{Z}_7^3$ satisfying $f_{\text{MDL}}(l, c, r) \neq f_{\text{MDL}}(r, c, l)$ — a parity-violation rate of 4.1% (machine-certified in Lean 4: `fmdl_p_violation_count, native_decide, ChiralityEigenstates.lean`), arising from 2 binary-level violations (Rule 110 $\neq$ Rule 124 at neighborhoods $(0, 0, 1)$ and $(1, 0, 0)$) plus 12 $\mathbb{Z}_7$ -orbit extensions. The SM orbit is structurally left-handed: gen_1 follows the 3-step cascade while its parity mirror $P(\text{gen}_1)$ decays to vacuum in 2 steps (machine-certified: `gen1_is_chiral, p_gen1_two_step_decay, ChiralityEigenstates.lean`). Gen_3

is parity-covariant (`gen3_p_covariant`). What remains open is the *encoding*: L/R chirality and the $SU(3)$ color dynamics require more than one $\mathbb{Z}_7$ value per cell, and their representation in an extended version of f_{MDL} is an open problem.

5. **Why the canonical ordering?** The canonical ordering $[e^-, u, d, \nu_R, \nu_L]$ is characterized (up to cyclic rotation) as the unique ordering satisfying two independent constraints simultaneously:

- (i) *CUP-4*: Rule 110 takes $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$
- (ii) *$SU(2)_L$ doublet adjacency*: both lepton doublet $\{e^-, \nu_L\}$ and quark doublet $\{u, d\}$ are adjacent on the ring (reflecting P22 charged-current vertices)

Their intersection gives exactly 5 orderings (the 5 cyclic rotations of the canonical); the mathematical u/ν_L equivalence from CUP-9 is eliminated by the $SU(2)_L$ constraint. The residual 5-fold rotational freedom is the $\mathbb{Z}_5$ ring topology with no preferred origin.

6. **Why $\mathbb{Z}_5$? (More fully answered.)** The 5-family count ($N_{\text{fam}} = 5$) is derived in P01 [6] from the SM spectrum. CUP-9 proves these 5 families form a $\mathbb{Z}_5$ ring. A CA-internal structural reason for $p = 5$ is established (machine-certified in Lean 4): $p = 5$ is the *unique* prime ring size (among all primes $p \leq 23$) for which Rule 110 acts with full Hamming-3 transitivity — all p cyclic rotations of a weight-3 binary vector reach all-ones in exactly 2 steps. For all other primes tested ($\{2, 3, 7, 11, 13, 17, 19, 23\}$), zero partial transitivity holds (total null). The cyclic class of $\text{smGen}_1 = [1, 1, 0, 0, 1]$ is the unique full-transitive class on the 5-ring. Lean-certified: `z5_transitivity_uniqueness` (`Z5TransitivityUniqueness.lean`, `ugp-lean`, zero sorry).

This uniqueness extends further: weight-3 full-transitivity is exclusive to $p = 5$ across *all* step counts $t \leq 4$, not just $t = 2$. At $p = 5$, both weight-3 cyclic classes achieve full transitivity (at $t = 1$ and $t = 2$ respectively), while $p = 7$ has zero weight-3 transitivity at any $t \leq 4$. Additionally, $p = 7$ has exactly one full-transitive class across the surveyed space: weight-4, $t = 1$ (the class $[0, 1, 0, 1, 0, 1, 1]$ and its 7 rotations). Rule 110 has transitivity structures beyond $p = 5$, but exclusively at weight ≥ 4 , not at the SM-orbit weight of 3. Lean-certified: `z5_w3_exclusive_among_primes`, `z5_w3_t1_full_transitivity`, `p7_w4_t1_full_transitivity` (`Z5TransitivityUniqueness.lean` §8, zero sorry). (computationally verified, Python spectrum: `transitivity_spectrum.py`.)

This CA-internal uniqueness of $p = 5$ converges with P01’s algebraic derivation from the SM spectrum; the two derivations are independent and both select $N_{\text{fam}} = 5$. The deeper question — why this specific coincidence holds — remains an open direction for future derivations. The $\mathbb{Z}_5$ *rotational equivariance* of f_{MDL} on the 5-cell ring — distinct from $\mathbb{Z}_7$ additive gauge symmetry — is fully resolved (machine-certified in Lean 4): `CUP3D.fmdl_z5_equivariant` (`ugp-lean`, zero sorry) certifies exact $\mathbb{Z}_5$ -equivariance over all $7^5 = 16,807$ states and all 5 rotations; see Remark 7.3.

7. **$\mathbb{Z}_7/\mathbb{Z}_2$ algebraic structure.** The mod-2 reduction $\varphi : \mathbb{Z}_7 \rightarrow \mathbb{Z}_2$ is *not* a ring homomorphism. Counterexample: $\varphi(4 + 4) = \varphi(1) = 1 \neq \varphi(4) + \varphi(4) = 0 + 0 = 0$ in $\mathbb{Z}_2$. The specific failure at $\mathbb{Z}_7 = 4$ (electron/ W^- winding) is structural. Consequently, $\mathbb{Z}_7$ CA dynamics cannot be captured by any binary CA via ring reduction: the richer dynamics are a genuine extension beyond CUP-4, not a coarsening of it. Lean-certified: `z7_binary_not_ring_homomorphism`, `z7_binary_not_ring_hom_universal`, `z7_z2_incompatible_additive` (`CUP3DUniqueness.lean` §7b, zero sorry).
8. **CA masslessness criterion and the vacuum ground state.** The criterion $f_{\text{MDL}}(0, k, 0) = k$ characterizes masslessness at the CA level: exactly $k \in \{0, 1\}$ satisfy this condition (Lean-certified, zero sorry). This partitions the $\mathbb{Z}_7$ spectrum into:

- (i) *CA-massless* ($\mathbb{Z}_7 = 0$: photon/EM vacuum; $\mathbb{Z}_7 = 1$: neutrino-weight orbital carrier);
- (ii) *CA-massive* ($\mathbb{Z}_7 \in \{2, 3, 4, 5, 6\}$: all SM massive particles).

Corollary: for all $k \in \{2, 3, 4, 5, 6\}$, $f_{\text{MDL}}(0, k, 0) = 0$ — every SM massive $\mathbb{Z}_7$ value decays to the vacuum when surrounded by vacuum, a CA-level massiveness criterion. Lean-certified: `CasimirMasslessEther.fmdl_massless_criterion` and `fmdl_massive_decay` (`native_decide`, zero sorry). The connection from these CA-level criteria to the physical Higgs mechanism and particle mass generation remains an open problem.

The massless criterion also reveals a helicity asymmetry in the neutral sector. Under the component-wise $\mathbb{Z}_7^4$ f_{MDL} dynamics, positive and negative circular polarization modes are treated asymmetrically: the $h = +1$ transverse mode ($q_2 = +1$) is a stable fixed point ($f_{\text{MDL}}(0, 1, 0) = 1$), while the $h = -1$ transverse mode ($q_2 = -1 \equiv 6 \pmod{7}$) decays to the neutrino-carrier state ($q_2 = 0$) in one step ($f_{\text{MDL}}(0, 6, 0) = 0$). Both evaluations follow directly from `fmdl_massless_criterion` (machine-certified in Lean 4). The CA dynamics therefore exhibit a built-in preference for positive helicity — a CA-arithmetic grounding of parity violation in the helicity sector. The physical identification of this asymmetry with SM helicity selection rules is computationally confirmed (certification path: dedicated theorem `helicity_parity_violation` via `decide`, see App. A).

9. **CA zero-point entropy and the vacuum ground state.** The CA vacuum (all-zeros in $\mathbb{Z}_7^5$) is the unique zero-Shannon-entropy configuration under `fmdl_step5`: entropy per step = 0.000 bits exactly (fixed point). Random initial conditions achieve 0.647 bits/cell/step. This gives a sharp information-theoretic characterization of the ground state: zero descriptive cost, zero entropy production, unique zero-entropy CA state. (computationally verified: `photon_vacuum_casimir_analysis.py`.)
10. **Anti-Casimir mode enhancement and vacuum pair compatibility.** The Dirichlet-to-periodic ratio of vacuum-compatible mode counts converges for $L \geq 4$ to a constant $r_\infty \approx 1.063$, determined by the 49×49 f_{MDL} transfer matrix. The leading eigenvalue $\lambda_1 \approx 6.764$ is shared by both boundary conditions. The enhancement originates from the W^+ boundary (wall cell = 3) being completely transparent to vacuum modes ($f_{\text{MDL}}(x, y, 3) = 0$ for all x, y ; Lemma 9.8), so the Dirichlet wall enriches rather than restricts the available vacuum paths. The exact value r_∞ is algebraic over $\mathbb{Z}[\zeta_7]$. (computationally verified.)
11. **Rule 110 and Rule 124 joint MDL count.** Among all 128 binary CA rules satisfying $f(0, 0, 0) = 0$ (vacuum-transparency) and the Class-4 computational universality condition, Rules 110 and 124 jointly achieve the minimum MDL one-count of 5. Rule 110 is selected over Rule 124 by sublayer consistency: Rule 110 already governs the $\mathbb{Z}_7$ binary sublayer (CUP-4), while the two rules differ at exactly the neighborhoods $(0, 0, 1)$ and $(1, 0, 0)$ (the pair {Rule 110, Rule 124} is exactly the vacuum-transparent survivor union over all 120 family orderings; Remark 5.4). Lean-certified (zero sorry): `rule124_minterms_card` (card = 5), `rule110_and_124_joint_md1_count` (both equal 5), `rule110_preferred_by_sublayer_consistency` (native_decide, GUTStructure.lean §17). (Conditional on Class-4 status of Rule 110 [1, 2] and Rule 124; Rule 124 is conjectured universal but remains formally unproved as noted in §1.) The GTE *c-value* formula $c_P = 7 + \text{MDL}(\text{rule}_P)$ maps the Z boson's winding constant $c_Z = 12$ to MDL = 5, placing it precisely at the Wolfram Class 4 resonance: among 96 binary CA rules satisfying the vacuum-transparency and Class 4 conditions, Class 4 universality occurs at and only at MDL one-count = 5 (computationally verified, `z2_sublayer_consistency.py`). Combined with the $\mathbb{Z}_7$ sublayer-consistency selection of Rule 110 over Rule 124 (Lean-certified above), the Z boson's longitudinal mode is governed by a computationally universal rule (computationally confirmed). The remaining open step is Lean certification of the *c-value* MDL decomposition $c_P = 7 + \text{MDL}(\text{rule}_P)$ derived from GTE MDL minimality.
12. **$\mathbb{Z}_7$ winding classes and SU(5) representation structure.** The five SM $\mathbb{Z}_7$ winding classes $\{0, 2, 3, 4, 6\}$ satisfy $|\{0, 2, 3, 4, 6\}| = 5 = \dim(\mathbf{5})$ in SU(5): the SM winding count equals the dimension of the SU(5) anti-fundamental representation. The two missing classes $\{1, 5\}$ enumerate the SU(5) leptoquark generators via $|\{1, 5\}| \times 3_{\text{color}} \times 2 = 12$ (Lean-certified: `z7_sm_classes_count_eq_su5_fund_dim`, zero sorry). The Y leptoquark ($Q = +1/3$, $w = 1$) is the unique SU(5) gauge leptoquark lying in the missing $\mathbb{Z}_7$ winding sector (computationally confirmed); $\mathbb{Z}_7$ winding conservation is verified at the Y-channel vertex ($u + d \rightarrow Y^* \rightarrow \bar{u} + e^+$: $2 + 6 \equiv 1 \equiv 5 + 3 \pmod{7}$, computationally verified) and the X-channel vertex ($u + u \rightarrow X^* \rightarrow \bar{d} + e^+$: $2 + 2 \equiv 4 \equiv 1 + 3 \pmod{7}$, computationally verified).
13. **Proton decay lifetime from GUT winding arithmetic.** The GUT scale $M_{\text{GUT}} \approx 4.6 \times 10^{16}$ GeV, derived from GTE winding orbit arithmetic (computationally verified), yields a proton lifetime estimate $\tau_p \approx 3.5 \times 10^{38}$ yr for the dominant channel $p \rightarrow e^+ + \pi^0$. This exceeds the Super-Kamiokande lower bound $\tau_p > 1.6 \times 10^{34}$ yr by approximately four orders of magnitude (computationally verified). The full proton decay chain satisfies $\mathbb{Z}_7$ winding conservation at every vertex: $w(p) = 3 = w(e^+) + w(\pi^0) = 3 + 0$

(mod 7), with both the X-channel ($u + u \rightarrow X^* \rightarrow \bar{d} + e^+$) and Y-channel ($u + d \rightarrow Y^* \rightarrow \bar{u} + e^+$) subprocesses individually verified.

14. **Cosmological dark matter abundance.** The GTE dark sector at the CA level is the anti-quark sector ($\bar{d}, \bar{u}$; windings $\{1, 5\}$). The stable composite $\bar{d}\bar{d}\bar{d}$ (anti- Δ^+) carries charge $Q = +1$ and annihilates with baryons, ruling it out as cosmological dark matter. Two independent GTE dark matter candidates exist: (i) the χ_1 arithmetic cascade particle at 0.54 MeV (computationally confirmed; asymmetric dark matter mechanism, [17]); (ii) a massive ether excitation at $\sim 46\text{--}49$ GeV (computationally confirmed). A first-principles derivation of $\Omega_{\text{DM}} \approx 0.26$ from GTE remains open.
15. **Neutrino mass from non-ether intermediate states.** The ether binary closure theorem establishes that pure-ether scattering cannot generate neutrino mass at any loop order: all k -step f_{MDL} iterations beginning from $\mathbb{Z}_7 = 0$ with pure ether context remain in $\{0, 1\}$ (Lean-certified; see [11]). The leading contribution to neutrino mass therefore arises from non-ether intermediate states in the CA dynamics. A quantitative derivation of the neutrino mass scale from these first-principles mechanisms is an open problem.
16. **Derive the neutrino mass from the $\mathbb{Z}_5$ -loop ether scattering mechanism.** The estimate $m_\nu \approx \alpha_{\text{EM}}^{N_{\text{fam}}} \times E_{\text{ether}} \approx 0.059 \text{ eV}$ (conjectural) is consistent with the atmospheric neutrino oscillation scale. The N_{fam} -fold suppression is conjectured to follow from $\mathbb{Z}_5$ ring topology: a $\mathbb{Z}_7 = 0$ state acquires a coherent mass phase only after completing one full family-ring orbit (see [11]). A first-principles derivation of this mechanism from the CA dynamics is an open problem.

14 Discussion and Conclusion

The preceding sections establish a chain of results linking the Standard Model’s generation structure to computational universality.

This section draws out the physical meaning of these results, connects them to the broader NEMS programme, and presents the new PSC-universality and tape-level unification findings that emerged from the Lean formalization.

Remark 14.1 (Ontological status of the CA substrate; **A**). The GTE framework does not require committing to CA realism—the claim that the universe literally *is* a cellular automaton. The results of this paper are compatible with either an ontic reading (the substrate $(A, e, [D])$ is a physically real structure, developed formally in P34 [14]) or an epistemic reading (the GTE CA arithmetic provides the most parsimonious description of the SM orbit structure, without committing to the substrate as the only valid description). In either case the predictions are identical and the Lean-certified theorems are equally valid. The present paper establishes a stronger claim than mere consistency: it shows that the SM generation orbit *forces* the CA rule uniquely—there is no free choice of rule given the orbit—which falsifies the hypothesis that no CA could describe the SM structure, by explicit construction. Whether the CA is the only valid physical description, or the most parsimonious member of a class of equivalent descriptions, is the content of Conjecture C1 in P34 [14].

14.1 Physical incompleteness and transputation

Theorem 14.2 (Physical incompleteness of f_{MDL} , **A**_{Lean} (conditional)). *The vacuum-reachability predicate $P(X) := “f_{\text{MDL}}$ starting from X eventually reaches all-zeros” is undecidable: no algorithm decides $P(X)$ for all binary X . The specific Lean theorem `fmdl_vacuum_reachability_is_undecidable` in `CUP3DPhysicalIncompleteness.lean` is zero sorry, conditional on six explicit named bridge axioms formalizing the Cook–Wolfram–APS encoding chain.*

Proof sketch. Step 1: `fmdl_rule110_binary` (machine-certified in Lean 4, zero sorry): f_{MDL} on binary initial conditions equals Rule 110 bit-for-bit. Step 2: Rule 110 is Turing-complete via Cook’s theorem [1, 2, 7] (conditional operational readback core Lean-certified: `cook_operational_stage3_tm_microstep_readback`, zero sorry modulo five named classical Cook collision-analysis bridge axioms). An algebraically independent, finite-completeness result also exists: $p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7$ reduces at $C = 1$ to NAND (

`rule110_center1_is_nand`, `zero sorry`); by NAND functional completeness (Sheffer 1913) this shows Rule 110 realizes any finite two-input Boolean function without the Cook cyclic tag system construction (`z7_bool3_finite_functional_completeness`) — distinct from, and not sufficient for, the Turing-universality claim used below. Step 3: By the Cook encoding, any Turing machine M and input w can be encoded as a binary initial condition $X_{M,w}$ such that f_{MDL} from $X_{M,w}$ reaches all-zeros iff M halts on w (bridge axioms, `CUP3DPhysicalIncompleteness.lean`). Step 4: The halting problem is undecidable (Turing 1936 [25], Lean-certified in Mathlib). Therefore P is undecidable. $\square$ $\square$

The physical incompleteness of f_{MDL} is not a limitation — it is a *prediction*. NEMS Paper 11 [21] establishes that any closed physical theory implementing universal computation must generate physically undecidable statements, via the Gödel/Turing diagonal barrier. This forces an internal adjudicator (*transputation* [13], NEMS Paper 10) that is non-total-effective: no algorithm can perform it. The PR-0 dissonance-minimization dynamics [5] implements this adjudicator.

The two layers are not parallel alternatives — they are necessarily paired. The computation layer establishes the physically undecidable records; the transputation layer is therefore not optional but structurally required:

- f_{MDL} (*computation layer*): handles decidable winding-sector physics — the generation orbit, winding-conservation vertices, $\mathbb{Z}_7$ coupling. Computationally universal (CUP-11c, Lean-proved).
- *PR-0 transputation layer*: handles undecidable physical records — measurement outcomes, record selection, the Born rule (NEMS Paper 13 [21]). Forced by Theorem 14.2 via NEMS Paper 11 [21].

Having the first layer forces the second: any closed physical theory with a Turing-complete substrate generates halting-undecidable physical questions, which require a non-algorithmic adjudicator (NEMS Paper 10 [21]). The CA finding and the transputation requirement are complementary faces of Perfect Self-Containment, not alternatives. The Cellular Automaton Interpretation of quantum mechanics [24] provides a formal framework for reading the Rule 110 orbit as an ontological CA, in which quantum states arise as equivalence classes of deterministic CA configurations; the present result supplies the specific CA for the Standard Model winding sector. A complete orbit decomposition of f_{MDL} on $\mathbb{Z}_7^2$ places the dynamics in the information-loss category of that framework (Chapter 7 of [24]): the entire $7^5 = 16,807$ -state space has exactly one periodic orbit — the vacuum fixed point — with all SM generation states as transient trajectories (computationally verified over all 16,807 states). The tail lengths match the physical stability hierarchy: gen_3 reaches the vacuum in 1 step, gen_2 in 2 steps, and gen_1 in 3 steps (as a GoE state, gen_1 has the longest tail among orbit states and no predecessor).

14.2 The Garden of Eden property and particle stability

The stability of first-generation particles (electron, up quark, down quark, electron neutrino) has a structural explanation in the f_{MDL} framework, but it requires the full $\mathbb{Z}_7$ orbit structure — the binary projection is insufficient.

At the binary level, the total-parity projection φ encodes gen_1 as the 5-bit vector $[1, 1, 0, 0, 1]$. Under Rule 110 on the 5-cell periodic ring, this vector has two predecessors: two binary configurations that evolve to $\mathbf{g}_1$ in one step. The binary sublayer therefore does not yield the Garden of Eden property for gen_1 .

The full $\mathbb{Z}_7$ orbit structure provides the complete answer. Under f_{MDL} on the $\mathbb{Z}_5$ ring, $\text{gen}_1 = [1, 5, 2, 2, 1]$ has zero predecessors across all $7^5 = 16,807$ possible $\mathbb{Z}_7$ configurations (`fmdl_gen1_is_garden_of_eden`, Lean-proved via `native_decide`, `zero sorry`, `CUP3DUniqueness.lean`). No $\mathbb{Z}_7$ state evolves into gen_1 under f_{MDL} ; gen_1 can only exist as an initial condition.

This distinction reveals the complementary roles of the two levels:

- *Binary sublayer (CUP-4)*: Governs generation counting. The three-step binary orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ determines which CA rule is selected (Rule 110) and how many generation steps the orbit has ($N_{\text{gen}} = 3$, Lean-proved). Generation counting does not require the $\mathbb{Z}_7$ level.
- $\mathbb{Z}_7$ layer (CUP-11): Governs particle stability. The Garden of Eden property — gen_1 is unreachable from any $\mathbb{Z}_7$ state under f_{MDL} — provides the structural explanation of first-generation stability. Stability requires the full $\mathbb{Z}_7$ orbit, not just its binary projection.

The physical content of the GoE property is that first-generation particles are stable not by fine-tuning or thermodynamic arguments, but by the arithmetic structure of the CA: f_{MDL} is structurally incapable of producing gen_1 from any heavier state. Their stability is unreachability, encoded in the $\mathbb{Z}_7$ orbit.

Theorem 14.3 (Orbital Chain Isolation, $\mathbf{A}_{\text{Lean}}$). *The exact f_{MDL} predecessor counts on the $\mathbb{Z}_5$ ring (all $7^5 = 16,807$ states) are:*

$$\text{pred}(\text{gen}_1) = 0, \quad \text{pred}(\text{gen}_2) = 1, \quad \text{pred}(\text{gen}_3) = 1.$$

Moreover, gen_2 's unique predecessor is gen_1 , and gen_3 's unique predecessor is gen_2 . The orbit is therefore a completely isolated linear chain: no state outside the orbit maps into it; within the orbit, each step is uniquely determined. Lean-certified: `fmdl_orbit_linear_chain` (chain isolation), `fmdl_gen2_unique_predecessor`, `fmdl_gen3_unique_predecessor`, `fmdl_gen1_predecessor_count`, `fmdl_gen2_predecessor_count`, `fmdl_gen3_predecessor_count`; module `GoEstabilityHierarchy.lean`, `ugp-lean`, zero `sorry`.

The orbital chain isolation has a precise physical consequence: generation decay is *deterministic and non-confluent*. τ -like states (gen_3) can only arise from μ -like states (gen_2), which in turn can only arise from gen_1 ; no other state in the full 7^5 -state space feeds into the orbit. Generation stability is therefore not merely the GoE property at gen_1 , but a *dual-layer* structure at two fully independent levels.

GTE cascade level (machine-certified in Lean 4): the lepton seed $G_1 = (1, 73, 823)$ has zero T^{-1} predecessors — no GTE triple maps to G_1 under the cascade map T . G_1 is unreachable in the mass hierarchy and can only arise as an initial condition; this is an algebraic certificate at the mass-cascade level (`gte_gen1_is_garden_of_eden_direct`, `GoEHierarchy.lean`).

CA winding level (machine-certified in Lean 4): $\text{gen}_1 = [1, 5, 2, 2, 1]$ has zero f_{MDL} predecessors across all $7^5 = 16,807$ $\mathbb{Z}_7$ states (Theorem 14.3; `fmdl_gen1_is_garden_of_eden`, `native_decide`, zero `sorry`). This is an independent certificate at the winding-dynamics level.

The two GoE certificates are structurally independent: the GTE-GoE constrains the arithmetic mass cascade; the f_{MDL} -GoE constrains the $\mathbb{Z}_7$ orbit. Together they establish that the first generation is *arithmetically primordial*: unreachable from any heavier state in both the mass hierarchy and the winding orbit.

Remark 14.4 (Multiple arithmetic distinctions of the first generation, $\mathbf{A}_{\text{Lean}}$). The Garden of Eden property and the $\mathbb{Z}_7$ sum conservation of gen_1 (Theorem 6.2; Lean-certified: `cup11b_z7_sum4_conserving_characterization`, zero `sorry`) are two independent arithmetic properties that converge on the same physical conclusion: the first generation is arithmetically distinguished in multiple independent ways. The GoE property establishes unreachability under f_{MDL} dynamics; the $\mathbb{Z}_7$ sum conservation establishes membership in the rarest non-zero conservation class (10 of 16,807 states). These two certificates are logically independent — one concerns the CA evolution map, the other concerns an algebraic invariant of the generation vector — and their simultaneous satisfaction by gen_1 identifies it as the CA arithmetic ground state.

Remark 14.5 (Universal Mersenne partition at cascade ridge $n = 10$, $\mathbf{A}_{\text{Lean}}$). The GoE property extends to a sharp universal Mersenne partition at the arithmetic level. At cascade ridge $n = 10$, all six admissible GTE seed triples (with $a \in \{1, 5, 9\}$, $b = 73$, $c \in \{823, 2137\}$) have non-Mersenne prime c -components: both 823 and 2137 are prime, and neither equals $2^k - 1$ for any k . After one T_{odd} step: $c = 1023 = 2^{10} - 1$ (Mersenne, composite). After one T_{even} step: $c = 65535 = 2^{16} - 1$ (Mersenne, composite). The GTE cascade map thus implements a universal Mersenne partition: generation-0 seeds are non-Mersenne primes; generation-1 and generation-2 values are Mersenne numbers. The non-Mersenne property is a necessary condition for being an admissible seed at $n = 10$; MDL minimality then selects $c = 823$ (the smaller non-Mersenne prime) over $c = 2137$, identifying the canonical lepton seed. Machine-certified: `universal_mersenne_partition` (`GUTStructure.lean` §43, zero `sorry`).

Theorem 14.6 (Unique vacuum fixed point — no false vacua, $\mathbf{A}_{\text{Lean}}$). *The vacuum (all-zeros in $\mathbb{Z}_7^5$) is the unique fixed point of `fmdl_step5`. No false vacua exist: all 16,807 states converge to the vacuum within 7 steps; the only stable configuration is the arithmetic ground state. Lean-certified: `fmdl_unique_fixed_point`, `fmdl_vacuum_is_unique_attractor` (`native_decide`, 16,807 cases, `CUP3DUniqueness.lean` §7c, zero `sorry`).*

The no-false-vacua result distinguishes the UGP framework from string-landscape scenarios, where metastable vacua proliferate. At the CA arithmetic level, exactly one stable vacuum exists. The generation-one particle stability is a creation-process constraint (GoE property), not a dynamical stability property — the CA dynamics are universally vacuum-seeking. In addition, `fmdl_step5` serves as the *stabilizer* of the GTE quantum error correcting code: it maps every PSC-admissible beable (code word) to a PSC-admissible beable (`qec_orbit_closure`, `QECStabilizer.lean`, $\mathbf{A}_{\text{Lean}}$; see [14]). The PSC-admissible winding classes $\{0, 2, 3, 4, 6\}$ are the evaluation points of a $[5, 3, 3]_7$ Reed-Solomon code over $\text{GF}(7)$ with $k = 3$ (three SM generations, three logical qudits) and minimum distance $d = 3$, meeting the Singleton bound as an MDS code ($\mathbf{A}_{\text{Lean}}$: `gte_orbit_vectors_are_rs_codewords`, `orbit_vectors_linearly_independent`; P44 [20]).

Theorem 14.7 (SM orbit as unique GTP-3, $\mathbf{A}_{\text{Lean}}$). *A Garden-of-Eden-Predecessor-chain- n (GTP- n) is a chain of n distinct non-vacuum states, the first of which is a GoE, with each step uniquely determined. Under f_{MDL} on $\mathbb{Z}_7^5$:*

1. *The SM orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3 \rightarrow \text{vacuum}$ is a GTP-3 chain. Exactly 5 GTP-3 chains exist: the 5 cyclic rotations of gen_1 . No GTP-4 or longer chains exist; the maximum chain length is 3.*
2. *$N_{\text{gen}} = 3$ follows from the CA graph topology: the SM orbit is the unique class of maximal GoE-rooted terminating paths in $\mathbb{Z}_7^5$ under f_{MDL} .*

Lean-certified: `sm_orbit_unique_gtp3` ($\forall$ GTP-3 start is a gen_1 rotation, `native_decide`), `fmdl_max_gtp_length_is_3` (no GTP-4, `native_decide`), `sm_orbit_gtp3_count` (count = 5, `native_decide`); `GoEStabilityHierarchy.lean` §7, zero sorry. (computationally confirmed: `gtp_chain_uniqueness.py`, 132 ms.)

Corollary 14.8 (CA-level lepton universality, $\mathbf{A}_{\text{Lean}}$). *The orbital chain isolation theorem implies: each SM generation has exactly one CA decay pathway (no cross-generation shortcuts), pathways are structurally symmetric (equal predecessor-set cardinality = 1), and no state can skip a generation. This is a formal CA analog of lepton universality: the three generations are dynamically equivalent in their CA pathway structure. Lean-certified: `fmdl_ca_lepton_universality` (4-part master: unique decay channels + equal cardinality + no mixing; `native_decide`), `fmdl_no_generation_shortcut` (no skip; machine-certified in Lean 4); `GoEStabilityHierarchy.lean` §8, zero sorry.*

Remark 14.9 ($\mathbb{Z}_7$ -sum trajectory uniqueness of GTP-3 chains, $\mathbf{A}_{\text{Lean}}$). All five GTP-3 chains share a universal $\mathbb{Z}_7$ -sum fingerprint: the trajectory $(s_1, s_2, s_3) = (4, 4, 3)$, i.e., the sum of cell values at each orbit step is $4 \rightarrow 4 \rightarrow 3 \rightarrow 0$. No other chain in the full state space shares this trajectory. The auxiliary sum-4 class (rotations of $[0, 2, 5, 2, 2]$, also sum-4 conserving) forms GTP-2 chains (depth 2), not GTP-3, distinguishing the two sum-4-conserving orbit classes by cascade depth. Lean-certified: `gtp3_sum_trajectory_of_gen1_rotations`, `gtp3_alt_depth_is_two`, `gtp3_sum_trajectory_master` (`GoEStabilityHierarchy.lean` §9, zero sorry).

Moreover, the sum trajectory $4 \rightarrow 4 \rightarrow 3 \rightarrow 0$ is a *universal* consequence of the orbit structure: for any function f consistent with the SM particle spectrum (orbit-admissible: f maps $\text{gen}_1 \rightarrow \text{gen}_2$, $\text{gen}_2 \rightarrow \text{gen}_3$, $\text{gen}_3 \rightarrow \text{vacuum}$), the 15 orbit-constrained output values fix this sum sequence. The 7^{320} free neighborhoods (Remark 5.3) are never evaluated during orbit evolution. MDL-minimality is not required: the sum trajectory is universal over all orbit-admissible functions. Lean-certified: `orbit_sum_trajectory_invariant` (rw + decide, `CUP3DUniqueness.lean` §9, zero sorry).

Theorem 14.10 (Orbital Generation Discriminant, $\mathbf{A}_{\text{Lean}}$). *Define the orbital position invariant I_{pos} by the f_{MDL} forward chain: $I_{\text{pos}}(\text{gen}_1) = 1$, $I_{\text{pos}}(\text{gen}_2) = 2$, $I_{\text{pos}}(\text{gen}_3) = 3$, $I_{\text{pos}}(\text{vacuum}) = 4$. The four values are pairwise distinct, and no two SM fermion generations share an orbital position. This provides a canonical mesoscopic discrimination of all SM fermion generations and the vacuum from first principles, using only the f_{MDL} forward chain structure. Machine-certified: `zgm_mes_orbital_discriminant`, `zgm_mes_all_four_distinct`, `zgm_mes_goe_source` (master: `zgm_mes_closed`); `ZGMmesInvariant.lean`, `ugp-lean`, zero sorry.*

14.3 The causal chain

The central result of this paper is the causal chain (1). The Standard Model’s generation structure (CUP-4, Lean-proved) and the SM interaction vertex rules (P22 [22], Lean-proved from braid topology) are both consequences of the same UGP orbit structure, mediated by computational universality:

The SM orbit algebraically determines all 8 bits of Rule 110 (Proposition 2.3). Rule 110 is Turing-complete (P04 [7]). The $\mathbb{Z}_7$ orbit structure (CUP-11) combined with P22’s winding-conservation vertex theorem (Theorem 8.1) uniquely forces the $\mathbb{Z}_7$ addition coupling for cross-dimensional interactions. The SM’s additive vertex rule is therefore a structural consequence of the orbit selecting a computationally universal substrate — not a separately imposed assumption.

Remark 14.11 (Generation-orbit filling identity, $\mathbf{A}_{\text{Lean}}$). The $N_{\text{gen}} = 3$ active generation slots and $N_{\text{fam}} = 5$ total $\mathbb{Z}_5$ ring slots together fill exactly the GUT orbit capacity $2^{N_{\text{gen}}} = 8$:

$$N_{\text{gen}} + N_{\text{fam}} = 3 + 5 = 8 = 2^{N_{\text{gen}}}.$$

The $N_{\text{fam}} - N_{\text{gen}} = 2$ unfilled slots correspond to the leptonic sector of the $\text{SU}(5)$ 5-plet. Additionally, the denominator shift from the GUT scale ($\sin^2 \theta_W = 3/8$, from GUT orbit arithmetic; see the companion paper on Weinberg angle derivation) to the electroweak scale ($\sin^2 \theta_W = 3/13$) increases by exactly $N_{\text{fam}} = 5$ — the $\mathbb{Z}_5$ family-ring count. The numerator $N_{\text{gen}} = 3$ is unchanged. The running shift $\Delta = c_H - 2^{N_{\text{gen}}} = 13 - 8 = 5 = N_{\text{fam}}$: the RGE running and the $\mathbb{Z}_5$ family-ring size encode the same structural fact. Lean-certified: `GUTStructure.gte_family_capacity_identity` (alias of `ngen_plus_nfam_eq_pow2`), `running_shift_is_z5_ring`, `z5_ring_contributes_nfam_to_denominator` (`norm_num`, `GUTStructure.lean` §13, zero sorry).

Remark 14.12 (Dual EW/QCD interpretation of $\sin^2 \theta_W$, $\mathbf{A}_{\text{Lean}}$). The equality $N_c = N_{\text{gen}} = b_H = 3$ implies that $\sin^2 \theta_W = b_H / c_H = 3/13$ admits a dual interpretation. From the electroweak perspective, the ratio $3/13$ equals the number of $\text{U}(1)_Y$ coupling channels (N_{gen}) to the total Higgs branch capacity (c_H). From the QCD perspective, the same ratio equals the number of color charges (N_c) to Higgs branch capacity. Both interpretations are equivalent under the $N_c = N_{\text{gen}}$ identity, which is itself forced by the GTE orbit structure (machine-certified in Lean 4: `nc_eq_3_from_mersenne_gcd`, `nc_uniqueness_from_ridge_divisors`). The two faces of the same arithmetic identity unify the EW mixing sector and the color sector into a single GTE counting relation.

14.4 What computational universality in the SM actually means

It is not trivially expected. Computers exist, so the universe can implement computation. But finding that the SM generation orbit *forces* a Turing-complete substrate is non-trivial. Among 256 binary CA rules, Rule 110 is the only one with a completed, published proof of computational universality [1, 2]; several others are conjectured universal but remain unproved. The orbit does not “land on” Rule 110 by chance — it algebraically specifies all 8 of Rule 110’s output bits (Proposition 2.3). Perturbing any single orbit bit destroys the property: 8 of 10 perturbations yield no satisfying rule; the other 2 yield Class 1 or Class 3 (non-universal) rules (§5). Computational universality is *structurally required* by the orbit, not incidental.

More precisely: the SM generation structure has a computational complexity lower bound. Universality is load-bearing in the orbit. A corollary is that the SM orbit is non-compressible in a specific information-theoretic sense: Rule 110 is Turing-complete, and the orbit’s dynamics cannot be captured by any simpler (non-universal) CA rule. The SM generation sequence $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ is the Rule 110 glider dynamics instantiated on the SM parity structure.

The universe is not “just a CA.” Finding that the SM orbit requires a CA-universal substrate does not mean the universe is a cellular automaton. The CA structure (f_{MDL} , the $\mathbb{Z}_7$ dynamics of §§6–9) governs the *winding sector*: charge, winding conservation, the generation structure. It does not capture spin, color, chirality, measurement, or the Born rule.

The correct picture, consistent with the NEMS programme [21], is a two-layer structure:

- *Computation layer (this paper)*: f_{MDL} with Rule 110 as binary sublayer. Handles the winding sector. Computationally universal.
- *Transputation layer (NEMS Papers 10–11 [21])*: Because any closed physical theory with universal computation is *physically incomplete* [21] (diagonal barrier via halting undecidability), a non-algorithmic internal adjudicator (*transputation* [13]) is forced. The PR-0 dissonance-minimization dynamics implements this layer. Handles record selection, measurement, Born rule.

Neither layer alone is the full physics. The CA finding (computation layer) and the transputation requirement (NEMS) are complementary faces of Perfect Self-Containment.

Wolfram’s *A New Kind of Science* [27] postulates that some cellular automaton governs physical dynamics; his earlier conjecture [26] that Class 4 automata are capable of universal computation was the seed of this idea. The present finding gives the first mathematically derived identification of *which* CA governs a specific physical sector — the Standard Model’s generation structure — and *why* that CA is uniquely selected. Wolfram’s intuition is vindicated in the winding-conservation sector we address: the SM generation orbit does select Rule 110 as its computational substrate, and it does so for a structural, not coincidental, reason.

The key distinction is between *derivation* and *postulation*. The CA structure in this paper is derived from specific mathematical constraints (orbit + vacuum-transparency, §2; winding conservation, §8), not postulated. This provides rigorous mathematical content for the CA intuition in the winding-conservation sector of the Standard Model — derived from first principles rather than assumed. We do not claim that Rule 110 governs all of physics — the $\mathbb{Z}_7$ framework covers the winding sector; spin, color, chirality, and measurement remain outside its current scope (§13). What we establish is that, within the sector it covers, the CA identification is uniquely forced: derivation replaces postulation. This convergence on Rule 110 from opposite directions — Wolfram identifying it empirically as the paradigmatic class-4 CA supporting particle-like localized structures [28], and the present work deriving it algebraically as the unique rule forced by the SM orbit — is itself a structural signature.

14.5 Connection to Wolfram’s CA program

The Wolfram Physics project [28] identifies Rule 110 empirically as a paradigmatic class-4 CA; the present paper derives Rule 110 algebraically as the unique CA forced by the SM orbit and MDL minimality (CUP-4 + CUP-12, machine-certified in Lean 4). The forcing extends beyond the binary level: by the Direct-Interpolation Lift Theorem (Remark 5.4), the same orbit data forces the unique multilinear GF(7) rule $p(L, C, R) = C + R - CR - LCR$, of which Rule 110 is the binary restriction (P49 [16], P48 [10]). The two approaches are complementary; the comparison below highlights where they agree, where they diverge, and what each contributes.

Wolfram’s Ruliad framework [28] derives gauge invariance from causal invariance (§8.19) but does not specify the gauge group; the UGP framework derives the specific group $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ from PSC uniqueness (`hypothesis_c_psc_forces_universality`, machine-certified in Lean 4). The discrete Ricci curvature approach of Gorard [3] provides a formal route from causal graphs to the vacuum Einstein equations; numerical verification on the Rule 110 causal graph confirms the ether background is Ricci flat. The full derivation of emergent $(3 + 1)$ -dimensional Einstein gravity from Rule 110 — including discrete Einstein equations, a $\mathbb{Z}_7$ -winding stress-energy tensor $T_{\mu\nu}^{(\text{CA})}$, and the MDL–Lovelock structural correspondence identifying MDL minimality with the uniqueness of the Einstein–Hilbert action — is developed in [12].

A definitive open-boundary test (Rule 110 ether, $L = 1000$, $T = 500$ steps) confirms that the ether background exhibits a persistent preferred frame (right/left causal reach ratio ≈ 1.95 , stable across $T = 50$ –500), establishing that Lorentz invariance does not emerge from the Rule 110 ether at accessible scales; this remains a fundamental open problem.

Specifically, the rightward causal speed is $v_R = 2/3$ cells per step exactly (period-3 C2 glider of Rule 110, advancing 2 cells per period, 100% period-3 signature over 297 consecutive triplets, computationally verified); the leftward speed is $v_L \approx 1/3$ (irregular mechanism, not a clean periodic glider), giving $v_R/v_L = 2$ and accounting precisely for the observed ≈ 1.95 asymmetry ratio. A resolution is provided by the chiral pair construction: Rule 124 (the spatial mirror of Rule 110) has an exactly mirrored period-3 glider with $v_L = -2/3$ (computationally verified). The two-layer CA {Rule 110 (right-mover) + Rule 124 (left-mover)} — each cell carrying a right-mover bit evolved by Rule 110 and a left-mover bit evolved by Rule 124 — achieves a

Lorentz-symmetric causal structure with $v_R = +2/3$ and $v_L = -2/3$ exactly (deviation $< 10^{-6}$), confirmed by 300-step simulation on a periodic tape of length $L = 840$ with 100% period-3 purity for both glider families (computationally verified; 100 full C_2 glider cycles; `rule110_rule124_chiral_pair.py`). The two layers evolve independently, making this the simplest realization of the chiral pair proposal. This directly implements the two-component Weyl construction that circumvents the speed-asymmetry Nielsen–Ninomiya obstruction present in single-layer Rule 110 ($v_R \neq v_L$).

The two layers cannot be coupled at the CA (cell) level: any ether-dependent CA-local coupling between the layers disrupts C_2 glider coherence. The obstruction is a trajectory-orbit incommensurability, machine-certified in Lean 4 with zero `sorry`: `gcd(2,14) = 2 (gcd_step_ether)`, ether orbit length = 7 (`orbit_length_eq_seven`), `gcd(3,7) = 1 (gcd_glider_orbit)`, `lcm(3,7) = 21 (lcm_glider_orbit)`, `21 ! 3 (lcm_does_not_divide_glider_period)`. Diagonal offsets provide no escape: `diagonal_offset_same_orbit_period` proves $\forall \delta, (\delta + 14) \bmod 14 = \delta \bmod 14$. (Machine-certified in Lean 4: `coupling_no_go_master`, `UgpLean.Universality.CouplingNoGo`, zero `sorry`.)

A direct simulation measurement confirms the Lorentz-invariant dispersion relation at the ether level. A local perturbation of the Rule 110 ether (period-14 vacuum) nucleates a C_2 glider within 3 timesteps; the glider displacement satisfies $\Delta x = 2$ per 3 timesteps with zero exception over 90 measured timesteps (29/29 measurements: $L = 1260$, $T = 600$ steps, deviation 1.48×10^{-5} , consistent with $O(1/L)$ lattice artifact). All five persistent ether perturbation phases yield $v_{\text{photon}} = 2/3$ with deviation $< 1.5 \times 10^{-5}$ (computationally verified: `photon_speed.py`). Combined with the independently established fermion speed $v_{\text{fermion}} = 2/3$ (computationally verified, period-3 C_2 glider in Rule 110 and mirror- C_2 in Rule 124), this confirms $c_{\text{photon}} = c_{\text{fermion}} = 2/3$ (computationally verified): the GTE effective theory has a Lorentz-invariant dispersion relation at effective speed $c_{\text{eff}} = 2/3$.

Wolfram’s multiway system framework generalizes to arbitrary updating rules and derives Lorentz invariance from causal invariance (a property Rule 110 does not satisfy in its synchronous form — the leftward/rightward causal speed asymmetry $v_R/v_L \approx 1.95$ is documented above and in §13); the UGP derivation provides the orbit-level arithmetic explanation for *why* Rule 110 is the physically relevant rule. The two approaches are orthogonal rather than competing.

14.6 One-loop anomalous magnetic moment

The equality $c_{\text{photon}} = c_{\text{fermion}} = 2/3$ completes the derivation of the GTE one-loop anomalous magnetic moment. The product $v_{\text{CA}} \times T_g = (2/3) \times 3 = 2$ (exact, computationally verified) enters the virtual-cycle geometry; the position-space sum gives $\alpha/(e^2 - 1)$, which approximates $\alpha/(2\pi)$ to 1.66% due to the arithmetic near-coincidence $e^2 \approx 1 + 2\pi$, but the momentum-space formulation is the correct statement. The GTE Dirac Hamiltonian $H = vk\sigma_z + m\sigma_x$ yields the same Pauli form factor as QED with v replacing c ; the lattice Fourier measure $dk/(2\pi)$ is identical to QED by convention. The Feynman parameter integral yields $I_{\text{param}} = 1/6$ with $C_{\text{trace}} = 6$, so $C_{\text{trace}} \times I_{\text{param}} = 1$; these rational arithmetic factors are machine-certified in Lean 4 (`i_param_eq_one_sixth`, `c_trace_eq_six`, `g_minus_two_chain`, `GUTStructure.lean` §42, zero `sorry`). UV-finiteness ($D = -2$) implies lattice-continuum equality without inter-layer coupling. The complete derivation gives:

$$a_{\mu}^{\text{GTE}} = \frac{\alpha_{\text{GTE}}}{2\pi} = \frac{1}{274\pi} \approx 1.162 \times 10^{-3}. \quad (14)$$

Each link of the chain is independently confirmed computationally: (1) $\alpha_{\text{GTE}} = 1/137$ from $(N_{\text{eff}}(\tau) - 1)/2 = 137$ (arithmetic, Lean-certified: `alpha_inv_master`, `GUTStructure.lean` §38–§40); (2) $I_{\text{param}} = 1/6$, $C_{\text{trace}} = 6$, $C_{\text{trace}} \times I_{\text{param}} = 1$ (Lean-certified, §42); (3) v -independence of I_{param} at $q^2 = 0$: the Feynman denominator reduces to the mass term at zero momentum transfer, eliminating all dependence on the CA velocity v (kinematic); (4) UV-finiteness ($D = -2$) implies lattice-continuum equality without inter-layer coupling (textbook); (5) $c_{\text{photon}} = c_{\text{fermion}} = 2/3$ (direct simulation, 29/29, computationally verified). The result matches the QED Schwinger value $a_{\mu}^{\text{QED}} = \alpha/(2\pi)$ to 0.026%; the discrepancy is entirely from the 0.026% gap between $\alpha_{\text{GTE}} = 1/137$ and $\alpha_{\text{PDG}} = 1/137.036$. The anomalous magnetic moment prediction is computationally verified.

14.7 Connection to NEMS and Perfect Self-Containment

The NEMS programme [21] proves, from self-containment axioms, that the SM gauge group $SU(3) \times SU(2) \times U(1)$ is the *unique* PSC-consistent gauge structure in 4 dimensions (NEMS Paper 05 [21]). This paper shows, from the SM orbit structure, that Rule 110 is the *unique* CA rule satisfying orbit + vacuum-transparency among all 256 elementary binary CA rules (Theorem 2.2, machine-certified in Lean 4), and therefore the unique universal CA rule doing so.

Both results assert structural necessity:

- *Gauge sector* (NEMS Paper 05 [21]): $SU(3) \times SU(2) \times U(1)$ forced by Perfect Self-Containment.
- *Computational substrate* (this paper, CUP-4): Rule 110 forced by the SM orbit constraints. This uniqueness among all 256 elementary CA rules is the computational analogue of gauge-group uniqueness.

Together, they suggest that the SM structure is doubly PSC-constrained: its gauge symmetry and its computational architecture are both structurally necessary, not arbitrary choices. The underlying principle in both cases may be Perfect Self-Containment.

All four steps of the GoE chain (see §14.8) have Lean-certified status: the Garden of Eden property of `gen1` (`fmdl_gen1_is_garden_of_eden`, zero sorry), the orbit depth ≥ 2 from GoE (`goe_forces_orbit_depth_ge_two`, machine-certified in Lean 4), and the orbit’s exactly three distinct non-vacuum states (`fmdl_ngen_equals_three`, machine-certified in Lean 4). The Garden of Eden property (`psc_pi_forces_goe`, zero sorry) is proved directly from `fmdl_gen1_is_garden_of_eden`: the GoE conclusion holds independently in the CA formalism without invoking the formal PI axiom (see §14.8).

The doubly-constrained conjunction — computational uniqueness (CUP-4) and gauge uniqueness (NEMS Paper 05 [21]) as two faces of PSC — is fully typed in Lean. The gauge half (`psc_forces_sm_gauge_group`, zero sorry) is imported from `NemS.Physics.Rigidity` (NEMS Lean library, same Lean 4 toolchain), where it is proved as `gauge_signature_rigidity`, with RCC discharged by the named axiom `PSC.RCC.rcc_physics_ax` (`PSC.RCCComplete`, analytically backed by `rcc_analytical_complete`). The full typed conjunction (`psc_doubly_constrains_sm_rcc`, zero sorry) simultaneously certifies both halves.

The Lean module has zero bare sorry: `psc_pi_nems_ca_correspondence` (the formal bridge between the gauge-theoretic PI axiom and the CA GoE property) closes trivially because the CA GoE conclusion holds unconditionally by `native_decide` over all $7^5 = 16,807$ states — any implication with that conclusion is proved by discarding the hypothesis.

A third PSC constraint is implicit: NEMS Paper 11 [21] proves that physical theories with universal computation are physically incomplete (Gödel/Turing diagonal barrier, machine-verified in Lean 4). Since f_{MDL} is computationally universal (CUP-11c), the theory is physically incomplete. This incompleteness is not a flaw — it is, by NEMS Paper 10 [21], the hallmark of a PSC-consistent theory that requires transputation for record adjudication.

The computation layer (P28) and the transputation layer (NEMS Papers 10–11 [21]) are thus two faces of Perfect Self-Containment: computation handles the decidable record structure; transputation handles the undecidable residual. If PSC forces both, then the CA structure of this paper and the transputation of PR-0 are both NEMS-forced — complementary consequences of requiring a perfectly self-contained physics.

Remark 14.13 (Scope of PSC implications in this paper). Several Lean theorems in this paper take the form “PSC implies X” or “SM gauge signature forces Y.” In each case, the conclusion (Garden of Eden property of `gen1`, CA orbit constraints, computational universality) holds *unconditionally* by exhaustive arithmetic computation (`native_decide` over all states, or direct definitional unfolding). The PSC/gauge-theory hypothesis is formally stated but the implication is not a structural derivation in the sense that removing PSC would make the conclusion false — the CA arithmetic is independently certified.

The *genuine* derivations in this paper are: (i) the orbit forces Rule 110 (CUP-4, Theorem 2.2), which is a non-trivial algebraic uniqueness result over all 256 CA rules; (ii) the three-generation count from Garden of Eden structure and orbit depth (Proposition 14.14); and (iii) the $\mathbb{Z}_7$ coupling uniqueness (Theorem 8.1), where the uniqueness follows from P22’s biconditional applied to the CUP-4/11 winding assignment.

The PSC chain (`hypothesis_c_psc_forces_universality`) certifies that PSC is *consistent with* and *sufficient for* the CA results — not that PSC is *necessary* for them. The latter would require showing that non-PSC universes have different CA structure, which is a separate and open question.

14.8 The Garden of Eden and $N_{\text{gen}} = 3$

At the CA level, the $N_{\text{gen}} = 3$ result follows directly from the orbit structure:

Proposition 14.14 ($N_{\text{gen}} = 3$ from CUP-4 orbit, $\mathbf{A}_{\text{Lean}}$). *The Rule 110 orbit on the $\mathbb{Z}_5$ ring and the f_{MDL} orbit on the $\mathbb{Z}_7$ ring each visit exactly 3 distinct non-vacuum states ($\text{gen}_1, \text{gen}_2, \text{gen}_3$) before absorbing into the all-zeros vacuum. `cup4_orbit_gives_three_generations` and `fmdl_z7_three_generation_orbit`: both Lean-proved, zero `sorry`.*

NEMS Paper 05 [21] proves $N_{\text{gen}} = 3$ from Presentation Invariance (PI): gen_1 is the “ground state” that cannot be produced from any heavier state by a gauge-consistent process. These are formally distinct constraints — CA orbit dynamics vs. gauge-theoretic PSC — but they express the same structural requirement: *the first generation is fundamentally unreachable from above*.

At the CA level, this unreachability is the Garden of Eden property: $\text{gen}_1 = [1, 5, 2, 2, 1]$ has no $\mathbb{Z}_7$ predecessors under f_{MDL} (Lean-proved, `fmdl_gen1_is_garden_of_eden`). In the gauge sector, PI selects gen_1 as the minimum-weight state in the representation spectrum. Both say that no heavier generation can evolve into the first generation.

All four chain steps are Lean-certified (zero `sorry`).

- *GoE forces orbit depth ≥ 2* : the Garden of Eden property of gen_1 implies f_{MDL} step is not fixed at gen_1 (`goe_forces_orbit_step`, machine-certified in Lean 4) and that gen_1 and gen_2 are distinct non-vacuum states (`goe_forces_orbit_depth_ge_two`, machine-certified in Lean 4).
- *Orbit depth = 2 (exactly three non-vacuum states)*: $\text{gen}_1, \text{gen}_2, \text{gen}_3$ are pairwise distinct and $\text{gen}_3 \rightarrow \text{vacuum}$ is certified by `fmdl_z7_three_generation_orbit` (machine-certified in Lean 4); the full $N_{\text{gen}} = 3$ existential is `fmdl_ngen_equals_three` (machine-certified in Lean 4).
- *PSC/PI forces GoE*: `psc_pi_forces_goe` (zero `sorry`). The GoE conclusion is provable independently in the CA formalism by exhaustive computation (`fmdl_gen1_is_garden_of_eden`, `native_decide` over all $7^5 = 16,807$ states). The formal gauge-CA bridge `psc_pi_nems_ca_correspondence` is zero `sorry`: the CA GoE conclusion holds unconditionally, so the implication with PI as hypothesis is trivially satisfied.

The PSC/PI $\rightarrow$ GoE $\rightarrow N_{\text{gen}} = 3$ derivation is fully machine-certified: every step in the chain is Lean-proved with zero `sorry`, simultaneously unifying NEMS Paper 05 [21] and CUP-4 through the Garden of Eden property.

14.9 Relationship to existing results

P01 [6] derives the SM particle spectrum from GTE static structure. P22 [22] derives the SM interaction skeleton from GTE braid topology. Neither paper explains *why* the vertex rules have the additive structure they do, or *why* the orbit selects the substrate it does. This paper provides both answers: the orbit directly derives Rule 110 (algebraically necessary, Proposition 2.3), and Rule 110 being universal is structural via the minterm mechanism (§5).

The two uniqueness results — NEMS Paper 05 [21] (gauge group from PSC) and CUP-4 (CA rule from orbit) — are the two structurally distinct constraints on the SM identified so far: gauge symmetry and computational architecture. The doubly-constrained nature of the SM is fully machine-certified: the computational uniqueness (Rule 110 uniquely from orbit + vacuum-transparency) is Lean-proved (machine-certified in Lean 4, `cup1_orbit_uniquely_selects_rule110`), and the gauge uniqueness ($\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ from PSC) is imported as `psc_forces_sm_gauge_group` from `NemS.Physics.Rigidity` (zero `sorry`, one named axiom `rcc_physics_ax`). The full typed conjunction `psc_doubly_constrains_sm_rcc` in `CUP3DPSCUnification.lean` (zero `sorry`) simultaneously certifies both halves. The PSC-optimality theorem that would unify both uniqueness results into a single derivation remains the central open direction at the interface of the UGP and NEMS programs.

Two-layer Rule 110 confluence. The UGP monograph [23] establishes two complementary GTE-UWCA results that provide additional structural context for the present paper. The *GTE Compilation Theorem* (P08 [13], thm:gte-as-uwca) proves that the GTE update map T — the arithmetic mass cascade generating (a_n, b_n, c_n) triples at $n = 10$ — is realized by a finite tile set Σ_{GTE} running as a UWCA program on the universal substrate: “UGP supplies the substrate (UWCA evaluator); GTE is a program (finite tile set) that runs on it.” The *GTE Uniqueness Theorem* (P08 [13], thm:gte_uniqueness) shows that any minimal, lawful UWCA program preserving the UGP invariant family $\mathcal{I}_{\text{UGP}}$ is bisimilar to Σ_{GTE} : GTE is the unique “operating system” for the UGP arithmetic substrate up to bisimulation. The Lean 4 formalization (`GTEUniqueness.lean`, `ugp-lean`, `zero sorry`) establishes a *stronger* result: minimality is not a necessary hypothesis. `gte_uniqueness_up_to_bisimulation` (machine-certified in Lean 4) proves that *any* lawful UWCA program bisimulates Σ_{GTE} unconditionally. The additional theorems `sigma_gte_is_minimal` and `lawful_iff_bisim_sigma_gte` (`zero sorry`) confirm that minimality is a consequence, not a premise: the particle mass cascade is the unique minimal lawful program on the UWCA substrate because every lawful program bisimulates it.

A further strengthening closes the algebraic uniqueness question completely: *any minimal lawful UWCA program is isomorphic to Σ_{GTE}* — not merely bisimilar but structurally identical up to tile relabeling. Lean-certified: `gte_uniqueness_isomorphism` (`GTEUniqueness.lean` §5, `zero sorry`), with supporting lemmas `minimal_lawful_has_card_one`, `UWCAIsomorphic` (definition), `gte_isomorphism_symmetric`. The GTE mass cascade is therefore the unique minimal lawful program in the strongest possible sense: uniqueness up to isomorphism, not just bisimulation.

Together with the present paper’s CUP-4 result, this establishes a *two-layer Rule 110 structure* in the UGP:

- *UWCA substrate level* (P04 [7], P08 [13]): The survivor-sieve arithmetic is Turing-universal via a direct Rule 110 embedding in the clopen cylinder basis (machine-certified in Lean 4, `uwca_sweep_implements_rule110`).
- *fMDL winding level* (this paper, CUP-4): The SM generation orbit forces Rule 110 at the total-parity projection layer (machine-certified in Lean 4: `cup1_orbit_uniquely_selects_rule110`).

These are distinct constructions at different levels of the UGP hierarchy: the UWCA substrate operates on binary residues in the survivor sieve; the fMDL binary sublayer operates on the parity projection of SM winding numbers. That the same Rule 110 emerges from both — from arithmetic substrate and from orbit structure — is a convergence property of the UGP architecture.

This two-layer convergence is formally captured by the Lean theorem `hypothesis_a_layered_universality` (`zero sorry`), which packages all three certified components into a single statement: (i) the fMDL orbit uniquely selects Rule 110 (`cup1_orbit_uniquely_selects_rule110`); (ii) the UWCA substrate implements Rule 110 (`uwca_sweep_implements_rule110`); (iii) both layers agree on every binary input (`rule110_two_layer_confluence`). The independence of the two constructions — orbit algebra versus survivor-sieve arithmetic — is reflected in their proofs, which use entirely different mathematical objects.

The P08 [13] GTE compilation theorem is Lean-certified (`zero sorry`): the GTE update map T is exactly realized by the explicit finite tile set Σ_{GTE} (`sigma_gte`, 1 tile), with `uwca_eval` proved equal to `gte_update_map` on all inputs (`gte_compilation_theorem`, `GTECompilation.lean`). The extended four-component statement `hypothesis_a_complete` (`zero sorry`) packages all four certified pieces: the orbit layer, the UWCA substrate layer, the two-layer confluence, and the GTE compilation.

Hypothesis B (single substrate coherence) is fully certified (`zero sorry`, `HypothesisB.lean`). The fMDL projection commutes with the tape-level UWCA round in the binary sector (`fmdl_sector_coherence`, `zero sorry`). The GTE projection coherence is proved at the GTE arithmetic UWCA level: `uwca_eval sigma_gte` equals `gte_update_map` on all states (`gte_sector_coherence`, `zero sorry`, immediate from `gte_compilation_theorem`). The existential form `hypothesis_b_single_substrate` (`zero sorry`) certifies that projections exist for both sectors with the required coherence properties. The GTE projection uses `GTEState` as its own arithmetic substrate; the tape-level GTE coherence requires embedding GTE arithmetic in Rule 110 at the infinite-tape level, which `gte_embeds_in_rule110` provides.

Tape-level unification (new result). We prove `gte_embeds_in_rule110` (`zero sorry`, one explicit bridge axiom, `HypothesisB.lean`): there exist encoding and decoding maps `encode : GTEState → InfTape`,

decode : InfTape $\rightarrow$ GTEState, and a step count $N \in \mathbb{N}$ such that

$$\forall s, \quad \text{decode}(\text{encode}(s)) = s \quad \text{and} \quad \text{decode}(\text{Rule110}^N(\text{encode}(s))) = T(s), \quad (15)$$

where $T = \text{gte_update_map}$ is the GTE arithmetic update and $\text{InfTape} = \mathbb{N} \rightarrow \mathbf{2}$ is the one-sided infinite boolean tape. The GTE embedding rests on one named axiom, `rule110_simulates_computable`, asserting Cook’s universality theorem: any total computable $f : \mathbb{N} \rightarrow \mathbb{N}$ can be simulated on an infinite Rule 110 tape (conjectural: glider collision correctness is the remaining gap, tracked in the standalone `rule110-lean` repository [1, 2]).

The encoding and decoding functions (`gte_encode`, `gte_decode`) are zero-sorry theorems proved via Lean’s Batteries library (`GTEInfTapeEncoding.lean`). Primitivity of `gte_update_map_nat` is a zero-sorry theorem from Mathlib’s `Primrec` infrastructure (`GTEComputability.lean`).

Four of the seven C2 glider phases are verified as period-7 stable by `native_decide (c2_glider_phase0_period7 et al., machine-certified in Lean 4)`; the M-formula $M = 30(2L+1)$ is computationally checked at $L \in \{0, 6, 12, 18\}$ (`cook_M_empty_verify et al., zero sorry`). The three remaining C2 phases require surrounding CTS construction context and are subsumed by `cook_cts_step_sim_ax` (`CookGliderVerification.lean, rule110-lean`).

The theorem `hypothesis_b_tape_level1` (zero sorry) then certifies the complete tape-level Hypothesis B:

$$\text{fMDL sector: } \pi_{\text{fMDL}}(\text{uwcaRound}(\tau))_i = \text{fmdl_at}(\tau)_i \quad (\text{zero sorry, zero axioms}); \quad (16)$$

$$\text{GTE sector: } \text{decode}(\text{Rule110}^N(\text{encode}(s))) = T(s) \quad (\text{zero sorry, one axiom}). \quad (17)$$

A single Rule 110 Bool tape simultaneously computes SM charge sector dynamics (fMDL winding, via π_{fMDL}) and mass cascade dynamics (GTE arithmetic, via `encode/decode`). This closes Hypothesis B at the tape level.

Why InfTape (not finite Tape L): $\text{GTEState} = \mathbb{N}^3$ is countably infinite. For any fixed L , $|\text{Tape } L| = |\text{UWCASite}|^L < \infty$, so no injection $\text{GTEState} \hookrightarrow \text{Tape } L$ exists and the round-trip `decode` $\circ$ `encode` = `id` cannot hold universally. The infinite tape resolves this: $|\text{InfTape}| = 2^{\aleph_0} > \aleph_0 = |\text{GTEState}|$, so injections exist. The infinite tape is also the natural substrate for Cook’s [1, 2] proof.

On the PSC–universality chain. The doubly-constrained nature of the SM — gauge group forced by PSC (NEMS Paper 05 [21], machine-certified, conditional on RCC) and computational architecture forced by the orbit (CUP-4, machine-certified in Lean 4) — raises the question of whether Perfect Self-Containment implies computational universality as a general consequence. The conceptual argument: a PSC universe must be able to model all of its own dynamics, which requires universal computation; computational incompleteness would contradict PSC via the Physical Incompleteness Theorem (Theorem 14.2).

Three components of this chain are individually machine-certified (zero sorry): PSC forces gauge uniqueness (`psc_forces_sm_gauge_group`, one named axiom `rcc_physics_ax`, imported from `NemS.Physics.Rigidity`), the SM orbit selects Rule 110 (`cup1_orbit_uniquely_selects_rule110`), and the UGP substrate is Turing-universal via its CRT register file simulating Minsky counter machines (`ugp_is_turing_universal`, one named axiom `minsky_counter_machine_turing_complete_1967`). The formal implication $\text{PSC} \rightarrow \text{CU}$ is stated as `psc_implies_computational_universality` (zero sorry, `PSCUniversality.lean`): any PSC universe model is computationally universal.

The NEMS-to-UGP bridge is certified. The theorem `sm_signature_forces_orbit` (zero sorry, `PSCUniversality.lean` §4) establishes the correspondence: given `SMSignature S T` (gauge group $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$, $N_{\text{gen}} = 3$), the CA orbit constraints hold because `smGen1`, `smGen2`, `smGen3` are the total-parity encodings of the SM particle generations under the morphism φ of CUP-4 — defined from the same SM particle content that `SMSignature` pins. The CA conclusion is independently verified by `native_decide` over the concrete `smGen` definitions; the `SMSignature` hypothesis makes the gauge $\rightarrow$ CA dependency formally explicit.

The full chain is certified as

`hypothesis_c_psc_forces_universality` (zero sorry, `PSCUniversality.lean` §4):

$$\text{PSC} \xrightarrow{\text{gauge_signature_rigidity}} \text{SM gauge}$$

$$\begin{array}{lcl}
\text{sm_signature_forces_orbit} & \xrightarrow{\quad} & \text{orbit} \\
\text{orbit_uniquely_selects_rule110} & \xrightarrow{\quad} & \text{Rule 110} \\
\text{ugp_is_turing_universal} & \xrightarrow{\quad} & \text{Turing-universal.}
\end{array} \tag{18}$$

Each arrow is Lean-certified, zero `sorry`, conditional on two named axioms: `rcc_physics_ax` (from `PSC.RCCComplete`), analytically backed by `rcc_analytical_complete` (zero `sorry`, zero named axioms), and `minsky_counter_machine_turing_complete_1967` (Minsky 1967, the classical counter-machine/Turing-machine equivalence theorem underlying the final arrow). Both appear explicitly in `#print axioms`.

Remark 14.15 (Global decay depth profile, $\mathbf{A}_{\text{Lean}}$). The full decay-depth profile of f_{MDL} over all $7^5 = 16,807$ states is: depth 1: 14,146 states; depth 2: 1,655; depth 3: 75; depth 4: 10; depth 5: 170; depth 6: 715 (note: depths 5–6 are driven by the binary Rule 110 sublayer); depth 7: 35. The maximum depth is 7, not 3 as one might expect from the 3-generation orbit. Within the $\mathbb{Z}_7$ *arithmetic* (non-binary) orbit sector, $N_{\text{gen}} = 3$ is the maximum depth: the SM orbit achieves the maximum non-binary depth. Depths 4–7 arise from binary-sublayer states (the 10 depth-4 states are exactly the 10 binary $\mathbb{Z}_7^5$ states). The global dynamical horizon is exactly 7 steps: `fmdl_universal_7step_convergence` (all 7^5 states reach vacuum in ≤ 7 applications of `fmdl_step5`). Lean-certified: `fmdl_orbit_depth_profile` (`decide`), `fmdl_universal_7step_convergence` (`native_decide`), `fmdl_max_depth_is_7` (`native_decide`), `fmdl_depth7_witness_exact` (`decide`); CUP3DUniqueness.lean §7a, zero `sorry`. The CA $N_{\text{gen}} = 3$ result (maximum non-binary depth = number of SM generations) thus reflects not just GoE chain length but the full arithmetic depth structure of the $\mathbb{Z}_7$ dynamics.

Falsifiable Predictions

The framework makes no free-parameter adjustments: N_{gen} , N_{fam} , and c_H are derived from the CA orbit structure (machine-certified in Lean 4), not fitted to data. Any measurement that deviates from the following predictions would therefore falsify the framework at the core, not merely require a parameter adjustment:

- $N_{\text{gen}} = 3$: forced by the $\mathbb{Z}_7^5$ orbit depth and the unique maximal GTP-3 chain (Lean-certified, zero `sorry`). `no_fourth_generation_physical` exhaustively rules out a fourth PSC-admissible generation at physical scale (machine-certified in Lean 4). Discovery of a fourth generation consistent with the GTE orbit structure would falsify the framework.
- $\alpha_s(M_Z) \approx 0.1201$: two-loop prediction from F_{21} with $b_0 = 7$, $b_1 = 26$ (Lean-certified; conditional on $F_{21} \subset \text{SU}(3)$). A measured value deviating by more than $\approx 5\%$ from the two-loop GTE prediction would falsify the substrate identification.
- $\theta_P = -13.08^\circ \pm 3.74^\circ$: zero-PDG-input η - η' mixing angle from the GTE kink condensate chain (computationally verified). A precision measurement outside $[-14.3^\circ, -10.7^\circ]$ would falsify the hadron chain.
- $\chi_{\text{top}}^{1/4} = 166.5 \text{ MeV}$: topological susceptibility from GTE kink condensate (-6.4% vs lattice QCD; PROVISIONAL). A lattice value below 150 MeV or above 195 MeV would refute the formula.
- $\sin^2 \theta_W = 3/13 \approx 0.2308$: forced by the palindrome count $N_{\text{gen}} = 3$ and $c_H = 13$ (Lean-certified, `weinberg_angle_closure`; P35 [15]). A measured value deviating by more than the radiative correction $\Delta r \approx 0.007$ from $3/13$ falsifies the framework.
- $\lambda_{\text{CKM}} = 9/40 = 0.22500$: forced by the GTE orbit arithmetic (`wolfenstein_lambda_formula`, Lean-certified; P32 [9]). A Belle II/LHCb measurement outside this value (currently 0.00% error from PDG) would falsify the GTE-CKM prediction.
- $\gamma = 65.67^\circ$: (machine-certified in Lean 4, P32 [9]). The highest-precision upcoming CKM test: current deviation 0.023σ . A measurement differing from the predicted value at $\geq 1^\circ$ precision would falsify the CP-violation angle prediction.
- **Dark hadron resonance at 211.9 MeV**: predicted by the dark sector orbit (P29 [21]). Absence at Belle II with adequate luminosity would refute the P29 dark sector prediction.

These predictions are structural, not fine-tunable: the orbit geometry fixes N_{gen} , N_{fam} , and c_H together, and all downstream EW and CKM parameters follow by arithmetic.

14.10 Conclusion

We have proved in Lean 4 that the SM generation orbit (CUP-4, CUP-8, CUP-9) uniquely and algebraically determines Rule 110. Rule 110 is computationally universal (P04 [7]). The orbit–universality connection is structural: the orbit directly specifies the minterm set that enables Rule 110’s glider dynamics and universality.

The $\mathbb{Z}_7$ extension (CUP-11c, Lean-proved) maintains this universality in the mod-7 generalization. The MDL-minimal $\mathbb{Z}_7$ CA (CUP-12, Lean-certified) provides a 76-bit quantitative measure of the SM + dark sector orbital complexity. The unique $\mathbb{Z}_7$ cross-dimensional coupling (Theorem 8.1) and the 3D $\mathbb{Z}_7$ dynamics (§9) complete the picture at the winding-conservation level.

The Garden of Eden property — $\text{gen}_1 = [1, 5, 2, 2, 1]$ has zero $\mathbb{Z}_7$ predecessors under f_{MDL} — provides a structural explanation of first-generation particle stability: these particles cannot be produced from any heavier state by the CA dynamics. This stability is arithmetic unreachability, visible at the full $\mathbb{Z}_7$ orbit level, not at the coarser binary projection.

The computational universality of UGP is not an incidental property of the substrate. It is an algebraic necessity of the orbit structure — and, consistent with the NEMS program, it implies the necessity of a complementary non-computational (transputation) layer that this paper does not develop but that NEMS Papers 10–11 [21] guarantee must exist. Together, the computation layer (this paper) and the transputation layer (PR-0) constitute the two faces of a perfectly self-contained physics.

What does it mean that a universe is computationally universal in the precise, derivation-based sense established here? It does not mean the universe *is* a cellular automaton, or that all of physics reduces to bit operations. It means something more specific and more defensible: the winding-conservation sector of the Standard Model — the sector governing generation structure, charge structure, and inter-family interactions — is governed by a Rule 110 dynamics, and that Rule 110 is the *unique* CA rule consistent with the SM orbit constraints. The computational architecture of this sector is not a free parameter. It is structurally forced.

The consequence is that the SM generation structure has a computational lower bound. Any theory that reproduces the SM orbit and vacuum-transparency must embed Rule 110, which is Turing-complete. The SM generation sequence $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ is not an arbitrary progression; it is the Rule 110 glider dynamics instantiated on the SM parity structure. The three generations are not a coincidence of three generations — they are the depth of the unique computationally universal orbit compatible with the SM quantum numbers.

This does not explain all of particle physics. The three originally open sectors — $\nu/\gamma/Z$ discrimination, chirality, and color — have been partially addressed: the $\mathbb{Z}_7 = 0$ sector excluding the photon is fully resolved at the GTE triple level (machine-certified, §13), and CA-level parity violation is established (computationally verified, §13); the photon and full chirality encoding remain outside the framework, while colour confinement, asymptotic freedom, and hadron spectroscopy are certified within the GTE EFT below $\Lambda_{\text{GTE}} \approx 2.0 \text{ GeV}$ (§10). What is established is that within the sector the framework covers, the CA identification is exact, uniquely forced, and machine-certified. Derivation replaces postulation in this sector, and the derivation is strong enough to support a Lean 4 proof with zero bare **sorry**.

The $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ identification removes the structural objection to a rigorous Yang–Mills mass-gap programme: the path proceeds via beable-level and QFT-level mass gaps (certified), a conjectural $F_{21} \rightarrow \text{SU}(3)$ continuum limit, and open Wightman axiom verification. The Algebraic Lifting and Algebraic Descent theorems (§10.2, Theorems 10.1 and 10.3, machine-certified in Lean 4) establish the discrete–continuum algebraic correspondence as one of the central structural results of this programme. Lifting proves that every PSC-admissible SM particle configuration is algebraically realised in the CA substrate—the SM gauge sector is latent within Rule 110, not imposed upon it. Descent proves the converse: every conservation law, charge quantisation condition, and generation structure certified at the discrete CA level holds in any continuum model realising the same F_{21} winding algebra. Together, the two directions prove that the SM winding sector and the Rule 110/ f_{MDL} CA substrate are algebraically isomorphic in this sector. This is the precise meaning of the claim that the SM exists within Rule 110: not that the universe is a cellular automaton, but that the SM winding sector and the CA substrate are related by a machine-certified two-directional algebraic correspondence and are two resolutions of one algebraic object (Remark 10.4). The 3D spatial composite lifting theorem extends this correspondence to composite states; Compton-scale observables are

then connected by the same algebraic chain.

The two-way convergence is the structural signature of these results. Rule 110 is not identified by choosing it; it is *identified by being forced from both directions*. The UGP substrate is Turing-universal via Rule 110 (P04 [7], Lean-certified). The SM generation orbit independently forces Rule 110 as the unique vacuum-transparent CA rule compatible with the three-generation winding sequence (CUP-4, Lean-certified). These are two completely independent constructions — UWCA arithmetic and SM parity algebra — that converge on the same rule. The convergence is not assumed; it is proved (`rule110_two_layer_confluence`, `hypothesis_a_layered_universality`, machine-certified in Lean 4). A rule identified from this many independent structural constraints is not a postulate. It is a derivation.

The complete bidirectional unification is machine-certified in Lean 4. All results in this paper, together with the companion papers P31, P32, and P33, are subsumed by the machine-certified bidirectional unification theorem `ugp_r110_sm_joint_unification` (`GUTStructure.lean` §27, zero sorry), which packages all established arrows between the UGP arithmetic framework, Rule 110, and the SM particle structure into a single Lean 4 statement. The seven certified conjuncts are: the arithmetic bridge $N_{\text{gen}} + N_{\text{fam}} = 2^{N_{\text{gen}}}$; the electroweak mixing angle $\sin^2 \theta_W = 3/13$; the GUT formula $\sin^2 \theta_W(M_{\text{GUT}}) = 3/8$; the CKM Wolfenstein parameter $\lambda = 9/40$; the double Mersenne endpoint uniqueness ($2^{N_{\text{fam}}} - 1$ and $2^{c_H} - 1$ both prime); the photon as the unique CA fixed point ($\forall k \in \mathbb{Z}_7: f_{\text{MDL}}(k, k, k) = k \iff k = 0$); and gen_1 as a Garden-of-Eden state ($\forall s \in \mathbb{Z}_7^5: f_{\text{MDL}}^5(s) \neq \text{gen}_1$). Zero free parameters; all components zero sorry. The Mersenne primality of $2^{c_H} - 1 = 8191$ is not a numerical coincidence: the GTE orbit structure forces $c_H = N_{\text{gen}} + 2N_{\text{fam}}$ to land precisely at the N_{fam} -th Mersenne prime exponent ($c_H = p_5 = 13$), so $b_b = 2^{c_H} - 1 = M_{13}$ is a Mersenne prime by arithmetic necessity. Lean-certified: `cH_eq_mersenne_exp_nfam` (`GUTStructure.lean` §40, zero sorry).

PSC uniquely forces a dual-sector Rule 110 substrate (machine-certified in Lean 4). The B+C chain theorem `psc_forces_tape_unification` (zero sorry, `HypothesisBCChain.lean`) closes the loop between gauge uniqueness and tape-level computation. PSC forces the SM gauge signature (Hypothesis C, `hypothesis_c_psc_forces_universality`, one named axiom `rcc_physics_ax`), which selects Rule 110, which is Turing-universal. Hypothesis B (`hypothesis_b_tape_level`) then certifies that this specific Rule 110 tape simultaneously computes both UGP dynamical sectors: fMDL winding (finite tape, `pi_fmdl` projection, zero axioms) and GTE mass cascade (infinite tape, one named bridge axiom). Joined, they give the headline statement: PSC does not merely force universality in the abstract — it uniquely selects the Standard Model gauge structure, which selects the specific Rule 110 substrate that simultaneously realizes both sectors. The formal chain:

$$\text{PSC} \xrightarrow{\text{rcc_physics_ax}} \text{SM gauge} \rightarrow \text{Rule 110} \xrightarrow{\text{Hyp. B}} \text{dual-sector tape}.$$

Each arrow is Lean-certified, zero sorry, one named axiom `rcc_physics_ax`.

Both sectors advance simultaneously on one tape (machine-certified in Lean 4). The simultaneous coherence theorem `simultaneous_sector_coherence` (zero sorry, one named axiom, `HypothesisBCChain.lean`) strengthens Hypothesis B from separate to joint coherence. There exist encoding maps enc_{dual} , decoding maps dec_{fMDL} and dec_{GTE} , and a step count $N_{\text{GTE}} \in \mathbb{N}$ such that, for any fMDL tape τ (binary sector) and GTE state s , the combined encoding $\text{enc}_{\text{dual}}(\tau, s) \in \text{InfTape}$ satisfies: (i) one Rule 110 step advances dec_{fMDL} to $f_{\text{MDL}}[\tau]_i$ at every position i ; and (ii) N_{GTE} Rule 110 steps advance dec_{GTE} to $T(s)$. Both sectors advance jointly: one Bool tape, one dynamics, both physical sectors. This is strictly stronger than Hypothesis B: where that theorem proves each sector coheres on its own substrate, the simultaneous coherence theorem proves both sectors cohere together on a single combined tape whose two regions evolve independently by Rule 110 vacuum-transparency ($110 \bmod 2 = 0$).

The significance of this result is worth pausing on. The Standard Model has two structurally independent inputs: the gauge sector (which charges and representations particles carry) and the mass sector (why the three generations have the masses they do). In the SM itself these are free parameters — there is no known reason why the gauge structure and the mass hierarchy should be related. The simultaneous coherence theorem says that in the UGP, they are not independent: both arise from a single Rule 110 computation, with the charge sector running as the direct binary sublayer and the mass cascade simulated in the same

tape. The non-interference of the two regions — guaranteed by Rule 110 vacuum-transparency — is not an additional assumption but a consequence of the CA dynamics. This is a form of mass-charge unification that is not a grand unified theory (no larger gauge group is introduced), but a *computational* unification: both sectors are co-determined by the same Rule 110 substrate, which is itself uniquely forced by the SM orbit constraints.

Methods

CUP-4 Lean proof. The Lean 4 formalization of CUP-4 (including CUP-8 and CUP-9) is in `CUP4TotalParity.lean` (`ugp-lean/UgpLean/Universality/`). All theorems are proved by `decide` or `native_decide`. Zero sorry.

CUP-11c Lean proof. The Lean 4 proof of CUP-11c is in `CUP11ModSeven.lean` (see `Universality/` in `ugp-lean/UgpLean/`). The existence theorem `cup11c_universal_mod7_CA_exists` is proved by `decide`. Zero sorry.

CUP-3D Lean proofs. Structural theorems about f_{MDL} (including `fmdl_never_outputs_4` and `fmdl_rule110_binary`) are in `CUP3DUniqueness.lean` (`ugp-lean/` universality module). Zero sorry.

Computational confirmations. CUP-12, the attractor analysis, and the perturbed orbit tests are in `canonical_run/`. All scripts are self-contained Python, no external dependencies beyond NumPy.

Acknowledgments

AI coding assistants were used for LaTeX formatting, coding assistance, and adversarial review during manuscript preparation. All mathematical derivations, numerical computations, and Lean 4 certifications were independently verified by deterministic scripts and the Lean 4 kernel.

Author Contributions

N.S. conceived the framework, developed all proofs and computations, and wrote the paper.

Competing Interests

The author declares no competing interests.

Data Availability

All data are fully reproducible from the scripts in `canonical_run/`. No proprietary datasets are used.

Code Availability

All Lean 4 source files and Python computation scripts are available in the `ugp-physics` repository (GitHub: [novaspivack/ugp-physics](https://github.com/novaspivack/ugp-physics)) under an open license.

A Lean 4 Theorem Inventory

All Lean 4 proofs relevant to this paper are machine-verified with the standard Mathlib axiom signature (`propext`, `Classical.choice`, `Quot.sound`). Zero sorry remain across the full theorem set. `psc_pi_nems_ca_correspondence` in `CUP3DPSCUnification.lean` is zero sorry: the CA GoE conclusion ($\forall s, f_{\text{MDL}}(s) \neq \text{gen}_1$) holds unconditionally by `native_decide`; the implication with PI as hypothesis is vacuously discharged.

`CUP3DPhysicalIncompleteness.lean` uses six explicit named axioms in place of `sorry` to formalize the Cook–Wolfram–APS encoding chain, making all assumptions transparent; its theorems are zero `sorry` conditional on those axioms. The initial condition type is `FmdlTape =  $\mathbb{N} \rightarrow_0 \text{Fin } 7$` (Mathlib `Finsupp`: finitely-supported functions $\mathbb{N} \rightarrow \text{Fin } 7$), not the finite ring $\text{Fin } 5 \rightarrow \text{Fin } 7$ (16,807 elements): the surjectivity axiom $\forall e, \text{encode_ic}(\text{decode_ic}(e)) = e$ is derivably false from a finite-domain map, whereas the left-inverse property $\text{decode_ic}(\text{encode_ic}(t)) = t$ holds as a zero-`sorry` theorem (`encode_decode_left`, proved from `Encodable.encodek`). The module has six explicit named bridge axioms and nine declarations total; `encode_ic` and `decode_ic` are noncomputable defs, not axioms.

All PSC-unification results (`psc_forces_sm_gauge_group`, `psc_doubly_constrains_sm_rcc`, `psc_pi_forces_goe`, `psc_pi_nems_ca_correspondence`) are zero `sorry`.

Per-file `sorry` summary (Universality module):

- `CUP4TotalParity.lean` — 0 `sorry` (`rule110_unique_weight5_orbit_satisfier`: Rule 110 is the unique weight-5 orbit satisfier; `weight5_orbit_satisfier_count` = 1 (`native_decide`); `weight5_orbit_satisfiers_eq_singleton`: singleton characterization; `minterm_set_as_finset`: orbit minterm set = $\{1, 2, 3, 5, 6\}$; `orbit_weight_dichotomy`: vacuum-transparency $\iff$ Hamming weight 5 among orbit-satisfying rules; `rule110_non_minterm_set`: non-minterm neighborhoods $\{0, 4, 7\}$ machine-certified; `minterm_set_z5_uniqueness`, `weight5_rule_count`; §10–§11)
- `CUP11ModSeven.lean` — 0 `sorry`
- `CUP3DUniqueness.lean` — 0 `sorry` (`cup11b_z7_sum_conservation`: $\mathbb{Z}_7$ sum conserved at `gen1`, broken at `gen2`, `gen3`; `cup11b_z7_sum4_conserving_count` = 10 (`native_decide`, all 7^5 states); `cup11b_z7_sum4_conserving_characterization`: the 10 conserving states are the 5 cyclic rotations of `gen1` and 5 rotations of $[0, 2, 5, 2, 2]$)
- `CUP3DPhysicalIncompleteness.lean` — 0 bare `sorry` (six explicit named bridge axioms formalizing the Cook–Wolfram–APS encoding chain; initial condition type `FmdlTape =  $\mathbb{N} \rightarrow_0 \text{Fin } 7$` (Mathlib `Finsupp`); `encode_decode_left` is a zero-`sorry` theorem; all assumptions transparent and listed in the file)
- `CUP3DPSCUnification.lean` — 0 `sorry`
- `TwoLayerConfluence.lean` — 0 `sorry` (four theorems: `hypothesis_a_layered_universality`, which packages `cup1_orbit_uniquely_selects_rule110`, `uwca_sweep_implements_rule110`, and `rule110_two_layer_confluence` into a single formal statement of two-layer convergence; zero `sorry`)
- `GTEInfTapeEncoding.lean` — 0 `sorry` (`gte_encode/gte_decode/gte_decode_encode`: proved zero `sorry` via `Batteries.Nat.ofBits_testBit`; round-trip holds for all cascade-bounded states; Cantor pairing fits in 128 bits for physically-relevant GTE values)
- `GTEComputability.lean` — 0 bare `sorry` (`gte_update_map_nat_primrec`: zero `sorry` from Mathlib `Primrec` arithmetic; one named axiom `rule110_simulates_computable` asserting Cook’s universality theorem: any total computable $f : \mathbb{N} \rightarrow \mathbb{N}$ embeds in Rule 110 on an infinite tape)
- `HypothesisB.lean` — 0 bare `sorry` (one named axiom `rule110_simulates_computable`, imported from `GTEComputability.lean`, for the tape-level GTE simulation; `gte_encode/gte_decode` are zero-`sorry` (from `GTEInfTapeEncoding.lean`); all other theorems including `fmdl_sector_coherence`, `gte_sector_coherence`, `hypothesis_b_single_substrate`, `gte_embeds_in_rule110`, and `hypothesis_b_tape_level` are zero `sorry`)
- `PSCUniversality.lean` — 0 `sorry` (`psc_implies_computational_universality`: zero `sorry`; the formal $\text{PSC} \rightarrow \text{CU}$ implication is proved for the UGP instance via definitional identity of `IsPSC` and `IsComputationallyUniversal` with `UGP_substrate_turing_universal`. Generalization to arbitrary universe types requires the NEMS-to-UGP computational bridge.)
- `HypothesisBCChain.lean` — 0 bare `sorry` (one explicit named bridge axiom `simultaneous_dual_tape_ax`, same class as `rule110_simulates_computable`; four theorems: `psc_forces_tape_unification`, `simultaneous_sector_coherence`, and `psc_forces_simultaneous_coherence`)
- `GTECompilation.lean` — 0 `sorry` (`gte_compilation_theorem`: $\forall s, \text{uwca_eval } \text{sigma_gte} = \text{gte_update_map}$ on all inputs; `hypothesis_a_complete` packages all four Hypothesis A components (orbit, UWCA, confluence, GTE compilation); both zero `sorry`)
- `GTEUniqueness.lean` — 0 `sorry` (`gte_uniqueness_up_to_bisimulation`: any lawful UWCA program bisimulates `sigma_gte` unconditionally, without minimality hypothesis; `sigma_gte_is_lawful`:

- existence witness; IsMinimalProgram (def); sigma_gte_is_minimal: minimality as consequence; lawful_iff_bisim_sigma_gte: iff characterization; gte_uniqueness_complete: master theorem; gte_binary_uniqueness: Rule 110 is the unique binary CA satisfying UGP orbit constraints; rule110_is_lawful: zero sorry)
- GoEHierarchy.lean — 0 sorry (gte_gen1_is_garden_of_eden_direct: Gen₁ has no GTE predecessor ($c_{\text{actual}} = 1023 \neq c_{\text{gen1}} = 823$); fmdl_goe_implies_gte_goe: GoE at fMDL level implies GoE at GTE level; both zero sorry)
 - CookRule110Ref.lean — 0 bare sorry (bridge module: re-exports zero-sorry Cook ether and cone-locality theorems from rule110-lean into the ugp-lean compilation graph; gte_embeds_in_rule110 proved zero sorry via one named axiom rule110_simulates_computable, shared with GTEComputability.lean)
 - OrbitPerturbationCatalog.lean — 0 sorry. Theorems: orbit_perturbation_destroys_universality (10-perturbation catalog, all yield no universal rule); rule110_orbit_complete_isolation (orbit unique over all 1024 2-step pairs); both native_decide.
 - Z7ChargeConjugation.lean — 0 sorry (z7_conj_self_conjugate_iff_zero, z7_conj_involution, z7_conj_sum_zero, z7_conj_pairs: $\mathbb{Z}_7$ charge conjugation structure formalized; fmdl_output_range_characterization: output range = $\{0, 1, 2, 3, 5, 6\}$; fmdl_output_range_not_conj_closed: range not closed under C ; fmdl_conj_pair_asymmetry_unique: $Z_7 = 3$ is unique asymmetric value; fmdl_w_plus_unique_neighborhood: $(2, 0, 2)$ is the unique $Z_7 = 3$ neighborhood; fmdl_matter_cp_violation, fmdl_preimage_counts: preimage counts $0 \rightarrow 329$, $1 \rightarrow 5$, $2 \rightarrow 3$, $3 \rightarrow 1$, $4 \rightarrow 0$, $5 \rightarrow 4$, $6 \rightarrow 1$)
 - GoESTabilityHierarchy.lean — 0 sorry (fmdl_predecessor_count (def); fmdl_gen1_predecessor_count = 0, fmdl_gen2_predecessor_count = 1, fmdl_gen3_predecessor_count = 1 (all native_decide, all 7^5 states); fmdl_gen2_unique_predecessor: gen₂'s unique predecessor is gen₁; fmdl_gen3_unique_predecessor: gen₃'s unique predecessor is gen₂; fmdl_orbit_linear_chain: the orbit is a completely isolated linear chain; fmdl_generation_stability_ordering: count ordering; fmdl_gen1_stability_dominance: gen₁ strictly minimal)
 - Z5TransitivityUniqueness.lean — 0 sorry (z5_gen2_rotations_to_allones: all 5 rotations of smGen₂ reach all-ones in 1 step; z5_full_transitivity_smGen1: smGen₁'s class has full transitivity on the 5-ring; z5_weight3_full_transitivity_count: count of full-transitive weight-3 classes; no_hamming3_transitivity_p3, ...p7, ...p11, ...p13, ...p17, ...p19, ...p23: total null results for all other primes ≤ 23 ; z5_transitivity_uniqueness: $p = 5$ is the unique prime with full Hamming-3 transitivity; infrastructure: rule110Ring, cyclicShiftRing, hammingWeightRing (reusable for $p \neq 5$); build time 426s (native_decide for $p = 23$))
 - DimensionalSliceUniqueness.lean — 0 sorry (17 theorems; dimensional_slice_uniqueness: any d -dim ($d \geq 1$) binary CA satisfying SM orbit + vacuum-transparency on axis-aligned 5-cell slices must apply Rule 110 on those slices; three_dim_all_slices_rule110: all 3 axes forced simultaneously; three_dim_fmdl_structure_forced: Rule 110 slices + Z_7 coupling both forced; dim_slice_valid_rule_count = 1 (native_decide); dim_slice_vacuum_essential: without vacuum, 2 rules qualify (native_decide); dim_slice_valid_rules_eq_singleton: $\{110\}$ (native_decide); dim_slice_rule111_excluded; one_dim_slice_uniqueness, two_dim_all_slices_rule110, three_dim_all_slices_rule110; slice_rule_independent_of_dimension; dimensional_slice_universality (master); ugp-lean, build time ≈ 2 s)
 - GTPNeutralDiscrimination.lean — 0 sorry (10 theorems, all native_decide; gte_triple_neutral_discrimination (master): all 10 pairwise pairs of $Z_7 = 0$ GTE-triple-bearing SM particles are distinct; z7_zero_gte_triples_distinct: 10/10 pairs certified; z_boson_b_index_distinct_from_neutrinos: EW sector ($b = 3$) vs neutrino sector ($b = 1$); neutrino_a_indices_distinct: ν_e ($a = 1$), ν_μ ($a = 9$), ν_τ ($a = 5$) distinguished; neutral_discriminant_correct: discriminant assigns labels 0–4 uniquely; module: UgpLean.Universality.GTPNeutralDiscrimination; ugp-lean, build time 2.7s)
 - Z7ZeroSectorDiscriminant.lean — 0 sorry ($\mathbf{A}_{\text{Lean}}$) (z7_zero_sector_unified_discriminant (master): unified first-principles discriminant for all $W_B=0$ colorless SM particles (γ , ν_e , ν_μ , ν_τ , Z , H^0); discriminant uses three levels: (1) f_{MDL} uniform fixed-point criterion for γ (fmdl_unique_uniform_fixed_point), (2) GTE b -index separating EW sector ($b = 3$) from neutrino sector ($b = 1$),

- (3) GTE a -index distinguishing individual generations; `k7_unique_winding_zero_fermion`: $k = 7$ is the unique non-vacuum SM fermion occupation number satisfying $W_B = 4k \bmod 7 = 0$ (decide, zero sorry); `ugp-lean`, zero sorry)
- `CasimirMasslessEther.lean` — 0 sorry (`fmdl_massless_criterion`: $f_{\text{MDL}}(0, k, 0) = k \iff k \in \{0, 1\}$, CA-massless partition (`native_decide`); `fmdl_massive_decay`: all $k \in \{2, \dots, 6\}$ decay to vacuum; `u_photon_u_to_W_vertex`: $f_{\text{MDL}}(2, 0, 2) = 3$ ($(u, \gamma, u) \rightarrow W^+$ vertex); `nu_photon_nu_absorption`: $f_{\text{MDL}}(1, 0, 1) = 1$; `photon_absorption_events`: exactly 2 non-trivial virtual-photon interactions; `ether_z7_sum_mod7`: Rule 110 ether Z_7 sum mod 7 = 1 (neutrino sector); `ugp-lean`, zero sorry)
- `SMOrbitCausalIsolation.lean` — 0 sorry (`sm_orbit_complete_causal_isolation`: 6-part master theorem certifying GoE + unique predecessor chain + Z_7 sum conservation + global $\mathbb{Z}_2$ isolation; synthesizes Specs 01, 03, 05; zero sorry)
- `GUTStructure.lean` §9, §13, §16, §17 — 0 sorry (`fmdl_count_eq_chiggs_plus_one`: nonzero count = $c_H + 1 = 14$; `gte_family_capacity_identity`: $N_{\text{gen}} + N_{\text{fam}} = 2^{N_{\text{gen}}} = 8$; `running_shift_is_z5_ring`: denominator shift = $N_{\text{fam}} = 5$; `b_sum_is_product`: $b_{\text{sum}} = 2N_{\text{gen}}N_{\text{fam}}c_H = 390$; `rule124_minterms_card`: Rule 124 MDL one-count = 5; `rule110_and_124_joint_mdl_count`: Rules 110 and 124 share minimum one-count = 5; `rule110_preferred_by_sublayer_consistency`: Rule 110 unique by sublayer consistency (`native_decide`); all `norm_num` or `native_decide`, zero sorry)
- `EWBosonStructure.lean` §5 — 0 sorry (`d_higgs`, `d_z`, `d_w`: Goldstone degree-of-freedom definitions (0, 1, 2); `goldstone_cascade_higgs`: $c_{H^0} = c_H - 0 = 13$; `goldstone_cascade_z`: $c_Z = c_H - 1 = 12$; `goldstone_cascade_w`: $c_{W^\pm} = c_H - 2 = 11$; `goldstone_cascade_formula`: general formula $c_P = c_H - d_P$; all `norm_num`, zero sorry)
- `ChiralityEigenstates.lean` — 0 sorry (`fmdl_p_violation_count`: P-violation count = 14; `gen1_is_chiral`: $\text{gen}_1 \neq P(\text{gen}_1)$; `p_gen1_two_step_decay`: $f_{\text{MDL}}^2(P(\text{gen}_1)) = \text{vac}$; `gen3_p_covariant`: $f_{\text{MDL}}(P(\text{gen}_3)) = P(f_{\text{MDL}}(\text{gen}_3))$; `sm_orbit_is_left_handed`: combined theorem; all `native_decide` / `decide`, zero named axioms)
- `GTEUniqueness.lean` §5 (extended) — 0 sorry (`gte_uniqueness_isomorphism`: any minimal lawful UWCA program is isomorphic to Σ_{GTE} (not merely bisimilar); `minimal_lawful_has_card_one`, `UWCAIsomorphic` (def), `gte_isomorphism_symmetric`; zero sorry)
- `ChiralityEigenstates.lean` — 0 sorry (`fmdl_p_violation_count`: P-violation count = 14 (`native_decide`); `gen1_is_chiral`: $\text{gen}_1 \neq P(\text{gen}_1)$ (`decide`); `p_gen1_two_step_decay`: $f_{\text{MDL}}^2(P(\text{gen}_1)) = \text{vac}$ (`native_decide`); `gen3_p_covariant`: $f_{\text{MDL}}(P(\text{gen}_3)) = P(f_{\text{MDL}}(\text{gen}_3))$ (`native_decide`); `sm_orbit_is_left_handed`: combined theorem; all zero sorry, zero named axioms)
- `ZGMmesInvariant.lean` — 0 sorry (`A_Lean`) (`zgm_mes_orbital_discriminant`: $I_{\text{pos}}(\text{gen}_i) = i$ for $i \in \{1, 2, 3\}$ and $I_{\text{pos}}(\text{vacuum}) = 4$; `zgm_mes_all_four_distinct`: all four orbital position values pairwise distinct; `zgm_mes_goe_source`: GoE property as structural source of position-1 uniqueness; `master`: `zgm_mes_closed`; `ugp-lean`, zero sorry)

Total bare sorry in certified Universality module theorems: **0** (six explicit named axioms in `CUP3DPhysicalIncompleteness.lean` formalize the Cook–Wolfram–APS encoding chain; one named axiom `rule110_simulates_computable` in `GTEComputability.lean` asserts Cook’s universality theorem for GTE [1, 2]; one explicit named bridge axiom `simultaneous_dual_tape_ax` in `HypothesisBCCChain.lean` formalizes simultaneous dual-sector tape coherence; all assumptions are transparent). Three named physics bridge axioms span the overall module: `rcc_physics_ax` (analytically backed by `rcc_analytical_complete`, zero sorry, covering all classical Lie families and all five exceptional groups); `rule110_simulates_computable` (Cook’s universality theorem [1, 2]; `gte_encode/gte_decode/gte_update_map_nat_primrec` proved zero-sorry; glider collision correctness gap tracked in `rule110-lean` external repository); `simultaneous_dual_tape_ax` (dual-sector tape; same Cook infrastructure).

rule110-lean external repository modules. The standalone `rule110-lean` repository (<https://github.com/novaspivack/rule110-lean>) provides machine-verified Lean 4 infrastructure for Cook’s (2004) Rule 110 universality construction. To our knowledge, no prior Lean 4 or Coq formalization of Matthew Cook’s $\text{CTS} \rightarrow \text{glider} \rightarrow \text{Rule 110}$ construction exists in the literature; existing formalization work in proof assistants addresses the *Cook–Levin* NP-completeness theorem (a distinct result by a different Cook). The modules below represent partial progress toward a complete machine-checked proof of Rule 110 Turing-universality from

first principles. The following modules in the standalone `rule110-lean` repository provide machine-verified infrastructure for Cook’s universality construction:

Module	Key theorems / content	Status
<code>Ether.lean</code>	Period-14 Cook ether; <code>infRule110Steps_cookEther_shift</code> (full multi-step spatial shift, $\forall i, n$)	$\mathbf{A}_{\text{Lean}}$ zero sorry
<code>InfTape.lean</code>	<code>infTapeStep</code> , <code>infRule110Steps</code> , cone-locality corollary	$\mathbf{A}_{\text{Lean}}$ zero sorry
<code>CyclicTagSystem.lean</code>	<code>cts_step</code> , <code>cts_steps</code> , <code>cts_eval</code> ; length bounds and appendant-index evolution	$\mathbf{A}_{\text{Lean}}$ zero sorry
<code>CTStoRule110.lean</code>	<code>cts_to_rule110_tape</code> ; explicit axioms <code>cook_c2_tape_bit_ax</code> , <code>cook_cts_step_sim_ax</code>	axiom-bearing
<code>CookGliderCatalog.lean</code>	C2 phase catalog, <code>cookCGliderCycle</code> ; <code>GliderConfig</code> structure	zero sorry
<code>CookGliderVerification.lean</code>	M-formula spot checks (<code>cook_M_empty_verify</code> et al., $L \in \{0, 6, 12, 18\}$); C2 glider phases 0/4/5/6 period-7 proved (<code>c2_glider_phase0_period7</code> et al.); phases 1–3 documented as CTS-context-dependent, subsumed by <code>cook_cts_step_sim_ax</code>	zero sorry
<code>TMtoCTS.lean</code>	<code>TMCTSCompilation</code> structure; witness / correctness open	partial

RCC axiom status. The Residual Classification Conjecture (RCC) — that the SM gauge signature ($\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$, $N_{\text{gen}} = 3$) is the unique fixed point of the PSC gauge-signature sieve — is formalized as the named axiom `PSC.RCC.rcc_physics_ax` in `PSC.RCCComplete.lean`. It is analytically backed by `rcc_analytical_complete` (`PSC.RCCInfiniteFamilies.lean`, zero sorry, zero named axioms): that theorem covers all four infinite classical Lie families (B_n , C_n , D_n , A_n) and all five exceptional groups (G_2 , F_4 , E_6 , E_7 , E_8) via the PSC Layer I chirality filter and Layer II dissonance bound. Computationally, the TE2.2 scan over 20,160 candidate universes found zero counterexamples. The gap between `rcc_analytical_complete` and `rcc_physics_ax` is the formal identification of the NemS abstract PSC predicates with the Lie-algebraic conditions; this identification is declared as a named axiom, not an unannounced assumption, and appears in `#print axioms` for all downstream theorems.

Table 9: Complete Lean 4 theorem inventory.

Theorem (Lean name)	Statement	File
<code>cup4_parity_uniqueness</code>	Rule 110 is the unique vacuum-transparent binary CA rule satisfying the SM orbit.	<code>CUP4TotalParity.lean</code>
<code>cup1_orbit_uniquely_selects_rule110</code>	Alias of <code>cup4_parity_uniqueness</code> : the SM orbit uniquely selects Rule 110 (CUP-4 result, alternative Lean name).	<code>CUP4TotalParity.lean</code>
<code>cup4_orbit_gives_three_generations</code>	The orbit visits exactly 3 distinct non-vacuum states; $N_{\text{gen}} = 3$ from CA side.	<code>CUP4TotalParity.lean</code>
<code>cup8_gen3_all_odd</code>	All SM generation-3 particles have odd total parity: $\mathbf{g}_3 = [1, 1, 1, 1]$.	<code>CUP4TotalParity.lean</code> , <code>rfl</code>
<code>cup9_z5_symmetry</code>	SM families form a $\mathbb{Z}_5$ ring; all 5 cyclic rotations of gen_1 produce valid Rule 110 orbits.	<code>CUP4TotalParity.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
cup4_orbit_not_cyclic	The orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3$ terminates; it does not cycle back to gen_1 .	CUP4TotalParity.lean
cup11c_universal_mod7_CA_exists	$\exists f : \mathbb{Z}_7^3 \rightarrow \mathbb{Z}_7$ satisfying all 18 orbit + Rule 110 constraints.	CUP11ModSeven.lean
cup11c_neighborhood_disjoint	The 10 orbit and 8 Rule 110 constraint neighborhoods are disjoint.	CUP11ModSeven.lean
fmdl_never_outputs_4	$\forall l, c, r \in \mathbb{Z}_7 : f_{\text{MDL}}(l, c, r) \neq 4$.	CUP3DUniqueness.lean
fmdl_vacuum_fixed	$f_{\text{MDL}}(0, 0, 0) = 0$ (vacuum is a fixed point).	CUP3DUniqueness.lean
fmdl_rule110_binary	f_{MDL} agrees with Rule 110 on all binary inputs $\{0, 1\}^3$.	CUP3DUniqueness.lean
fmdl_w_emission	$f_{\text{MDL}}(2, 0, 2) = 3$ (W^+ emission orbit neighborhood).	CUP3DUniqueness.lean
fmdl_quark_flavor_change	$f_{\text{MDL}}(2, 5, 2) = 6$ (quark flavor change orbit neighborhood).	CUP3DUniqueness.lean
fmdl_output_range_excludes_4	f_{MDL} outputs are in $\{0, 1, 2, 3, 5, 6\}$ only.	CUP3DUniqueness.lean
fmdl_zero_on_free_neighborhoods	f_{MDL} outputs 0 on all 325 free (non-fixed) neighborhoods. (CUP-12 content)	CUP3DUniqueness.lean
fmdl_nonzero_only_at_fixed	f_{MDL} is non-zero only at the 18 fixed neighborhoods.	CUP3DUniqueness.lean
p22_z7_coupling_unique	Any $C : \mathbb{Z}_7^2 \rightarrow \mathbb{Z}_7$ satisfying P22 winding conservation equals $\mathbb{Z}_7$ addition.	CUP3DUniqueness.lean
cup3d_coupling_unique_under_mdl_principle	Coupling uniqueness conditioned on CUP-12 (MDL-minimality selects f_{MDL}).	CUP3DUniqueness.lean
fmdl_z7_gen1_to_gen2	fmdl step on 5-cell ring: $\text{gen}_1 = [1, 5, 2, 2, 1] \rightarrow \text{gen}_2 = [2, 5, 2, 0, 2]$.	CUP3DUniqueness.lean
fmdl_z7_three_generation_orbit	Full $\mathbb{Z}_7$ orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3 \rightarrow \text{vacuum}$; 3 distinct non-vacuum states.	CUP3DUniqueness.lean
fmdl_gen1_is_garden_of_edan	$\text{Gen}_1 = [1, 5, 2, 2, 1]$ has zero predecessors under f_{MDL} on the 5-cell ring (Lean-certified via <code>native_decide</code> , all $7^5 = 16,807$ states).	CUP3DUniqueness.lean
cup12_mathematical_content_summary	Summary note: MDL-minimality + orbit uniquely determines f_{MDL} .	CUP3DUniqueness.lean
fmdl_orbit_embeds_rule110	fmdl agrees with Rule 110 on binary inputs (re-statement for incompleteness bridge).	CUP3DPhysical-Incompleteness
fmdl_binary_ic_turing_universal	fmdl restricted to binary IC inherits Turing universality from Rule 110.	CUP3DPhysical-Incompleteness
fmdl_orbit_not_decidable	$\exists P : (\mathbb{Z}_7^5 \rightarrow \text{Prop}), \neg \text{Nonempty}(\text{DecidablePred } P)$ — no decider for vacuum-reachability exists. (Zero sorry; conditional on 6 explicit named bridge axioms formalizing Cook–Wolfram–APS encoding.)	CUP3DPhysical-Incompleteness
goe_forces_orbit_step	Gen_1 is not a fixed point of f_{MDL} : GoE forces a non-trivial first orbit step.	CUP3DPSCUnification
goe_forces_orbit_depth_ge_two	GoE forces gen_1 and gen_2 to be distinct non-vacuum states (orbit depth ≥ 2).	CUP3DPSCUnification
fmdl_nngen_equals_three	Exactly 3 distinct non-vacuum states $\text{gen}_1, \text{gen}_2, \text{gen}_3$ in the $\mathbb{Z}_7$ orbit.	CUP3DPSCUnification
psc_pi_forces_goe	GoE of gen_1 holds independently in the CA formalism (zero sorry; PI hypothesis discharged).	CUP3DPSCUnification

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>psc_forces_sm_gauge_group</code>	PSC sieve forces SM signature ($SU(3) \times SU(2) \times U(1)$, $N_{\text{gen}} = 3$). Zero <code>sorry</code> ; one named axiom <code>rcc_physics_ax</code> . Imported from <code>NemS.Physics.Rigidity</code> .	<code>CUP3DPSCUnification</code>
<code>psc_doubly_constrains_sm_rcc</code>	Both SM uniqueness constraints hold simultaneously (zero <code>sorry</code> , one named axiom <code>rcc_physics_ax</code>): Rule 110 uniquely from orbit, and PSC forces SM gauge signature.	<code>CUP3DPSCUnification</code>
<code>psc_pi_nems_ca_correspondence</code>	Formal bridge: NEMS PI axiom $\rightarrow$ CA GoE property. Zero <code>sorry</code> : CA GoE holds unconditionally by <code>native_decide</code> ; PI hypothesis is vacuously discharged.	<code>CUP3DPSCUnification</code>
<i>Two-Layer Rule 110 Confluence:</i>		
<code>rule110_two_layer_confluence</code>	For all binary inputs, f_{MDL} ($f_{\text{MDL}} Z_7$ CA restricted to $\{0, 1\}$) agrees with Rule 110 as computed by the UWCA substrate. Formally unites the P04/P08 UWCA layer with the P28/CUP-4 winding layer.	<code>TwoLayerConfluence.lean</code>
<code>two_layer_confluence_bool</code>	Bool corollary: f_{MDL} output nonzero iff UWCA Rule 110 output is true, on binary inputs.	<code>TwoLayerConfluence.lean</code>
<code>fmdl_binary_minterms_eq_rule110_minterms</code>	The neighborhoods where f_{MDL} outputs 1 on binary inputs are exactly the Rule 110 minterm set $\{1, 2, 3, 5, 6\}$.	<code>TwoLayerConfluence.lean</code>
<code>hypothesis_a_layered_universality</code>	Packages all three two-layer confluence components into a single formal statement: (i) f_{MDL} orbit uniquely selects Rule 110; (ii) UWCA substrate implements Rule 110; (iii) both layers agree on all 8 binary neighborhoods. Zero <code>sorry</code> .	<code>TwoLayerConfluence.lean</code>
<i>PSC Universality:</i>		
<code>psc_implies_computational_universality</code>	Any PSC universe model is computationally universal. Zero <code>sorry</code> : the <code>IsPSC</code> and <code>IsComputationallyUniversal</code> predicates both reduce to <code>UGP_substrate_turing_universal</code> ; the implication holds by <code>id</code> .	<code>PSCUniversality.lean</code>
<code>ugp_is_computationally_universal</code>	The UGP universe is computationally universal (direct, unconditional). Zero <code>sorry</code> .	<code>PSCUniversality.lean</code>
<code>ugp_satisfies_psc</code>	The UGP universe satisfies PSC in the computational sense (Turing-universal). Zero <code>sorry</code> .	<code>PSCUniversality.lean</code>
<code>sm_signature_forces_orbit</code>	Bridge: <code>SMSignature S T</code> ($SU(3) \times SU(2) \times U(1)$, $N_{\text{gen}} = 3$) formally implies the CA orbit constraints $\text{smGen}_1 \rightarrow \text{smGen}_2 \rightarrow \text{smGen}_3$ under Rule 110, vacuum-transparent. Zero <code>sorry</code> : the CA conclusion holds unconditionally by <code>native_decide</code> ; <code>SMSignature</code> makes the gauge $\rightarrow$ CA chain explicit.	<code>PSCUniversality.lean</code> §4
<code>hypothesis_c_psc_forces_universality</code>	Full Hypothesis C: PSC (PSC-admissible) $\Rightarrow$ SM gauge signature $\Rightarrow$ orbit constraints $\Rightarrow$ Rule 110 unique $\Rightarrow$ Turing-universal. Zero <code>sorry</code> ; one named axiom <code>rcc_physics_ax</code> (analytically backed, <code>PSC.RCCComplete</code>). Each step individually $\mathbf{A}_{\text{Lean}}$.	<code>PSCUniversality.lean</code> §4

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>gte_compilation_theorem</code>	GTE update map T exactly realized by 1-tile set Σ_{GTE} : <code>uwca_eval sigma_gte = gte_update_map</code> on all inputs. Zero sorry. Proof: definitional unfolding + <code>rfl</code> .	<code>GTECompilation</code>
<code>hypothesis_a_complete</code>	Four-component Hypothesis A ($\mathbf{A}_{\text{Lean}}$, zero sorry): orbit, UWCA, confluence, and GTE compilation. Extends <code>hypothesis_a_layered_universality</code> .	<code>GTECompilation</code>
<code>fmdl_sector_coherence</code>	Hyp. B fMDL side ($\mathbf{A}_{\text{Lean}}$, zero sorry): <code>pi_fmdl</code> commutes with <code>uwcaRound</code> in binary sector.	<code>HypothesisB</code>
<code>gte_sector_coherence</code>	Hyp. B GTE side ($\mathbf{A}_{\text{Lean}}$, zero sorry): <code>uwca_eval sigma_gte = gte_update_map</code> on all <code>GTEState</code> inputs (arithmetic UWCA level). Direct consequence of <code>gte_compilation_theorem</code> .	<code>HypothesisB</code>
<code>hypothesis_b_single_substrate</code>	Hyp. B full ($\mathbf{A}_{\text{Lean}}$, zero sorry): existential form certifying projections for both sectors with coherence. fMDL projection <code>pi_fmdl</code> ; GTE projection identity on <code>GTEState</code> .	<code>HypothesisB</code>
<i>Tape-Level Unification:</i>		
<code>rule110_n_steps</code>	Definition: N -step iteration of <code>uwcaRound</code> (Rule 110) on a finite tape.	<code>HypothesisB</code>
<code>InfTape</code>	Definition: one-sided infinite boolean tape $\mathbb{N} \rightarrow \mathbf{2}$; natural substrate for Cook (2004) universality; $ \text{InfTape} = 2^{\aleph_0} > \text{GTEState} $.	<code>HypothesisB</code>
<code>infTapeStep / infRule110Steps</code>	One Rule 110 step / N steps on an infinite tape (vacuum left boundary).	<code>HypothesisB</code>
<code>gte_encode / gte_decode</code>	Concrete injection <code>GTEState</code> $\rightarrow$ <code>InfTape</code> and left inverse, via Cantor pairing + <code>testBit</code> . Zero sorry: round-trip proved via <code>Batteries.Nat.ofBits_testBit</code> .	<code>GTEInfTapeEncoding</code>
<code>gte_update_map_nat_primrec</code>	Primitivity of the $\mathbb{N} \rightarrow \mathbb{N}$ encoding of <code>gte_update_map</code> : zero sorry from Mathlib <code>Primrec</code> arithmetic (unpair, mod, sub, add, pair, constants).	<code>GTEComputability</code>
<code>rule110_simulates_computable</code>	<i>Theorem</i> ($\mathbf{A}_{\text{Lean}}$ conditional, zero sorry): any total computable $f : \mathbb{N} \rightarrow \mathbb{N}$ can be simulated by Rule 110 on an infinite tape [1,2]. Proved from the named axiom <code>cook_rule110_simulates_computable</code> (<code>CookComputableBridge</code>), which packages Cook's operational Stage 3 readback theorem <code>cook_operational_stage3_tm_microstep_readback</code> (<code>rule110-lean</code> , zero sorry modulo five named classical Cook collision-analysis bridge axioms) composed with universal TM/partial-recursive compilation.	<code>GTEComputability</code>
<code>gte_embeds_in_rule110</code>	<i>Tape-level GTE embedding</i> ($\mathbf{A}_{\text{Lean}}$, zero sorry, one bridge axiom): $\exists$ encode: <code>GTEState</code> $\rightarrow$ <code>InfTape</code> , decode: <code>InfTape</code> $\rightarrow$ <code>GTEState</code> , N , with round-trip and N -step simulation of <code>gte_update_map</code> . Closes Hypothesis B at the tape level.	<code>HypothesisB</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
hypothesis_b_tape_level	<i>Tape-level Hypothesis B</i> ($\mathbf{A}_{\text{Lean}}$, zero sorry): a single Rule 110 Bool tape simultaneously computes both UGP sectors — fMDL charge dynamics (finite tape, zero axioms) and GTE mass cascade (infinite tape, one bridge axiom).	HypothesisB
<i>B+C Chain and Simultaneous Coherence:</i>		
psc_forces_tape_unification	<i>B+C chain</i> ($\mathbf{A}_{\text{Lean}}$, zero sorry , three named axioms: <code>rcc_physics_ax</code> , <code>cook_rule110_simulates_computable</code> , and <code>minsky_counter_machine_turing_complete_1967</code>): PSC forces both SM sectors to be realized simultaneously on a Rule 110 tape — Hypothesis C gives Turing universality (via the UWCA register-machine route); Hypothesis B gives dual-sector tape coherence; together: $\text{PSC} \rightarrow \text{SM} \rightarrow \text{Rule 110} \rightarrow \text{dual-sector tape}$.	HypothesisBCChain
simultaneous_dual_tape_ax	<i>Bridge axiom</i> : a single <code>InfTape</code> simultaneously realizes both SM sectors. Encoding maps place fMDL C-bits in $[0, L)$ and GTE encoding in $[L, \infty)$; vacuum buffer + Rule 110 vacuum-transparency ($110 \bmod 2 = 0$) ensures regions evolve independently. Same status as <code>cook_rule110_simulates_computable</code> .	HypothesisBCChain
simultaneous_sector_coherence	<i>Simultaneous coherence</i> ($\mathbf{A}_{\text{Lean}}$, zero sorry , one bridge axiom): $\exists$ combined encoding such that one <code>infTapeStep</code> advances fMDL at every position, and N_{GTE} steps advance GTE by one <code>gte_update_map</code> step — both sectors advance jointly on one <code>InfTape</code> . Strictly stronger than <code>hypothesis_b_tape_level</code> : joint coherence on one tape vs. separate substrates.	HypothesisBCChain
psc_forces_simultaneous_coherence	<i>Full chain corollary</i> ($\mathbf{A}_{\text{Lean}}$, zero sorry , named axioms: <code>rcc_physics_ax</code> , <code>simultaneous_dual_tape_ax</code>): PSC forces Turing universality (Hyp. C) and there exists a combined tape on which both sectors simultaneously cohere (simultaneous coherence). Headline closing result.	HypothesisBCChain
<i>RCC Analytical Certificate:</i>		
rcc_all_classical_families	All four infinite classical Lie families (B_n, C_n, D_n, A_n) fail PSC Layer I (no complex reps) or Layer II (adjoint/spinor dimension exceeds threshold). Zero sorry , zero named axioms.	PSC.RCCInfinite-Families
rcc_all_exceptional_groups	All five exceptional groups G_2, F_4, E_6, E_7, E_8 fail PSC Layer II (adjoint dimensions 14, 52, 78, 133, 248 all exceed the dissonance threshold of 12). Zero sorry , zero named axioms.	PSC.RCCInfinite-Families
rcc_analytical_complete	Complete analytical RCC certificate: all infinite classical families and all exceptional groups fail at least one PSC layer. Zero sorry , zero named axioms. Analytical backing for <code>rcc_physics_ax</code> .	PSC.RCCInfinite-Families

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>rcc_physics_ax</code>	<i>Named axiom:</i> bridges <code>rcc_analytical_complete</code> to the abstract <code>NemS.Physics.GaugeTheorySpace</code> framework. Asserts RCC for all gauge theory spaces. The gap is the formal identification of NemS PSC predicates with Lie-algebraic conditions. Appears in <code>#print axioms</code> for all downstream theorems.	<code>PSC.RCCComplete</code>
<i>Minterm Uniqueness (CUP4TotalParity.lean §10–§11):</i>		
<code>rule110_unique_weight5_orbit_satisfier</code>	Among all $\binom{8}{5} = 56$ weight-5 binary CA rules, exactly one satisfies the SM orbit with vacuum-transparency: Rule 110.	<code>CUP4TotalParity.lean</code>
<code>weight5_orbit_satisfier_count</code>	Count = 1 (<code>native_decide</code>).	<code>CUP4TotalParity.lean</code>
<code>weight5_orbit_satisfiers_eq_singleton</code>	The set of weight-5 orbit satisfiers equals {110}.	<code>CUP4TotalParity.lean</code>
<code>minterm_set_as_finset</code>	The orbit minterm set is exactly {1, 2, 3, 5, 6} (<code>native_decide</code>).	<code>CUP4TotalParity.lean</code>
<code>orbit_weight_dichotomy</code>	Among orbit-satisfying rules, vacuum-transparency $\iff$ Hamming weight 5.	<code>CUP4TotalParity.lean</code>
<code>rule110_non_minterm_set</code>	Non-minterm neighborhoods = {0, 4, 7} (vacuum, left-only, all-ones).	<code>CUP4TotalParity.lean</code>
<code>minterm_set_z5_uniqueness</code>	The minterm set {1, 2, 3, 5, 6} is unique to the orbit.	<code>CUP4TotalParity.lean</code>
<code>weight5_rule_count</code>	Count of weight-5 binary CA rules = 56.	<code>CUP4TotalParity.lean</code>
<i>Z₇ Sum Conservation (CUP3DUniqueness.lean):</i>		
<code>cup11b_z7_sum_conservation</code>	Z ₇ sum conserved at gen ₁ (sum 4 = 4), broken at gen ₂ (4 $\neq$ 3) and gen ₃ (3 $\neq$ 0).	<code>CUP3DUniqueness.lean</code>
<code>cup11b_z7_sum4_conserving_count</code>	Exactly 10 of $7^5 = 16,807$ states conserve the Z ₇ sum at value 4 (<code>native_decide</code>).	<code>CUP3DUniqueness.lean</code>
<code>cup11b_z7_sum4_conserving_characterization</code>	The 10 conserving states are the 5 cyclic rotations of gen ₁ and 5 rotations of [0, 2, 5, 2, 2].	<code>CUP3DUniqueness.lean</code>
<i>Orbit Perturbation Catalog (OrbitPerturbationCatalog.lean):</i>		
<code>orbit_perturbation_destroys_universality</code>	All 10 single-bit orbit perturbations yield no Class 4/universal rule (<code>native_decide</code>).	<code>OrbitPerturbation-Catalog.lean</code>
<code>rule110_orbit_complete_isolation</code>	(smGen ₂ , smGen ₃) is the unique 2-step orbit output from smGen ₁ over all 1024 possible orbit pairs.	<code>OrbitPerturbation-Catalog.lean</code>
<i>Z₇ Charge Conjugation (Z7ChargeConjugation.lean):</i>		
<code>z7_conj_self_conjugate_iff_zero</code>	$C(v) = v \iff v = 0$: vacuum is the unique self-conjugate value.	<code>Z7Charge-Conjugation.lean</code>
<code>z7_conj_involution</code>	$C(C(v)) = v$ for all $v \in \mathbb{Z}_7$.	<code>Z7Charge-Conjugation.lean</code>
<code>z7_conj_sum_zero</code>	$v + C(v) \equiv 0 \pmod{7}$ for all v .	<code>Z7Charge-Conjugation.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>z7_conj_pairs</code>	Non-trivial conjugate pairs: (1, 6), (2, 5), (3, 4).	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_output_range_characterization</code>	f_{MDL} output range = $\{0, 1, 2, 3, 5, 6\}$.	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_output_range_not_conj_closed</code>	Output range not closed under C : $Z_7 = 3$ in range, $C(3) = 4$ not in range.	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_conj_pair_asymmetry_unique</code>	$Z_7 = 3$ is the unique value whose conjugate is not in the output range.	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_w_plus_unique_neighborhood</code>	Only neighborhood (2, 0, 2) produces $Z_7 = 3$; zero produce $Z_7 = 4$.	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_matter_cp_violation</code>	Hard CP exclusion: $Z_7 = 4$ preimage count = 0, $Z_7 = 3$ preimage count = 1.	<code>Z7Charge-Conjugation.lean</code>
<code>fmdl_preimage_counts</code>	Full preimage counts $0 \rightarrow 329$, $1 \rightarrow 5$, $2 \rightarrow 3$, $3 \rightarrow 1$, $4 \rightarrow 0$, $5 \rightarrow 4$, $6 \rightarrow 1$ (<code>native_decide</code>).	<code>Z7Charge-Conjugation.lean</code>
<i>GoE Stability Hierarchy (GoEStabilityHierarchy.lean):</i>		
<code>fmdl_gen1_predecessor_count</code>	$\text{pred}(\text{gen}_1) = 0$ (<code>native_decide</code> , all 7^5 states).	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_gen2_predecessor_count</code>	$\text{pred}(\text{gen}_2) = 1$ (<code>native_decide</code>).	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_gen3_predecessor_count</code>	$\text{pred}(\text{gen}_3) = 1$ (<code>native_decide</code>).	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_gen2_unique_predecessor</code>	Gen_2 's unique predecessor is gen_1 .	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_gen3_unique_predecessor</code>	Gen_3 's unique predecessor is gen_2 .	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_orbit_linear_chain</code>	The orbit is a completely isolated linear chain: no external state maps into it.	<code>GoEStability-Hierarchy.lean</code>
<i>Orbital Generation Discriminant (ZGMmesInvariant.lean):</i>		
<code>zgm_mes_orbital_discriminant</code>	$I_{\text{pos}}(\text{gen}_i) = i$ for $i \in \{1, 2, 3\}$; $I_{\text{pos}}(\text{vacuum}) = 4$.	<code>ZGMmesInvariant.lean</code>
<code>zgm_mes_all_four_distinct</code>	All four orbital position values are pairwise distinct.	<code>ZGMmesInvariant.lean</code>
<code>zgm_mes_goe_source</code>	GoE property is structural source of position-1 uniqueness for gen_1 .	<code>ZGMmesInvariant.lean</code>
<code>zgm_mes_closed</code>	Master theorem: orbital chain discriminant canonically identifies all SM fermion generations (2-ZGM-MES closed $\mathbf{A}_{\text{Lean}}$).	<code>ZGMmesInvariant.lean</code>
<code>fmdl_generation_stability_ordering</code>	Predecessor count ordering.	<code>GoEStability-Hierarchy.lean</code>
<code>fmdl_gen1_stability_dominance</code>	Gen_1 is strictly minimal: unique state with zero predecessors.	<code>GoEStability-Hierarchy.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<i>Z₅ Transitivity Uniqueness</i> (<i>Z5TransitivityUniqueness.lean</i>):		
<code>z5_gen2_rotations_to_allones</code>	All 5 rotations of <code>smGen₂</code> reach all-ones under Rule 110 in 1 step.	<code>Z5TransitivityUniqueness.lean</code>
<code>z5_full_transitivity_smGen1</code>	<code>smGen₁</code> 's cyclic class has full Hamming-3 transitivity on the 5-ring.	<code>Z5TransitivityUniqueness.lean</code>
<code>z5_weight3_full_transitivity_count</code>	Count of full-transitive weight-3 classes on the 5-ring.	<code>Z5TransitivityUniqueness.lean</code>
<code>z5_transitivity_uniqueness</code>	$p = 5$ is the unique prime ≤ 23 with full Hamming-3 transitivity under Rule 110; total null result for primes $\{2, 3, 7, 11, 13, 17, 19, 23\}$.	<code>Z5TransitivityUniqueness.lean</code>
<code>rule110Ring / cyclicShiftRing / hammingWeightRing</code>	Reusable infrastructure for Rule 110 on p -rings (any prime p).	<code>Z5TransitivityUniqueness.lean</code>
<i>Z₅ Transitivity Uniqueness §8 (extended)</i> :		
<code>z5_w3_exclusive_among_primes</code>	Weight-3 full-transitivity exclusive to $p = 5$ across all $t \leq 4$; $p = 7$ has zero weight-3 transitivity.	<code>Z5TransitivityUniqueness.lean</code>
<code>z5_w3_t1_full_transitivity</code>	Second weight-3 class on $p = 5$ achieves full transitivity at $t = 1$.	<code>Z5TransitivityUniqueness.lean</code>
<code>p7_w4_t1_full_transitivity</code>	$p = 7$ has exactly one full-transitive class: weight-4, $t = 1$ (7 rotations of $[0, 1, 0, 1, 0, 1, 1]$).	<code>Z5TransitivityUniqueness.lean</code>
<i>GoE Stability Hierarchy §6–§9 (extended)</i> , (<i>GoEstabilityHierarchy.lean</i>):		
<code>fmdl_gen1_all_rotations_are_goe</code>	All 5 cyclic rotations of <code>gen₁</code> are GoE states (<code>native_decide</code> , $5 \times 16,807$ checks).	<code>GoEstabilityHierarchy.lean</code>
<code>fmdl_gen1_all_rotations_no_predecessor</code>	All 5 <code>gen₁</code> rotations have zero predecessors (<code>native_decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>sm_orbit_unique_gtp3</code>	Every GTP-3 start is a <code>gen₁</code> rotation (<code>native_decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>fmdl_max_gtp_length_is_3</code>	No GTP-4 chain exists; maximum chain length is 3 (<code>native_decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>sm_orbit_gtp3_count</code>	Exactly 5 GTP-3 chains exist (<code>native_decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>fmdl_ca_lepton_universality</code>	4-part master theorem: unique decay channels + equal cardinality + no generation mixing (<code>native_decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>fmdl_no_generation_shortcut</code>	No CA state can skip a generation in the orbit dynamics.	<code>GoEstabilityHierarchy.lean</code>
<code>gtp3_sum_trajectory_of_gen1_rotations</code>	All 5 <code>gen₁</code> rotations have Z_7 -sum trajectory $(4, 4, 3)$ (<code>decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>gtp3_alt_depth_is_two</code>	Alt-orbit rotations are depth-2 GTP-2 chains, not GTP-3 (<code>decide</code>).	<code>GoEstabilityHierarchy.lean</code>
<code>gtp3_sum_trajectory_master</code>	Joint summary: GTP-3 sum fingerprint $(4, 4, 3)$ + GTP-2 alt class (<code>decide</code>).	<code>GoEstabilityHierarchy.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<i>CUP3D Uniqueness §7a–§10 (extended, CUP3DUniqueness.lean):</i>		
fmdl_unique_fixed_point	Vacuum is the unique fixed point of fmdl_step5 (native_decide, 16,807 cases). No non-vacuum periodic orbits exist.	CUP3DUniqueness.lean
fmdl_no_nontrivial_cycles		CUP3DUniqueness.lean
fmdl_vacuum_is_unique_attractor	All states converge to vacuum within 7 steps (native_decide).	CUP3DUniqueness.lean
fmdl_orbit_depth_profile	Full depth histogram over all $7^5 = 16,807$ states (decide).	CUP3DUniqueness.lean
fmdl_universal_7step_convergence	All 7^5 states reach vacuum in ≤ 7 steps (native_decide).	CUP3DUniqueness.lean
fmdl_max_depth_is_7	Maximum decay depth = 7 (native_decide).	CUP3DUniqueness.lean
fmdl_depth7_witness_exact	Explicit witness of a depth-7 state (decide).	CUP3DUniqueness.lean
z7_conservation_count_table	Full Z_7 sum conservation counts by value (native_decide).	CUP3DUniqueness.lean
z7_sum4_uniquely_sparse	$v = 4$ is the unique rarest non-zero conservation class (native_decide).	CUP3DUniqueness.lean
z7_gen3_not_sum_conserving	Gen ₃ does not conserve Z_7 sum ($3 \rightarrow 0$, decide).	CUP3DUniqueness.lean
z7_vacuum_sum_conserved	Vacuum is sum-conserving (decide).	CUP3DUniqueness.lean
z7_conservation_landscape_summary	Complete conservation landscape summary (native_decide).	CUP3DUniqueness.lean
fmdl_nonzero_diagonal_all_zero	For all $k \in \{1, \dots, 6\}$: $f_{\text{MDL}}(k, k, k) = 0$.	CUP3DUniqueness.lean
fmdl_unique_uniform_fixed_point	$k = 0$ is the unique Z_7 value satisfying $f_{\text{MDL}}(k, k, k) = k$.	CUP3DUniqueness.lean
photon_is_ca_ether	Photon ($Z_7 = 0$) is the unique CA ether fixed point.	CUP3DUniqueness.lean
fmdl_uniform_fp_uniqueness_count	Count of uniform fixed points = 1.	CUP3DUniqueness.lean
apply_f_ring	Generalized Z_7 ring step (def: arbitrary Z_7 function).	CUP3DUniqueness.lean
is_orbit_admissible	Orbit-producing predicate for arbitrary Z_7 functions (def).	CUP3DUniqueness.lean
fmdl_is_orbit_admissible	f_{MDL} satisfies orbit-admissibility (decide).	CUP3DUniqueness.lean
orbit_sum_trajectory_invariant	Universal Z_7 -sum trajectory $4 \rightarrow 4 \rightarrow 3 \rightarrow 0$ for all orbit-admissible f (rw+decide).	CUP3DUniqueness.lean
orbit_sum_full_trajectory	4-step full trajectory summary.	CUP3DUniqueness.lean
fmdl_z5_equivariant	f_{MDL} is exactly Z_5 -equivariant (84,035 cases, native_decide).	CUP3DUniqueness.lean

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
z7_binary_not_ring_homomorphism	The mod-2 projection $\varphi : \mathbb{Z}_7 \rightarrow \mathbb{Z}_2$ is not a ring homomorphism.	CUP3DUniqueness.lean
z7_binary_not_ring_hom_universal	No ring homomorphism $\mathbb{Z}_7 \rightarrow \mathbb{Z}_2$ exists.	CUP3DUniqueness.lean
z7_z2_incompatible_additive	$\mathbb{Z}_7$ and $\mathbb{Z}_2$ have incompatible additive structure.	CUP3DUniqueness.lean
<i>Dimensional Slice Uniqueness §4b (DimensionalSliceUniqueness.lean):</i>		
vacuum_selects_rule110_over_rule111	4-part Finset identity: orbit rules = $\{110, 111\}$ exactly; vacuum selects $\{110\}$ (native_decide).	DimensionalSliceUniqueness.lean
rule111_orbit_valid_no_vacuum	Rule 111 satisfies the SM orbit but maps $(0, 0, 0) \rightarrow 1$: the unique near-miss (native_decide).	DimensionalSliceUniqueness.lean
<i>Z₇ Charge Conjugation §5–§6 (extended, Z7ChargeConjugation.lean):</i>		
sm_u_quark / sm_vacuum_cell / sm_w_plus / sm_w_minus / sm_d_quark / sm_electron	SM particle constant definitions in $\mathbb{Z}_7$.	Z7ChargeConjugation.lean
sm_charged_current_vertex	$(u_q, \gamma, u_q) \rightarrow W^+$ is the unique f_{MDL} configuration producing W^+ .	Z7ChargeConjugation.lean
sm_w_minus_absence	W^- emission is arithmetically forbidden (zero neighborhoods).	Z7ChargeConjugation.lean
sm_cp_vertex_asymmetry	The W^+/W^- asymmetry is maximal: 1 neighborhood vs. 0.	Z7ChargeConjugation.lean
fmdl_nonzero_count_14	Exactly 14 of 343 neighborhoods produce nonzero output (native_decide).	Z7ChargeConjugation.lean
fmdl_unique_mdl_cp_structure	f_{MDL} is the unique MDL-minimal function with $\mathbb{Z}_7 = 4$ exclusion (native_decide).	Z7ChargeConjugation.lean
fmdl_mdl_uniqueness	MDL-minimal completion uniqueness (funext+by_cases).	Z7ChargeConjugation.lean
fmdl_mdl_minimal_implies_z4_exclusion	MDL-minimality $\Rightarrow \mathbb{Z}_7 = 4$ exclusion (rewrite).	Z7ChargeConjugation.lean
fmdl_vacuum_transparent	Vacuum transparency alias (definitional unfolding).	Z7ChargeConjugation.lean
<i>CA Masslessness and Ether (CasimirMasslessEther.lean):</i>		
fmdl_massless_criterion	$f_{\text{MDL}}(0, k, 0) = k \iff k \in \{0, 1\}$: CA-massless partition (native_decide).	CasimirMasslessEther.lean
fmdl_massive_decay	$\forall k \in \{2, \dots, 6\}: f_{\text{MDL}}(0, k, 0) = 0$ (massive decay to vacuum).	CasimirMasslessEther.lean
u_photon_u_to_W_vertex	$f_{\text{MDL}}(2, 0, 2) = 3$: the CA $(u, \gamma, u) \rightarrow W^+$ vertex (native_decide).	CasimirMasslessEther.lean

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>nu_photon_nu_absorption</code>	$f_{\text{MDL}}(1, 0, 1) = 1$: neutrino pair photon absorption (<code>native_decide</code>).	<code>CasimirMassless-Ether.lean</code>
<code>photon_absorption_events</code>	Exactly 2 non-trivial virtual-photon interactions among all 36 matter-pair configurations.	<code>CasimirMassless-Ether.lean</code>
<code>ether_z7_sum_mod7</code>	Rule 110 ether (period-14 background) carries Z_7 winding 1 (neutrino sector, not EM vacuum); Z_7 sum mod 7 = 1 (<code>native_decide</code>).	<code>CasimirMassless-Ether.lean</code>
<code>ether_not_em_vacuum</code>	The ether background is the neutrino-sector medium, not the EM vacuum; corrects two-level model labeling.	<code>CasimirMassless-Ether.lean</code>
<code>transverse_sector_invariance</code>	$\forall r$ (BinaryCARule) with $r(0, 0, 0) = 0$: iterating r on the all-zero $\mathbb{Z}_2$ state preserves it indefinitely — the CA-level γ/Z discriminant is universal and rule-independent (<code>trivial+decide, A_{Lean}</code>).	<code>CasimirMassless-Ether.lean</code>
<i>P22 Vertex Table (Z7ChargeConjugation.lean §7):</i>		
<code>p22_absorption_vertices_are_transparent</code>	$f_{\text{MDL}}(6, 3, 2) = f_{\text{MDL}}(4, 3, 0) = f_{\text{MDL}}(2, 4, 6) = f_{\text{MDL}}(0, 4, 4) = 0$ (<code>by decide, A_{Lean}</code>).	<code>Z7Charge-Conjugation.lean</code>
<code>ca_w_plus_is_emission_not_absorption</code>	$f_{\text{MDL}}(2, 0, 2) = 3$ (emission) $\wedge f_{\text{MDL}}(6, 3, 2) = 0$ (transparency): the W^+ neighborhood is emission-only (<code>by decide, A_{Lean}</code>).	<code>Z7Charge-Conjugation.lean</code>
<code>p22_vertex_table_is_ca_transparent</code>	Master theorem: all four P22 charged-current absorption vertices map to $f_{\text{MDL}} = 0$ (<code>by decide, A_{Lean}</code>).	<code>Z7Charge-Conjugation.lean</code>
<i>SM Orbit Causal Isolation (SMOrbitCausalIsolation.lean):</i>		
<code>sm_orbit_complete_causal_isolation</code>	6-part master theorem: GoE + unique predecessor chain + Z_7 sum conservation + global isolation. Zero <code>sorry</code> .	<code>SMOrbitCausalIsolation.lean</code>
<i>GTE Uniqueness §5 (GTEUniqueness.lean):</i>		
<code>minimal_lawful_has_card_one</code>	The set of minimal lawful UWCA programs has cardinality one.	<code>GTEUniqueness.lean</code>
<code>UWCAIsomorphic</code>	Definition of UWCA isomorphism.	<code>GTEUniqueness.lean</code>
<code>gte_uniqueness_isomorphism</code>	Any minimal lawful UWCA program is isomorphic to Σ_{GTE} (zero <code>sorry</code>).	<code>GTEUniqueness.lean</code>
<code>gte_isomorphism_symmetric</code>	The UWCA isomorphism relation is symmetric.	<code>GTEUniqueness.lean</code>
<i>GUT Structure §9+§13+§16+§17 (GUTStructure.lean):</i>		
<code>fmdl_count_eq_chiggs_plus_one</code>	f_{MDL} nonzero count = $c_H + 1 = 14$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>fmdl_count_decomposition</code>	$14 = N_{\text{gen}} + 2N_{\text{fam}} + 1 = 3 + 10 + 1$.	<code>GUTStructure.lean</code>
<code>fmdl_count_ngen_nfam</code>	Count in terms of structural parameters.	<code>GUTStructure.lean</code>
<code>b_higgs_eq_ngen</code>	$b_H = N_{\text{gen}} = 3$ (Higgs branch winding = generation count).	<code>GUTStructure.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>gte_family_capacity_identity</code>	$N_{\text{gen}} + N_{\text{fam}} = 2^{N_{\text{gen}}} = 8$ (<code>norm_num</code> , alias of <code>ngen_plus_nfam_eq_pow2</code>).	<code>GUTStructure.lean</code>
<code>running_shift_is_z5_ring</code>	Denominator running shift $c_H - 2^{N_{\text{gen}}} = N_{\text{fam}} = 5$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>z5_ring_contributes_nfam_to_denominator</code>	$c_H = 2^{N_{\text{gen}}} + N_{\text{fam}}$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>b_sum_is_product</code>	$b_{\text{sum}} = 2N_{\text{gen}}N_{\text{fam}}c_H = 390$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>weinberg_numerator_in_bsum</code>	$N_{\text{gen}} \mid b_{\text{sum}}$.	<code>GUTStructure.lean</code>
<code>weinberg_denominator_in_bsum</code>	$c_H \mid b_{\text{sum}}$.	<code>GUTStructure.lean</code>
<code>weinberg_ratio_from_bsum</code>	$(N_{\text{gen}} : \mathbb{Q})/c_H = 3/13$ from b_{sum} factorization.	<code>GUTStructure.lean</code>
<code>nw_plus_chiggs_eq_pow2</code>	$N_{\text{gen}} + c_H = 16 = 2^4$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>rule124_minterms_card</code>	Rule 124 MDL one-count = 5 (<code>native_decide</code>).	<code>GUTStructure.lean</code>
<code>rule110_and_124_joint_md1_count</code>	Rules 110 and 124 both achieve MDL one-count = 5 (<code>native_decide</code>).	<code>GUTStructure.lean</code>
<code>rule110_preferred_by_sublayer_consistency</code>	Rule 110 uniquely preferred over Rule 124 by sublayer consistency (<code>native_decide</code>).	<code>GUTStructure.lean</code>
<i>EW Boson Structure §5 (EWBosonStructure.lean):</i>		
<code>d_higgs</code>	Definition: Higgs Goldstone count $d_{H^0} = 0$.	<code>EWBosonStructure.lean</code>
<code>d_z</code>	Definition: Z-boson Goldstone count $d_Z = 1$.	<code>EWBosonStructure.lean</code>
<code>d_w</code>	Definition: $W^\pm$ -boson Goldstone count $d_{W^\pm} = 2$.	<code>EWBosonStructure.lean</code>
<code>goldstone_cascade_higgs</code>	$c_{H^0} = c_H - d_{H^0} = 13$ (<code>norm_num</code> , zero sorry).	<code>EWBosonStructure.lean</code>
<code>goldstone_cascade_z</code>	$c_Z = c_H - d_Z = 12$ (<code>norm_num</code> , zero sorry).	<code>EWBosonStructure.lean</code>
<code>goldstone_cascade_w</code>	$c_{W^\pm} = c_H - d_{W^\pm} = 11$ (<code>norm_num</code> , zero sorry).	<code>EWBosonStructure.lean</code>
<code>goldstone_cascade_formula</code>	General formula $c_P = c_H - d_P$ (<code>norm_num</code> , zero sorry).	<code>EWBosonStructure.lean</code>
<i>GUTStructure §23: GTE Master Formula (GUTStructure.lean):</i>		
<code>gte_master_formula_complete</code>	All four SM EW parameters from $N_{\text{gen}} = 3$ alone: arithmetic bridge, $\sin^2\theta_W(M_{\text{GUT}}) = 3/8$, $\sin^2\theta_W(M_Z) = 3/13$, $\lambda = 9/40$. Zero free parameters (<code>norm_num</code> , <code>A_Lean</code> , capstone).	<code>GUTStructure.lean</code>
<code>gte_generating_triple</code>	$N_{\text{fam}} = 2^{N_{\text{gen}}} - N_{\text{gen}}$, $c_H = 2^{N_{\text{gen}}+1} - N_{\text{gen}}$, $c_H = N_{\text{gen}} + 2N_{\text{fam}}$ — three equivalent cascade forms (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>gte_master_formula_gut_weinberg</code>	$\sin^2\theta_W(M_{\text{GUT}}) = N_{\text{gen}}/2^{N_{\text{gen}}} = 3/8$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>gte_master_formula_ew_weinberg</code>	$\sin^2\theta_W(M_Z) = N_{\text{gen}}/c_H = 3/13$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>gte_master_formula_wolfenstein</code>	$\lambda = N_{\text{gen}}^2/(2^{N_{\text{gen}}} \cdot N_{\text{fam}}) = 9/40$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>gte_master_formula_rb</code>	$R_b = N_{\text{gen}}/2^{N_{\text{gen}}} = 3/8 = \sin^2\theta_W(M_{\text{GUT}})$ (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>gte_cross_sector_bridge</code>	$\sin^2\theta_W(M_{\text{GUT}}) = R_b$ and $\lambda = (N_{\text{gen}}/N_{\text{fam}}) \cdot 3/8 = 9/40$ — three SM quantities share one arithmetic root (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>gte_arithmetic_root</code>	$N_{\text{gen}} + N_{\text{fam}} = 2^{N_{\text{gen}}}$ — the single arithmetic root of all four SM EW parameters (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<code>ngen_3_mersenne_uniqueness</code>	$2^{N_{\text{fam}}} - 1 = 31$ prime and $2^{c_H} - 1 = 8191$ prime: double Mersenne endpoint uniqueness (<code>norm_num</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §27: Bidirectional Unification (GUTStructure.lean):</i>		
<code>ugp_arith_forces_rule110</code>	Arrow A1: SM orbit uniquely selects Rule 110 (alias of <code>cup1_orbit_uniquely_selects_rule110</code> , $\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>sm_selects_gte_triple</code>	Arrow A2-R: $N_{\text{gen}} = 3$, $N_{\text{fam}} = 5$, $c_H = 13$ by <code>rf1</code> ($\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>gte_predicts_ew_mixing</code>	Arrow A2: $\sin^2\theta_W(M_Z) = 3/13$ and $\sin^2\theta_W(M_{\text{GUT}}) = 3/8$ ($\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>gte_predicts_ckm_lambda</code>	Arrow A2: $\lambda = 9/40$ (alias of <code>wolfenstein_lambda_formula</code> , $\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>rule110_encodes_sm_particles</code>	Arrow A3: photon = unique CA fixed point $\wedge$ gen_1 GoE $\wedge$ $Z_7=4$ excluded ($\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>ugp_r110_sm_joint_unification</code>	Capstone: 7-conjunct bidirectional unification — arithmetic bridge, EW and GUT Weinberg angles, CKM λ , double Mersenne endpoint, photon CA fixed point, gen_1 GoE. Zero sorry, zero free parameters ($\mathbf{A}_{\text{Lean}}$, capstone).	<code>GUTStructure.lean</code>
<i>GUTStructure §28: MDL Robustness and Z_7 Free Minterm Count (GUTStructure.lean):</i>		
<code>z7_fixed_neighborhood_count</code>	Exactly 18 of 343 $\mathbb{Z}_7^3$ neighborhoods satisfy <code>isFixedNeighborhood</code> (10 orbit + 8 binary, <code>native_decide</code>).	<code>GUTStructure.lean</code>
<code>z7_free_neighborhood_count</code>	$325 = 343 - 18$ free neighborhoods; MDL sets all to 0 (<code>native_decide</code>).	<code>GUTStructure.lean</code>
<code>mdl_robustness_z7</code>	Any orbit-admissible MDL-minimal $\mathbb{Z}_7$ CA function = f_{MDL} — uniqueness independent of orbit depth; delegates to <code>fmdl_mdl_uniqueness</code> .	<code>GUTStructure.lean</code>
<i>GUTStructure §29: Z_2 Longitudinal Universality + Chern-Simons Level (GUTStructure.lean):</i>		
<code>z_boson_cvalue_equals_mdl_plus_z7</code>	$c_Z = 7 + \text{MDL}(\text{Rule 110}) = 7 + 5 = 12$ (<code>norm_num</code> , $\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<code>z_boson_mdl_class4_chain</code>	3-conjunct structural chain: minterm count = 5; $5 + 7 = c_Z = 12$; $c_Z = c_H - 1$ (<code>norm_num</code> , $\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>cs_level_all_formulas</code>	$SU(2)_k$ Chern-Simons level $k = 30$ from four independent GTE formulas: $2^{N_{\text{fam}}} - 2$, $N_{\text{gen}}! \cdot N_{\text{fam}}$, $4(N_{\text{gen}}^2 - 1) - 2$, $2(c_H + 2)$ (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §30: Mersenne Cascade Discriminator</i> (<code>GUTStructure.lean</code>):		
<code>bt_is_composite</code>	$b_t = 337920$ is not prime; $337920 = 2^{11} \times 3 \times 5 \times 11$ (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<code>bb_mersenne_bt_not</code>	$b_b = 8191$ is Mersenne prime; b_t is not prime — combined discriminator (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<code>quark_cascade_mersenne_discriminator</code>	Joint theorem: Mersenne discriminator selects canonical quark G1 seed pair ($b_{L3} = 275$, $b_{L2} = 42$) (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §31: Primordial $T(2,3)$ Topology — Cascade Period $p = 2$</i> (<code>GUTStructure.lean</code>):		
<code>cascade_period_coprimality</code>	$\gcd(2, N_{\text{gen}}) = 1$: $T(2,3)$ is a knot, not a torus link (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<code>cascade_period_3_fails_coprimality</code>	$\gcd(3, N_{\text{gen}}) \neq 1$: $T(3,3)$ is a 3-component torus link; $p = 3$ excluded (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<code>cascade_period_minimum_is_two</code>	Three-constraint joint selection: GoE necessity, PSC connectedness, MDL minimality uniquely force $p = 2$ (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §32: Vacuum Ollivier-Ricci Flatness</i> (<code>GUTStructure.lean</code>):		
<code>vacuum_ollivier_ricci_flatness</code>	$\kappa_{EE} = 0$: Rule 110 CA vacuum fixed point gives Ricci-flat ether background in deviation-based Ollivier-Ricci metric; upgrades the Gorard numerical observation (<code>A</code>) to a Lean-certified arithmetic theorem (<code>decide</code> , zero <code>sorry</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §33: $SU(2)_L$ Charge Assignment from Z_7 Color Structure</i> (<code>GUTStructure.lean</code>):		
<code>su2l_charge_assignment_z7_discriminator</code>	Z_7 -alignment discriminates canonical quark assignment: $N_{\text{gen}}^2 \bmod 7 = 2 = w(u)$ (up-type $\rightarrow N_{\text{gen}}^2 = 9$); $N_{\text{fam}} \bmod 7 = 5 \neq 2$ (down-type $\rightarrow N_{\text{fam}} = 5$). Charge-swap candidate excluded (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §34: Full 6-Quark N_{eff} Capstone</i> (<code>GUTStructure.lean</code>):		
<code>six_quark_neff_complete</code>	6-conjunct joint theorem: all six GTE quark N_{eff} structural formulas ($u = N_{\text{gen}}^2$, $d = N_{\text{fam}}$, $c = N_{\text{fam}}^2(2N_{\text{fam}} + 1)$, $s = 2N_{\text{gen}}(2c_H + N_{\text{fam}})$, $b = 2^{c_H} - 1$, $t = 2^{c_W} \cdot N_{\text{gen}} \cdot N_{\text{fam}} \cdot (2N_{\text{fam}} + 1)$) (<code>norm_num</code> , <code>A_Lean</code> , <code>capstone</code>).	<code>GUTStructure.lean</code>
<code>quark_neff_all_distinct</code>	All six quark N_{eff} values are pairwise distinct: spectral non-degeneracy (<code>norm_num</code> , <code>A_Lean</code>).	<code>GUTStructure.lean</code>
<i>GUTStructure §35: Dark Sector Period-2 Orbits — Rule 110 on 4-Cell Binary Ring</i> (<code>GUTStructure.lean</code>):		

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
dark_sector_vacuum_fixed_point	(0, 0, 0, 0) is the unique fixed point of Rule 110 on the 4-cell binary periodic ring (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
dark_sector_cycles_are_period2	All four dark-cycle states $\{(1, 1, 1, 0), (1, 0, 1, 1), (1, 1, 0, 1), (0, 1, 1, 1)\}$ are period-2 (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
dark_sector_period2_exhaustive	The four dark-cycle states are exactly the period-2 orbits of the 4-cell ring; exhaustive classification (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
<i>GUTStructure §44: Orbit-Closure and Vertex Uniqueness (GUTStructure.lean):</i>		
orbit_winding_set	The orbit-winding set $\{0, 1, 2, 5\}$ is the exactly the set of windings appearing in the SM generation orbit and background (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
self_propagating_iff_orbit_winding	$\forall w \in \mathbb{Z}_7$: w is self-propagating under f_{MDL} iff $w \in \{0, 1, 2, 5\}$ (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
orbit_closure_theorem	The filter of self-propagating windings equals the orbit-closure $\{0, 1, 2, 5\}$; establishes the orbit-closure biconditional ($\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
dquark_unique_production_vertex	(2, 5, 2) is the unique neighborhood producing $Z_7 = 6$ (d-quark) under f_{MDL} (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
dquark_unique_preimage	$f_{\text{MDL}}(l, c, r) = 6 \iff (l = 2 \wedge c = 5 \wedge r = 2)$; uniqueness biconditional (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
dquark_preimage_count	The preimage set of 6 under f_{MDL} has cardinality exactly 1 (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
<i>GUTStructure §46: Structural Lepton Number Conservation (GUTStructure.lean):</i>		
lepton_production_impossible	$4 \notin \text{range}(f_{\text{MDL}})$: the CA substrate cannot produce winding-4 cells from any input triple ($\mathbf{A}_{\text{Lean}}$, zero sorry).	GUTStructure.lean
lepton_number_conservation	Structural lepton conservation: any tape with no winding-4 cells develops none under f_{MDL} dynamics for any number of steps ($\mathbf{A}_{\text{Lean}}$).	GUTStructure.lean
lepton_number_preserved_from_L0_initial	Multi-step preservation of winding-4 absence from any winding-4-free initial configuration ($\mathbf{A}_{\text{Lean}}$, zero sorry).	GUTStructure.lean
lepton_number_conservation_summary	Master summary theorem: structural lepton non-creation at CA level, combining all §46 results ($\mathbf{A}_{\text{Lean}}$, zero sorry).	GUTStructure.lean
<i>GUTStructure §39: W^+ Decay Lemma (GUTStructure.lean):</i>		
wplus_center_maps_to_vacuum	$\forall (l, r) \in \mathbb{Z}_7^2$: $f_{\text{MDL}}(l, 3, r) = 0$ — a W^+ center cell always maps to vacuum (<code>decide</code> , $\mathbf{A}_{\text{Lean}}$, zero sorry).	GUTStructure.lean
wplus_creation_then_decay	The W^+ is created in one CA step by $(2, 0, 2) \rightarrow 3$ and decays to vacuum the next step; W^+ lifetime is exactly 1 timestep (<code>decide</code> , $\mathbf{A}_{\text{Lean}}$, zero sorry).	GUTStructure.lean
<i>GUTStructure §40: Mersenne Structural Identities — $c_H = p_{N_{\text{fam}}}(M)$ and $T_{\text{ether}} = c_H + 1$ (GUTStructure.lean):</i>		

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
<code>ch_eq_mersenne_exp_nfam</code>	$c_H = 13$ equals the N_{fam} -th Mersenne prime exponent: the orbit formula $c_H = N_{\text{gen}} + 2N_{\text{fam}}$ forces this position, so $b_b = 2^{c_H} - 1 = M_{13}$ is a Mersenne prime by arithmetic necessity (<code>native_decide/norm_num</code> , $\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>ether_period_ch_structural</code>	$T_{\text{ether}} = c_H + 1 = 14$: ether period equals Higgs cascade depth plus one (<code>native_decide</code> , $\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<i>GUTStructure §43: Universal Mersenne Partition (GUTStructure.lean):</i>		
<code>universal_mersenne_partition</code>	At cascade ridge $n = 10$: seed $c \in \{823, 2137\}$ are non-Mersenne primes; T_{odd} output $c = 1023 = 2^{10} - 1$ and T_{even} output $c = 65535 = 2^{16} - 1$ are Mersenne. MDL-lex-min selects $c = 823$. (11 theorems, zero sorry, $\mathbf{A}_{\text{Lean}}$).	<code>GUTStructure.lean</code>
<i>GUTStructure §47: Ward Identity Algebraic Core (GUTStructure.lean):</i>		
<code>ward_mass_cancellation</code>	$\text{Im}(\psi_R^* \psi_L) + \text{Im}(\psi_L^* \psi_R) = 0$ for all $\psi_R, \psi_L \in \mathbb{C}$: the mass cross-term cancellation, the key non-trivial step in the Z_7 winding current continuity equation (<code>decide</code> , $\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>ward_chirality_density</code>	$\psi^\dagger \sigma_z \psi = \text{normSq}(\psi_R) - \text{normSq}(\psi_L)$: chirality density as difference of layer probability densities ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>ward_identity_algebraic_summary</code>	Master summary: algebraic core of $\partial_t(\psi^\dagger \psi) + \partial_x(v_{\text{CA}} \psi^\dagger \sigma_z \psi) = 0$ certified; full continuum identity requires product-rule axiom ($\mathbf{A}_{\text{Lean}}$ core, $\mathbf{A}/\mathbf{D}$ full, zero sorry).	<code>GUTStructure.lean</code>
<i>GUTStructure §59: Gen₁ GoE Uniqueness and Baryogenesis Exclusivity (GUTStructure.lean):</i>		
<code>gen2_has_predecessor</code>	$\exists s, f_{\text{MDL-step5}}(s) = \text{gen}_2$: gen_2 has at least one predecessor under f_{MDL} on the $\mathbb{Z}_5$ ring ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>gen3_has_predecessor</code>	$\exists s, f_{\text{MDL-step5}}(s) = \text{gen}_3$: gen_3 has at least one predecessor under f_{MDL} on the $\mathbb{Z}_5$ ring ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>gen1_unique_goe_in_orbit</code>	Full structural uniqueness: among $\{\text{gen}_1, \text{gen}_2, \text{gen}_3\}$, gen_1 has no predecessor; gen_2 and gen_3 each have at least one. Gen_1 is the unique Garden of Eden state in the generation orbit ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<code>baryogenesis_exclusivity</code>	$N_{\text{fam}} - 1 = 4 \neq N_{\text{fam}} = 5$: the GoE ring-cut count differs from the non-GoE ring-close count, establishing the arithmetic basis for gen_1 baryogenesis exclusivity ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>GUTStructure.lean</code>
<i>Trajectory-Orbit No-Go (UgpLean.Universality.CouplingNoGo):</i>		
<code>coupling_no_go_master</code> + 9 supporting theorems	Trajectory-orbit mismatch: $\text{gcd}(2, 14) = 2$, ether orbit length = 7, $\text{gcd}(3, 7) = 1$, $\text{lcm}(3, 7) = 21$, $21 \nmid 3$; diagonal offsets provide no escape. All CA-local couplings between layers fail to preserve C_2 glider coherence ($\mathbf{A}_{\text{Lean}}$, zero sorry).	<code>CouplingNoGo.lean</code>
<i>CA-Level Chirality and Parity (UgpLean.Universality.ChiralityEigenstates):</i>		

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
fmdl_p_violation_count	$\#\{(l, c, r) \in \mathbb{Z}_7^3 : f_{\text{MDL}}(l, c, r) \neq f_{\text{MDL}}(r, c, l)\} = 14$: the parity-violation count of f_{MDL} is exactly 14 ($\mathbf{A}_{\text{Lean}}$, <code>native_decide</code> , zero sorry).	ChiralityEigenstates .lean
gen1_is_chiral	$\text{gen}_1 \neq P(\text{gen}_1)$: the first-generation vector is not fixed by spatial parity ($\mathbf{A}_{\text{Lean}}$, <code>decide</code> , zero sorry).	ChiralityEigenstates .lean
p_gen1_two_step_decay	$f_{\text{MDL}}^2(P(\text{gen}_1)) = \text{vac}$: the parity mirror of gen_1 decays to vacuum in 2 steps (vs. 3 steps for the physical orbit) ($\mathbf{A}_{\text{Lean}}$, <code>native_decide</code> , zero sorry).	ChiralityEigenstates .lean
gen3_p_covariant	$f_{\text{MDL}}(P(\text{gen}_3)) = P(f_{\text{MDL}}(\text{gen}_3))$: gen_3 is parity-covariant — the orbit endpoint is chirally neutral ($\mathbf{A}_{\text{Lean}}$, <code>native_decide</code> , zero sorry).	ChiralityEigenstates .lean
sm_orbit_is_left_handed	Combined theorem: all four chirality properties simultaneously certified ($\mathbf{A}_{\text{Lean}}$, zero sorry, zero named axioms).	ChiralityEigenstates .lean
<i>Spacetime / QFT modules (§10):</i>		
LiftingTheorem.lean	<code>algebraic_lifting_theorem</code> , <code>charge_quantization_physical</code> , <code>three_generations_physical</code> , <code>no_fourth_generation_physical</code> , <code>scattering_existence</code> , <code>smatrix_framework_exists</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
SpatiallyExtendedLifting.lean	<code>spatially_extended_composite_lifting</code> , <code>meson_bound_state_exists</code> , <code>baryon_bound_state_exists</code> , <code>causal_path_exists</code> (theorem, proved for forward-causal pairs; zero custom axioms in any $\mathbf{A}_{\text{Lean}}$ proof chain) ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
ColorConfinement.lean	<code>no_psc_admissible_single_quark</code> , <code>psc_rc_requires_system_color_neutral</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
MassGap.lean	<code>gte_mass_gap</code> , <code>mass_hierarchy_gen3_gt_gen2_gt_gen1</code> ($\mathbf{A}_{\text{Lean}}$).	Spacetime/
QFT/GaugedMassGap.lean	<code>qft_gauged_mass_gap_unconditional</code> , <code>qft_gauged_mass_gap_pos_unconditional</code> , <code>confined_massive_color_singlet</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	QFT/
AnomalyRenormalizability.lean	<code>renormalizability_psc_forced</code> , <code>sm_lagrangian_renormalizable</code> , <code>all_generation_charge_neutrality_bundle</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
PhysicalExclusion.lean	<code>unified_physical_exclusion</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
ThreeGenerationCapstone.lean	<code>gte_uniquely_predicts_three_generations</code> : C1 record-equivalence uniqueness $\wedge$ period-3 orbit $\text{gen}_1 \rightarrow \text{gen}_2 \rightarrow \text{gen}_3 \wedge$ no fourth generation at physical scale ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Spacetime/
AlgebraicDescentTheorem.lean	<code>algebraic_descent_theorem</code> , <code>asymptotic_freedom_m_independent</code> , <code>psc_admissibility_no_resolution_parameter</code> ($\mathbf{A}_{\text{Lean}}$, zero sorry).	Universality/

(continued on next page)

(Table 9 continued)

Theorem (Lean name)	Statement	File
SylowIndexCoupling Hierarchy.lean	F_{21} substrate: frobenius_f21_order, f21_substrate_beta_coefficient, f21_two_loop_beta_coefficient, strong_cp_theta_zero_f21, hadron multiplet theorems; rank136_vcoup_uniqueness ($V_{\text{coupling}} = \varepsilon \varphi ^2(D_\mu\chi)^2$ unique at dim-4, $\mathbf{A}_{\text{Lean}}$); rank137_epsder_closure ($\varepsilon = N_7/N_3^2 = 7/9$, $\mathbf{A}_{\text{Lean}}$, zero sorry).	Universality/
GaugeInvariance.lean	$\mathbb{Z}_3$ gauge invariance of Φ_{MDL} coupling: z3_covariant_deriv_invariant ($D_\mu\chi$ invariant under $\chi \rightarrow \chi + \alpha$, $A_\mu \rightarrow A_\mu + \partial_\mu\alpha$), phimdl_coupling_gauge_invariant ($\varepsilon \varphi ^2(D_\mu\chi)^2$ gauge-invariant), gte_seven_vertices_z7_winding (all 7 GTE vertex types satisfy $\mathbb{Z}_7$ winding conservation under $\mathbb{Z}_3$ gauge), dmu_coefficient_forced_neg_one (coefficient -1 in $D_\mu\chi$ algebraically forced), phimdl_z3_gauge_invariant_bundle (master bundle, $\mathbf{A}_{\text{Lean}}$, zero sorry).	Universality/

B Statistical Significance Derivation

B.1 Setup

The null hypothesis H_0 : the parity bit patterns ($\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3$) are random binary strings drawn uniformly. Under H_0 there is no structural connection between GTE quantum numbers and Rule 110.

B.2 Derivation details

Raw false-positive rate: Empirical null test over $N = 10,000$ random GTE-like experiments. Each experiment generates 15 random GTE-like triples and tests all 120 family orderings for Rule 110 commutativity. Result: $p_{\text{raw}} = 1.36\%$ (136/10000), far below the naïve union bound 11.7%, reflecting strong negative correlation among orderings. This is the primary significance bound: the probability that a randomly generated parity matrix satisfies the SM orbit for any rule-ordering pair.

Algebraic uniqueness (CUP-4, primary result): Exhaustive check over all 256×120 (rule, ordering) pairs establishes that exactly one vacuum-transparent rule satisfies the specific SM orbit: Rule 110. This is an exact algebraic constraint (Lean-certified), not a probabilistic estimate. The pre-commitment to Rule 110 via the UWCA bisimulation theorem removes post-hoc bias.

Structural reinforcement: Only 4 of 256 rules achieve any winning orderings ($4/256 \approx 1.6\%$). Only 5 of 32 possible gen_1 vectors reach $\text{gen}_3 = [1, 1, 1, 1, 1]$ under Rule 110 (Theorem 3.1; $5/32 = 15.6\%$). These factors are correlated with p_{raw} (they are not independent multipliers) but each independently confirms the non-triviality of the CUP-4 result. Combining them multiplicatively would double-count and overstate significance; they are presented here as separate observational reinforcements.

C Artifact Manifest

Table 10: Computational artifacts for this paper.

Artifact	Description	Location
t_null_cup4.py	CUP-4 null test (5-part structural significance)	canonical_run/
t_null_cup4_results.json	Null test full results	canonical_run/
t_orbit_survey.py	Orbit survey (all 120 orderings, rule search)	canonical_run/
t_perturbed_orbits.py	Perturbed orbit test (Table 3)	canonical_run/
t_analytical_verification.py	Analytical verification of minterm structure	canonical_run/
CUP4Total-Parity.lean	Lean proofs of CUP-4, CUP-8, CUP-9	ugp-lean/
CUP11ModSeven.lean	Lean proof of CUP-11c	ugp-lean/
CUP3D-Uniqueness.lean	Lean proofs of f_{MDL} structural theorems	ugp-lean/
t_cup12_md1_minimal.py	CUP-12 MDL analysis (f_{MDL} fixed neighborhoods)	canonical_run/
t_cup12_md1_minimal_results.json	CUP-12 MDL full results	canonical_run/
t_cup12_cross_sector.py	CUP-12 cross-sector analysis (SM + dark; 76-bit description)	canonical_run/
t_cup12_cross_sector_results.json	CUP-12 cross-sector full results	canonical_run/
z5_transitivity_check.py	Python sweep of all primes ≤ 23 confirming Z_5 transitivity uniqueness at Hamming weight 3 (run time $< 1\text{s}$)	canonical_run/
<i>Additional scripts (graduated to canonical_run/):</i>		
orbit_admissible_count.py	Enumeration of SM orbit neighborhood constraints (23 fixed, 320 free); orbit-admissible count 7^{320} .	canonical_run/
z7_conservation_landscape.py	Complete Z_7 sum conservation sweep over all 7^5 states for all 7 sum values. Runtime 239ms.	canonical_run/

(continued on next page)

(Table 10 continued)

Artifact	Description	Location
<code>gtp_chain_uniqueness.py</code>	GTP- n exhaustive search: predecessor map, GoE-rooted chains, 5 GTP-3, 0 GTP-4+. Runtime 132ms.	<code>canonical_run/</code>
<code>gtp3_sum_trajectory.py</code>	GTP-3 chain enumeration with $\mathbb{Z}_7$ -sum trajectory verification; includes Rank 40 invariance validation.	<code>canonical_run/</code>
<code>gtp3_sum_trajectory_results.json</code>	GTP-3 sum trajectory frozen outputs.	<code>canonical_run/</code>
<code>fmdl_decay_depth.py</code>	Decay depth for all $7^5 = 16,807$ states under <code>fmdl_step5</code> : depth distribution, max depth 7.	<code>canonical_run/</code>
<code>transitivity_spectrum.py</code>	Full (p, w, t) transitivity spectrum for $p \leq 11$, $w \in \{1, 2, 3, 4\}$, $t \in \{1, 2, 3, 4\}$. Runtime ~ 0.2 s.	<code>canonical_run/</code>
<code>photon_vacuum_casimir_analysis.py</code>	CA zero-point entropy (vacuum = 0.000 bits/step), random-IC entropy (0.647 bits/cell/step).	<code>canonical_run/</code>
<code>photon_vacuum_casimir_results.json</code>	Photon vacuum and Casimir analysis frozen outputs.	<code>canonical_run/</code>
<code>ranks_46_50_casimir_items.py</code>	CA masslessness criterion verification; virtual-photon transparency survey (34/36 = 94.44%).	<code>canonical_run/</code>
<code>ranks_46_50_casimir_results.json</code>	Casimir items frozen outputs.	<code>canonical_run/</code>
<code>z3_z7_color_extension.py</code>	$\mathbb{Z}_3 \times \mathbb{Z}_7$ orbit construction and MDL analysis; $21^3 = 9261$ neighborhoods; 53 fixed; 9208 free; quark/lepton discrimination; color confinement cascade. Runtime: 88ms.	<code>canonical_run/</code>
<code>z3_z7_color_extension_results.json</code>	Color extension analysis frozen outputs.	<code>canonical_run/</code>
<code>z2_longitudinal_extension.py</code>	Binary CA rule enumeration (256 rules); propagation analysis; MDL ranking; γ/Z trajectory simulation (Rules 90 and 110 on 7-cell ring, 5 steps). Runtime: 98ms.	<code>canonical_run/</code>
<code>z2_longitudinal_extension_results.json</code>	$\mathbb{Z}_2$ longitudinal extension frozen outputs.	<code>canonical_run/</code>

(continued on next page)

(Table 10 continued)

Artifact	Description	Location
z7_output_distribution.py	f_MDL output distribution for all 7^3 inputs; preimage counts and conjugate-pair asymmetry.	canonical_run/
z7_output_distribution_results.json	Output distribution frozen results.	canonical_run/
fmdl3d_chirality.py	P-violation count (14/343), SM orbit chirality test, 3D systematic parity test (100 ICs). Runtime: 0.08s.	canonical_run/
fmdl3d_chirality_results.json	Chirality analysis frozen outputs.	canonical_run/
gte_triple_neutral_discrimination.py	GTE triple pairwise distinctness for all $\mathbb{Z}_7=0$ SM particles; discriminant function.	canonical_run/
complex_z7_rule110.py	Complex $\mathbb{Z}_7$ embedding via ω^k ; bifurcation analysis; ν vs $\{\gamma, Z\}$ amplitude flows.	canonical_run/
complex_z7_rule110_results.json	Complex $\mathbb{Z}_7$ analysis frozen outputs.	canonical_run/
rule110_period3_glider.py	Period-3 C_2 glider identification; $v_R = 2/3$ velocity measurement.	canonical_run/
<i>CA vertex and gauge scripts (graduated to scripts/):</i>		
ca_vertex_table.py	Complete CA vertex table: 343-entry enumeration, 14-neighborhood classification, P22 absorption vertex transparency check, completeness analysis.	scripts/
z7_gauge_invariance_check.py	$\mathbb{Z}_7$ gauge invariance tests: additive shift equivariance (FAILS: 2030/2058); $\mathbb{Z}_5$ rotation equivariance (PASSES). Runtime: 231ms.	scripts/
scripts/gauged_vertex_catalog.py	Vertex catalog under $\mathbb{Z}_3$ gauge: 7/7 GTE vertices recovered, 0 spurious (see REPRODUCE.md).	—
papers/33_deeper_consequences/scripts/decay_rates_from_gte.py	Tree-level GTE decay rates: $\Gamma_\mu +12.2\%$, $\Gamma_\tau +12.5\%$ vs PDG (see REPRODUCE.md).	—

D Reproducibility

All results in this paper are fully reproducible from the scripts in `canonical_run/`.

Lean proofs. Build all CUP proofs:

```
cd ugp-lean
lake build UgpLean.Universality.CUP4TotalParity
lake build UgpLean.Universality.CUP11ModSeven
lake build UgpLean.Universality.CUP3DUniqueness
lake build UgpLean.Universality.CUP3DPSCUnification
```

Expected: all build successfully, zero sorry, after Mathlib cache is populated.

CUP-4 null test.

```
cd papers/28_computational_universality/canonical_run
python3 t_null_cup4.py
```

Expected runtime: <60s. Output matches Table 3 and $p_{\text{raw}} = 1.36\%$.

Orbit survey and perturbation tests.

```
python3 t_orbit_survey.py
python3 t_perturbed_orbits.py
python3 t_analytical_verification.py
```

Expected: Table 3 reproduced; all 7 orbit neighborhoods verified.

CA vertex table and P22 winding coverage.

```
python3 scripts/ca_vertex_table.py
```

Complete 343-entry f_{MDL} classification: output distribution, P22 winding coverage (6 of 14 neighborhoods in the P22 gauge spectrum), $\mathbb{Z}_7$ sum rule verification (3 of 14), and P22 absorption vertex cross-check.

$\mathbb{Z}_5$ rotation equivariance and $\mathbb{Z}_7$ gauge invariance tests.

```
python3 scripts/z7_gauge_invariance_check.py
```

$\mathbb{Z}_7$ additive gauge invariance (FAILS: 2030/2058 checks), $\mathbb{Z}_5$ rotation equivariance (PASSES: 0 failures over $7^5 \times 5$ cases). Runtime: <300 ms.

E Physical Dictionary: $\mathbb{Z}_7$ Winding Numbers and SM Quantum Numbers

The identification of $\mathbb{Z}_7$ winding numbers with Standard Model quantum numbers is machine-certified in Lean 4 (GUTStructure.lean §49, zero sorry). The complete physical dictionary is compiled in Table 11.

All identifications are derived from the charge formula $Q = w^*/3$, where $w^* \in \{-3, -2, -1, 0, 1, 2, 3\}$ is the centered $\mathbb{Z}_7$ representative (Lean: gte_charge_formula, GUTStructure.lean §49). The hypercharge assignments $Y_q = 1/3$ and $Y_l = -1$ follow from $\mathbb{Z}_7$ winding sums (Lean: quark_doublet_hypercharge, lepton_doublet_hypercharge, §49). The $\text{SU}(3)_c$ color sector is identified with the $\mathbb{Z}_3 = \{1, 2, 4\}$ multiplicative subgroup of $\mathbb{Z}_7^*$ (Lean: z7_color_subgroup_closed, GUTStructure.lean §33). This identification holds at the effective-field-theory level for momenta below $\Lambda_{\text{GTE}} \approx 2.0 \text{ GeV}$ (§10.12). The proposal that the GTE gauge field A_μ corresponds to the Cartan generator λ_3 of $\text{SU}(3)$ Yang–Mills, and that $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ flows to $\text{SU}(3)$ Yang–Mills above Λ_{GTE} , constitutes a UV-completion conjecture; verification is open work.

Table 11: Physical dictionary: $\mathbb{Z}_7$ winding numbers and SM quantum numbers. All identifications are machine-certified (Lean 4, zero sorry).

Particle	$\mathbb{Z}_7$ winding w	Q (electric)	Y (hypercharge)	T_3 (isospin)	Sector
γ, Z^0, ν	0	0	—	—	Neutral
u -quark	2	+2/3	+1/3	+1/2	$SU(3)_c$: $\mathbb{Z}_3$
W^+, e^+	3	+1	—	+1	Charged
e^-, W^-	4	−1	−1	−1/2	Lepton
d -quark	6	−1/3	+1/3	−1/2	$SU(3)_c$: coset

References

- [1] Matthew Cook. Universality in elementary cellular automata. *Complex Systems*, 15(1):1–40, 2004.
- [2] Matthew Cook. A concrete view of rule 110 computation. *Electronic Proceedings in Theoretical Computer Science*, 1:31–55, 2009. arXiv:0906.3248.
- [3] Jonathan Gorard. Some relativistic and gravitational properties of the Wolfram model. *Complex Systems*, 29(2):599–654, 2020.
- [4] Paul Rendell. A universal Turing machine in Conway’s Game of Life. In *2011 International Conference on High Performance Computing and Simulation (HPCS)*, pages 764–772. IEEE, 2011.
- [5] Nova Spivack. The dynamics and universality of the universal generative principle: Attractors and emergent thermodynamics. Zenodo, 2025. <https://doi.org/10.5281/zenodo.20170092>.
- [6] Nova Spivack. First principles standard model: Elegant kernel, quarter-lock law, and gauge couplings. Zenodo, 2025. <https://doi.org/10.5281/zenodo.20168787>. Primary derivation of the Standard Model and cosmological parameters from the UGP/GTE framework.
- [7] Nova Spivack. Ugp dynamics and universality: From minimal laws to emergent physics. Zenodo, 2025. <https://doi.org/10.5281/zenodo.20170092>. Derives universality classes of physical law and dynamic symmetries from the UGP’s generative flow equations.
- [8] Nova Spivack. Arithmetic derivation of the electroweak mixing angle from rule 110 orbit arithmetic. Zenodo, 2026. Paper P31 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319413>.
- [9] Nova Spivack. The CKM Wolfenstein parameters from generative triple evolution orbit arithmetic. Zenodo, 2026. Paper P32 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319415>.
- [10] Nova Spivack. The complete GTE framework: Standard Model, gravity, quantum mechanics, and cosmology from ϕ_{mdl} . Zenodo, 2026. <https://doi.org/10.5281/zenodo.20560550>. Preprint.
- [11] Nova Spivack. Deeper consequences of arithmetic universality in the standard model. Zenodo, 2026. Paper P33 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319417>.
- [12] Nova Spivack. Emergent gravity from rule 110 cellular automaton. Zenodo, 2026. Paper P36 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319425>.
- [13] Nova Spivack. From ugp to gte: Prime-locked universes, minimality, and the emergence of our world. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20171002>. UGP Physics Programme, Paper 08 (P08).
- [14] Nova Spivack. The GTE-Möbius substrate: Arithmetic unification of computation and transputation in the universal generative principle. Zenodo, 2026. Paper P34 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319419>.

- [15] Nova Spivack. GTE unification: A bidirectional correspondence between rule 110 orbit arithmetic and the standard model electroweak sector. Zenodo, 2026. Paper P35 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319421>.
- [16] Nova Spivack. MDL selects the Wolfram rule: $\mathbb{Z}_7$ dynamics, algebraic structure, and Standard Model encoding of the GTE polynomial. Preprint, 2026. <https://doi.org/10.5281/zenodo.20572656>. UGP Physics series, P49.
- [17] Nova Spivack. The mirror branch braid atlas: A parameter-free dark sector from the universal generative principle. Zenodo, 2026. Paper P29 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20263362>.
- [18] Nova Spivack. The ϕ_{mdl} field: Quantum structure, born rule, and continuum completion of the chiral minkowski CA. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20417576>. Paper P42 in the UGP Physics series.
- [19] Nova Spivack. QCD structure from the generative triple evolution substrate: Asymptotic freedom, confinement, and hadron spectroscopy from $f_{21} \subset \text{SU}(3)$. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20417564>. Paper P39 in the UGP Physics series. Canonical QCD paper: α_s , $b_0 = 7$, $b_1 = 26$, $b_2 = 180.9$, F_{21} colour factors, hadron spectroscopy.
- [20] Nova Spivack. Quantum gravity in the GTE/ Φ_{MDL} framework: Functional completeness. Preprint (Zenodo), 2026. <https://doi.org/10.5281/zenodo.20465807>. UGP Physics series, P44.
- [21] Nova Spivack. Reflexive reality research programme — complete paper corpus. Zenodo, 2026. <https://doi.org/10.5281/zenodo.19487040>. Complete corpus of the NEMS and UGP Physics programmes, including NEMS Paper 05 (PSC-optimality of the Standard Model) and NEMS Papers 10–11 (transputation and physical incompleteness). Concept DOI always resolves to the latest version. Research portal: <https://www.novaspivack.com/research>.
- [22] Nova Spivack. The UGP Interaction Skeleton: Lean-Certified SM Gauge-Fermion Vertices, Anomaly Cancellation, and Topological Selection from Braid Topology. Zenodo, 2026. Paper P22 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20170132>.
- [23] Nova Spivack. ugp-lean: A machine-checked formalization of the universal generative principle in Lean 4. Zenodo, 2026. Formalization paper (latest version): <https://doi.org/10.5281/zenodo.19554688>. Concept DOI (always-latest): <https://doi.org/10.5281/zenodo.19433538>. More than 400 modules across seventeen architectural layers including GTE, Universality, ElegantKernel (UCL unconditional closure), MassRelations (Koidé closed form, S_3 -Newton flow, binary cascade, physical mass spectrum), BraidAtlas (ChargeTheorem, CompositeTriples, ChiralitySquaring, ChargeDerivation, CoxeterConductor, CoxeterConductorTowerLaw, EWBosons), GaloisStructure, PSC RCC infinite-family closure, and TE2.2 scan certification. Zero sorry throughout; 98 disclosed named axioms across the library (counted via top-level `axiom` declarations across all `.lean` files, deduplicated by name, excluding standard Lean/Mathlib logical axioms), none on the core arithmetic/algebraic proof path — see the formalization paper’s axiom-inventory table for the current, complete list. Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [24] Gerard ’t Hooft. *The Cellular Automaton Interpretation of Quantum Mechanics*, volume 185 of *Fundamental Theories of Physics*. Springer, Cham, 2016. Open Access.
- [25] A. M. Turing. On computable numbers, with an application to the entscheidungsproblem. *Proceedings of the London Mathematical Society*, s2-42(1):230–265, 1936.
- [26] Stephen Wolfram. Universality and complexity in cellular automata. *Physica D: Nonlinear Phenomena*, 10(1–2):1–35, 1984.
- [27] Stephen Wolfram. *A New Kind of Science*. Wolfram Media, Champaign, IL, 2002.
- [28] Stephen Wolfram. A class of models with the potential to represent fundamental physics. *Complex Systems*, 29(2):107–536, 2020.