

Formal and Computational Concordance on PSC-Selected Standard Model Structure: *Axiomatic Closure Theorems and Finite Universe Enumeration*

Nova Spivack

2026

Preprint.

Archival version

(Zenodo): <https://doi.org/10.5281/zenodo.20169970>

UGP series hub

(Zenodo): <https://doi.org/10.5281/zenodo.20168144>

Abstract

We present the computational certificate that complements the Two-Layer PSC Theorem [6, 9], together with a full constraint independence analysis and three new UGP-derived coupling-ratio predictions.

Formal backbone (Papers 03 and 05).

The PSC theorem chain— $\text{RC} \Rightarrow \text{NM}^* \Rightarrow \{\text{GUT exclusion, vector-like exclusion, CP violation required}\}$ —establishes as formal theorems that any self-contained 4D renormalizable gauge theory must have (i) $G = G_{\text{SM}}$ and (ii) $N_{\text{gen}} \geq 3$ (Layer I, forced by consistency). Presentation Invariance further selects $N_{\text{gen}} = 3$ as the unique minimal Layer I survivor (Layer II, optimality). Key steps are machine-checked in nems-lean and ugp-lean with zero `sorry` [11]. Full uniqueness is now unconditional: the Residual Classification (RCC) is a Lean-certified theorem [12] (`PSC.RCCInfiniteFamilies`, zero `sorry`).

Computational certificate (TE2.2 scan, this paper). An exhaustive enumeration of 20,160 candidate universe descriptions minimises a PSC dissonance functional $D[\Psi]$. Only 12 pass the hard PSC Layer I filters (0.06%); all 12 are SM-like. **The five hard filters ($C_1, C_6, C_8, C_{12}, C_{13}$) contain no reference to G_{SM}** — they encode dimensional consistency, holographic closure, unitary evolution, area law, and Kähler structure. The SM-targeted constraints (C_2, C_3) are soft penalties applied only at Layer II; the 12-of-20,160 figure is a result of the non-SM-targeted filters alone. The

SM tuple $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ achieves the unique Layer II minimum $D_{\text{min}} = 1.0667$ (Hessian $\lambda_{\text{min}} = 2.0 > 0$). The residual $D_{\text{min}} > 0$ is entirely due to C_4 (Quarter-Lock), a UGP-derived coupling-ratio prediction [11] satisfied by the SM to 95% at M_Z ; all other 13 constraints evaluate to zero at the SM point.

New UGP-derived predictions. We add three new constraints derived from UGP orbit invariants (not from SM data): the bare g_1^2/g_2^2 ratio (C_{15}), the bare g_3^2/g_2^2 ratio (C_{16}), and a multi-scale Quarter-Lock test (C'_4). These are machine-checked rationals in ugp-lean; the SM satisfies all three to within 2–7% after RG running.

Constraint independence. We give a full taxonomy of all 16 constraint terms (base C_1 – C_{14} plus extended-scan C_{15}, C_{16} ; §4.1): 2 are explicitly SM-targeted (C_2, C_3); C_5, C_9, C_{11} use principled computations (one-loop RG stability, anomaly cancellation — see §4.1); 1 is algebraically fitted to Λ_{obs} (C_{14}); 3 are genuine UGP-derived predictions (C_4, C_{15}, C_{16}); and 7 are generically PSC-derived. A pre-committed ablation study confirms that the SM is the D -minimizer under the full principled set; the 12-of-20,160 Layer I result holds under the 5 non-SM-targeted hard filters alone.

Extended gauge group scan. The extended scan covers Pati-Salam, E_6 , G_2 , and $\text{SU}(6)$; all major BSM candidates fail the PSC sieve. The continuous cosmological parameters $(\Lambda, \rho, \kappa, \tau)$ remain underdetermined within the SM family (four co-minimizers).

Contents

1 Introduction

2

2 The Two-Layer PSC Theorem: Formal

Side	3
2.1 PSC, RC, and NM*	3
2.2 The Two-Layer PSC Theorem	4
2.3 The PSC Sieve and Residual Classification Conjecture	5
2.4 Status of Lean Formalization	5
3 The TE2.2 Computational Certificate	5
3.1 Overview and Methodology	5
3.2 The Universe Parameter Space	6
3.3 The Dissonance Functional	6
3.4 Scan Methodology	7
3.5 Results	7
3.6 Scope and Interpretation	10
4 Theory–Computation Concordance	10
4.1 Methodological Independence: Extent and Limits	10
4.2 The Concordance Table	12
4.3 Why the Agreement Strengthens the Case	12
4.4 What the Concordance Establishes	14
5 Residuals and Controlled Tensions	14
5.1 Residual Classification (RCC)	14
5.2 The Discretization Gap	14
5.3 The Neutrino Sector	15
5.4 Vector-Like Fermion Representations	15
5.5 Falsifiability	15
6 Conclusion	16
A The Fourteen TE2.2 Constraint Terms	16
B Reproducibility	17
C Lean Formalization of the PSC Gauge Sieve	18

1 Introduction

A central question in fundamental physics is whether the Standard Model (SM) of particle physics is a contingent outcome of cosmic initial conditions, or whether it is a privileged—perhaps uniquely self-consistent—structure forced by deep internal requirements. The SM gauge group $G_{\text{SM}} = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$, its three generations of chiral fermions, and its renormalizability structure were historically assembled to fit experimental data, not derived from first principles. Yet several independent research

programmes have sought axiomatic or variational principles that would *constrain or select* this structure rather than merely accommodate it. The algebraic structure of the SM gauge group and its embedding in GUTs has been analyzed in detail by Baez and Huerta [1]; the octonion/exceptional-algebra approach provides one perspective on why $S(\text{U}(2) \times \text{U}(3))$ is structurally special. The present paper does not claim to resolve this question unconditionally; rather, it assembles two lines of evidence and shows that they converge on the same answer subject to explicitly stated conditions.

The Perfect Self-Containment (PSC) programme [5, 7] asks whether a universe can be fully self-explaining: every law, structure, and constant must be generated by the universe’s own dynamics without external inputs or arbitrary parameters. PSC is formalized as a conjunction of closure conditions—renormalizability, anomaly cancellation, information-theoretic viability, geometrodynamical consistency, and others—that collectively constrain the space of possible physical theories.

Paper 03 of the NEMS/PSC suite [9] established that PSC implies the Renormalizability Criterion (RC) and the No-Miracles structural stability condition (NM*), and derived from these the exclusion of GUT groups, vector-like fermion content, and CP-conserving theories. Paper 05 [6] proved the Two-Layer PSC Theorem, showing that a first layer of structural conditions forces $G = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ and $N_{\text{gen}} \geq 3$, while a second information-theoretic layer selects $N_{\text{gen}} = 3$ uniquely. Papers 20 and 21 [8, 10] formalized the PSC Sieve in Lean 4 and established that the Standard Model is the *only* ontologically legal foundation for a PSC universe (now unconditionally: the Residual Classification is a Lean-certified theorem, `PSC.RCCInfiniteFamilies` [12], zero `sorry`).

These results constitute a powerful formal case. However, the formal approach operates within an axiomatic system, and its conclusions depend on the completeness and correctness of the premise bundle. An independent *computational* check, using entirely different methodology, provides complementary evidence.

The TE2.2 scan [4] takes a different route. Rather than reasoning syntactically from axioms, it constructs a *dissonance functional* $D[\Psi]$ that numerically measures how far a candidate universe deviates from each of the fourteen PSC constraint con-

ditions. It then exhaustively evaluates $D[\Psi]$ over 20,160 discretized universe descriptions parameterized by spacetime dimension, gauge group, number of generations, number of observers, cosmological constant, information profit ratio, spatial curvature, and topology. The Standard Model gauge structure $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ emerges as the unique co-minimum family at $D_{\text{min}} = 1.0667$, and only 12 of the 20,160 candidates pass the hard PSC filters at all.

The key observation of this paper is that the formal and computational results are largely methodologically independent. The formal side reasons by syntactic deduction; the computational side reasons by exhaustive numeric evaluation; they share no common proof steps, code, or deductive machinery. Two of the fourteen scan constraints (C_2, C_3) do encode the SM SRRG fixed-point result—a partial theoretical prior—but the remaining twelve constraints, and crucially the hard PSC filters, are defined independently of the SM answer. The convergence of both methods on the same answer constitutes a non-trivial concordance that we characterize carefully in §4.1.

Organization. Section 2 surveys the formal side, restating the key theorems from Papers 03, 05, 20, and 21 without re-deriving them. Section 3 gives a detailed account of the TE2.2 computational certificate: the dissonance functional, the fourteen constraint terms, the scan methodology, and the results. Section 4 articulates the concordance argument and presents a structured comparison. Section 5 characterizes the residuals and controlled tensions. Section 6 concludes. Appendix A lists all fourteen TE2.2 constraint terms, and Appendix B gives reproducibility information.

Novel contribution. The novel contributions of this paper are:

- (i) **Computational certificate for the Two-Layer PSC Theorem.** The TE2.2 scan provides a verified finite certificate that the SM is the unique D -minimizer over 20,160 candidate universes, constituting numerical evidence for the RCC at the discretized level.
- (ii) **D_{SM} decomposition.** A per-constraint breakdown (Table 1) shows $D_{\text{SM}} = 1.0667$ is entirely due to C_4 (Quarter-Lock): 13 of 14 constraints evaluate to zero at the SM point.

- (iii) **Full constraint independence taxonomy.** Complete classification of all constraints by SM-specificity (§4.1), correcting the prior “2 of 14” characterisation to an accurate “5 SM-targeted, 3 UGP-derived predictions, 7 generic PSC-derived.”
- (iv) **New UGP-derived coupling-ratio predictions** (C_{15}, C_{16}): bare g_1^2/g_2^2 and g_3^2/g_2^2 from ugp-lean (machine-checked rationals), not derived from SM data.
- (v) **Extended BSM gauge group scan:** Pati-Salam, E_6 , G_2 , $SU(6)$, confirming all major BSM candidates fail the sieve.
- (vi) **Audit of C5 Landau pole and C14 algebraic fit:** C5 does not evaluate the SM’s $U(1)$ UV behavior; C14 uses a formula fitted to Λ_{obs} , not derived prior to it. Both are disclosed and the paper’s independence claim is revised accordingly.
- (vii) **Residual taxonomy** (RCC, discretization gap, neutrino sector, co-minimizer underdetermination of $\Lambda, \rho, \kappa, \tau$).

Conventions. We write $G_{\text{SM}} = SU(3) \times SU(2) \times U(1)$ for the Standard Model gauge group. N_{gen} denotes the number of fermion generations. The notation $\Psi = (d, G, N_{\text{gen}}, N_{\text{obs}}, \Lambda, \rho, \kappa, \tau)$ refers to the eight-parameter universe description used in the TE2.2 scan. All formal claims attributed to Papers 03–21 are cited and not re-derived here; the novel contributions are itemized in the paragraph above.

2 The Two-Layer PSC Theorem: Formal Side

We summarize the formal results established in Papers 03, 05, 20, and 21 of the NEMS/PSC suite. The reader is referred to those papers for full proofs; we state the theorems in the form needed for the concordance analysis.

2.1 PSC, RC, and NM*

Definition 2.1 (Perfect Self-Containment [9]). *A physical theory $\mathcal{T}$ is perfectly self-contained (PSC) if it satisfies all of the following simultaneously:*

- (i) **Renormalizability Criterion (RC)**: $\mathcal{T}$ is a renormalizable gauge theory in $d = 4$ spacetime dimensions;
- (ii) **Structural Stability (NM*)**: $\mathcal{T}$ is structurally stable against small deformations; no external tuning is required to maintain its gauge structure;
- (iii) **Anomaly Cancellation**: all gauge and mixed gravitational anomalies cancel;
- (iv) **Topological Viability (TV)**: the gauge bundle admits consistent instanton sectors;
- (v) **Spectral Asymmetry (SA)**: the Dirac spectrum distinguishes matter from anti-matter.

Remark 2.2 (Scope of the axiomatic approach). The PSC conditions in Definition 2.1 are not themselves derived from a more primitive framework; they constitute a proposed axiomatic system whose physical motivation is given in Papers 01–03 [7, 9]. The formal programme establishes what *follows from* PSC, not that PSC is the unique or correct set of self-consistency requirements. The question of whether the PSC axioms are the “right” axioms for describing a self-contained universe is a physical and philosophical question separate from the formal and computational results reported here.

Remark 2.3 (TV and SA nomenclature in the Lean formalization). The axiom names TV (Topological Viability) and SA (Spectral Asymmetry) as used here correspond to the abstract Layer I conditions; the current Lean formalization uses the names Thermodynamic Viability and Semantic Admissibility for the same structural positions (NemS/Physics/Rigidity.lean).

Paper 03 [9] establishes a chain of implications from PSC to the exclusion of large classes of theories.

Theorem 2.4 (PSC Exclusions [9]). *Let $\mathcal{T}$ be a PSC theory in four spacetime dimensions. Then:*

- (i) *GUT gauge groups ($SU(5)$, $SO(10)$, $E_6, \dots$) are excluded;*
- (ii) *Vector-like fermion representations are excluded;*
- (iii) *CP-conserving theories are excluded.*

These exclusions arise from the interaction of RC with NM*: GUT groups and vector-like representations fail structural stability under renormalization-group running, and CP-conservation is incompatible with the full anomaly structure required by SA.

2.2 The Two-Layer PSC Theorem

Paper 05 [6] tightens the exclusions of Theorem 2.4 into a positive selection result. The key insight is that the PSC conditions split into two logically distinct layers with different filtering power.

Definition 2.5 (Layer I and Layer II [6]). •

Layer I comprises the structural conditions: Renormalizability (RC), No-Miracles stability (NM*), Topological Viability (TV), Spectral Asymmetry (SA), and Anomaly Cancellation.

- **Layer II** comprises the information-theoretic conditions: Principle of Informational Completeness (PI) and the Minimum Description Length criterion (MDL).

Theorem 2.6 (Two-Layer PSC Theorem [6]). *Let $\mathcal{T}$ be a PSC theory.*

- (i) **Layer I selection**: *The Layer I conditions jointly force the gauge group $G = SU(3) \times SU(2) \times U(1)$ and the generation count $N_{\text{gen}} \geq 3$.*
- (ii) **Layer II selection**: *Among theories satisfying Layer I, the Layer II conditions (PI/MDL) uniquely select $N_{\text{gen}} = 3$.*

The proof of Layer I combines the exclusion results of Theorem 2.4 with a classification of the anomaly-free representations of simple and semisimple gauge groups in four dimensions. The combination of chirality and anomaly cancellation forces the $SU(3) \times SU(2) \times U(1)$ structure. The Layer II proof uses an MDL argument: fewer than three generations fails informational completeness (the theory cannot encode its own coupling constants), while more than three generations violates the minimum-description-length criterion (the additional structure is compressible and hence redundant under PSC).

Remark 2.7. The Two-Layer structure reflects two genuinely distinct necessity mechanisms. Layer I is a purely structural-algebraic argument; Layer II is an information-theoretic argument. Their combination produces the sharp $N_{\text{gen}} = 3$ selection.

Remark 2.8 ($N_{\text{gen}} \geq 3$ is forced, not merely optimal). A common misreading frames $N_{\text{gen}} = 3$ purely as an optimality result (Layer II). This is imprecise. *Layer I itself forces* $N_{\text{gen}} \geq 3$ as a consistency requirement [9]: the exclusion of CP-conserving theories (Theorem 2.4(iv)) follows from the NM^* condition, and CP violation in a four-dimensional chiral gauge theory requires at least three generations of fermions (Kobayashi–Maskawa mechanism). Thus $N_{\text{gen}} \geq 3$ is not a choice or an optimization; it is *forced* by structural self-containment. Layer II then adds the further claim that among all Layer I survivors, $N_{\text{gen}} = 3$ is uniquely MDL-minimal. $N_{\text{gen}} = 4$ is PSC-consistent but not PSC-optimal.

2.3 The PSC Sieve and Residual Classification Conjecture

Papers 20 and 21 [8, 10] extend the Two-Layer Theorem to a full formal sieve and address the question of uniqueness.

Definition 2.9 (PSC Sieve [8]). *The PSC Sieve is the formal decision procedure that takes as input a candidate gauge theory $\mathcal{T}$ and outputs ADMIT or REJECT based on sequential evaluation of all PSC closure conditions, with the Two-Layer Theorem applied at the gauge-structure and generation stages.*

Paper 20 [8] formalizes all sieve constraints in Lean 4 [11] and verifies that the SM passes the sieve while all known non-SM theories fail. The formalization uses the `ugp-lean` and `nems-lean` Lean 4 libraries and is publicly archived [11].

Within the PSC framework, residual classification asserts that no gauge-theory completion inequivalent to the SM survives the full PSC Sieve.

Definition 2.10 (Residual Classification (RCC) [8]). *No gauge theory other than the Standard Model (up to physically equivalent representations) passes the full PSC Sieve.*

Theorem 2.11 (Existential Rigidity [10, 12]). *Under Definition 2.10, established in Section 5.1 via TE2.2-style enumeration, extended gauge certification, and Lean PSC.RCCInfiniteFamilies, the Standard Model is the only ontologically legal foundation for a PSC universe: any PSC-consistent universe necessarily has gauge structure G_{SM} and $N_{\text{gen}} = 3$.*

Section 5 summarizes the consolidated components of this classification statement.

2.4 Status of Lean Formalization

A substantial portion of the formal PSC programme has been machine-checked in Lean 4. The formalization [11] includes:

- All Layer I structural conditions as typed Lean predicates;
- The anomaly cancellation check as a Lean decision procedure;
- The Renormalizability Criterion and NM^* as formal stability conditions on gauge group representations;
- The PSC Sieve as a formal function from theory descriptions to $\{\text{ADMIT}, \text{REJECT}\}$;
- Machine-verified certificates that G_{SM} passes the sieve with $N_{\text{gen}} = 3$.

The formalization achieves zero `sorry` statements and uses no non-standard axioms beyond the Lean 4 core. This provides the strongest available formal assurance that the axiomatic reasoning is internally consistent and that the stated theorems follow from their premises.

3 The TE2.2 Computational Certificate

We now describe the second, entirely independent line of evidence: the TE2.2 finite exhaustive enumeration.

3.1 Overview and Methodology

The TE2.2 scan [4] approaches the question of SM selection computationally. Rather than reasoning from axioms, it:

- (1) Constructs a scalar functional $D[\Psi]$ that numerically measures PSC constraint violations;
- (2) Parameterizes the space of candidate universes by eight discrete parameters;
- (3) Exhaustively evaluates $D[\Psi]$ over all 20,160 points in this parameterized space;
- (4) Identifies the global minimizer and examines all PSC-passing candidates.

The approach is strictly computational: no theorem is proved, but a computational certificate is produced. The certificate is reproducible and SHA-256 verified (see Appendix B).

3.2 The Universe Parameter Space

A universe description Ψ is an eight-tuple

$$\Psi = (d, G, N_{\text{gen}}, N_{\text{obs}}, \Lambda, \rho, \kappa, \tau), \quad (1)$$

where the components and their discrete ranges (as implemented in the TE2.2 scan code [4, 13]) are:

- $d \in \{2, 3, 4, 5, 6\}$: spacetime dimension (5 values);
- $G \in \{\text{U}(1), \text{SU}(2), \text{SU}(3), \text{SU}(2) \times \text{U}(1), \text{SU}(3) \times \text{SU}(2) \times \text{U}(1), \text{SU}(5), \text{SO}(10)\}$: gauge group (7 values);
- $N_{\text{gen}} \in \{1, 2, 3, 4\}$: number of fermion generations (4 values);
- $N_{\text{obs}} \in \{0, 1\}$: number of observer classes (2 values);
- $\Lambda \in \{0, 10^{-122}, 10^{-60}\}$: cosmological constant (Planck units, 3 values);
- $\rho \in \{0.5, 1.0, 1.13, 1.5\}$: information profit ratio (4 values);
- $\kappa \in \{0, 0.01, -0.01\}$: spatial curvature (3 values);
- $\tau \in \{\text{flat}, \text{hyperbolic}\}$: global topology (2 values).

The Cartesian product of these ranges yields $5 \times 7 \times 4 \times 2 \times 3 \times 4 \times 3 \times 2 = 20,160$ universe descriptions, which are exhaustively enumerated.

3.3 The Dissonance Functional

The central object of the TE2.2 scan is the *dissonance functional*:

$$D[\Psi] = \sum_{i=1}^{14} w_i \|C_i[\Psi]\|^2, \quad (2)$$

where $C_i[\Psi]$ is the violation of the i -th PSC constraint, the norm $\|\cdot\|$ is appropriate to the type of constraint (scalar absolute value, or a norm on a vector of violations), and $w_i > 0$ are positive weights calibrated so that all constraints contribute comparably at the boundary of the feasible region.

The 14 constraint terms are:

1. **Dimensional** (C_1): Violation of the $d = 4$ dimensional optimality condition. $C_1[\Psi] = |d - 4|$.
2. **SRRG Fixed Point** (C_2): Deviation of the gauge group and generation count from the renormalization-group fixed point ($G_{\text{SM}}, N_{\text{gen}} = 3$).
3. **SRRG Viability** (C_3): Failure of the gauge coupling functional $F[S]$ to be maximal on the self-renewing renormalization group (SRRG) fixed point.
4. **Quarter-Lock** (C_4): Deviation from the gauge-coupling ratio condition $\sqrt{3}g_1 \approx g_2$. This condition is derived from the UGP orbit invariant $k_M = k_G + \frac{1}{4}k_L^2$ (machine-checked as `quarterLockLaw` in `ugp-lean`, Zenodo 10.5281/zenodo.19433538) — a UGP-derived prediction that the SM approximately satisfies at the M_Z scale (5% deviation; see §3.5). This is distinct from the Weinberg angle $\sin^2 \theta_W$, which encodes a different relation between g_1 and g_2 .
5. **RG Flow Stability** (C_5): Positive rate of change of the coupling constants, $dc/dt > 0$, which violates RG flow stability.
6. **Kähler Structure** (C_6): Failure of the Fisher information metric on coupling-constant space to be symplectic (Kähler).
7. **Area Law** (C_7): Deviation of the entanglement entropy from the Bekenstein–Hawking area law with logarithmic correction: $S = A/(4\ell_P^2) + \beta \log A$.
8. **Unitary Evolution** (C_8): Failure of the CP-structure to ensure unitary time evolution.
9. **RIET Equivalence** (C_9): Violation of the Reflexive Information-Energy-Topology equivalence, i.e., failure of curvature, energy, entropy, and information to be mutually dual.
10. **Einstein Equations** (C_{10}): Deviation of the dynamical equations from $G_{\mu\nu} = 8\pi G T_{\mu\nu}$.
11. **Coherence Field** (C_{11}): Failure of the coherence field Ψ to couple consistently to all matter sectors.
12. **Information Profit** (C_{12}): Violation of $\text{Gen/Drain} \geq 1.13$, i.e., $\rho < 1.13$.
13. **Necessary Observers** (C_{13}): $N_{\text{obs}} < 1$.

14. **Lambda Relation** (C_{14}): Deviation of the cosmological constant from the structurally-fit value $\Lambda_{\text{PSC}} = \ln(\varphi)/\ln(2\pi) \approx 10^{-122}$ (Planck units), where φ is the golden ratio. This formula matches the observed Λ_{obs} ; an independent analytic derivation from PSC axioms alone has not yet been established (see §4.1).

The full analytic definitions of each C_i are given in Appendix A.

Remark 3.1. The constraint terms C_1 – C_{14} are not axioms—they are numerical functions of the universe parameters that vanish when a particular PSC condition is exactly satisfied. The functional $D[\Psi]$ is therefore a quantitative measure of total PSC-inconsistency, and its global minimum is achieved when PSC-inconsistency is simultaneously minimized across all fourteen dimensions.

3.4 Scan Methodology

The scan proceeds in three phases.

Step 1: Analytic constraint verification. Each of the 14 constraint functions C_i is implemented analytically in Python. The Standard Model point $\Psi_{\text{SM}} = (4, G_{\text{SM}}, 3, 1, 10^{-122}, 1.13, 0, \text{flat})$ is evaluated and $D[\Psi_{\text{SM}}]$ is computed. The Hessian $\nabla^2 D$ at Ψ_{SM} is computed numerically and its eigenvalues are analyzed.

Step 2: Finite exhaustive enumeration. The parameterized universe space is enumerated as a Cartesian product. For each of the 20,160 universe descriptions, $D[\Psi]$ is evaluated and the PSC pass/fail decision is recorded. A universe passes the hard PSC filters if each “hard” constraint (Kähler structure, unitary evolution, information profit, necessary observers, dimensional optimality) is satisfied within tolerance. Results are sorted by $D[\Psi]$ and the top-ranked universes are recorded.

Step 3: Extension to the continuum. An analytic extension argument (detailed in [4]) is used to argue that the discrete certificate extends to the continuum for the *continuous* parameters (Λ, ρ, κ) : (i) $D[\Psi]$ is continuous and coercive in those parameters; (ii) the PSC-admissible region for continuous parameters is compact; together these imply the discrete minimizer extends to a minimizer over the continuum

for those parameters. Note that the analytic extension relies on the continuity and coercivity of $D[\Psi]$, not on topological density of the finite grid: three sample points per axis is *not* topologically dense in $\mathbb{R}$, but is sufficient to establish the local minimum structure via the Hessian analysis (§3.5), which confirms $\lambda_{\min} = 2.0 > 0$ at Ψ_{SM} . For the *discrete* parameters $(d, G, N_{\text{gen}}, N_{\text{obs}}, \tau)$, no continuity argument applies; the discrete grid exhausts all values considered for these parameters, and the certificate is valid exactly as stated for them. The extension argument does not cover gauge groups outside the seven in the grid, nor generation counts above 4; the limitations are as stated in §5.2.

3.5 Results

The TE2.2 scan [4] produces the following computational certificate.

Summary statistics. The scan evaluated 20,160 universe descriptions in 0.14 seconds (throughput: $\approx 144,257$ universes/second). Of the 20,160 candidates:

- **12** pass the hard PSC filters (0.06%);
- All 12 PSC-passing universes have $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$;
- **20,148** are rejected by at least one hard PSC constraint.

Global minimizer. The canonical global minimum is achieved by $\Psi_{\text{SM}} = (4, G_{\text{SM}}, 3, 1, 10^{-122}, 1.13, 0, \text{flat})$, with

$$D_{\text{SM}} = D[\Psi_{\text{SM}}] = 1.066657903568035. \quad (3)$$

This value equals $D_{\min}$: no other universe description achieves a lower dissonance value. However, three other universe descriptions—sharing $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ but differing in topology ($\tau = \text{hyperbolic}$) or information profit ratio ($\rho = 1.5$ vs. 1.13)—achieve the same minimum $D_{\min} = 1.0667$. Thus, the global minimum at the level of the dissonance value is *degenerate*: four universe descriptions are co-minimizers. The SM is the unique minimizer only at the level of the *gauge-structure-and-generation parameter tuple* (d, G, N_{gen}) : no universe with $(d, G, N_{\text{gen}}) \neq (4, G_{\text{SM}}, 3)$ achieves $D[\Psi] < D_{\text{SM}}$ or equals $D_{\min}$.

The co-minimizer degeneracy has a specific structural origin: constraint C_{12} (Information Profit) penalizes only $\rho < 1.13$; any $\rho \geq 1.13$ gives exactly zero

violation. Thus the scan establishes a *lower bound* on the viable information profit ratio ($\rho \geq 1.13$), not a unique value. The continuous cosmological parameters ($\Lambda, \rho, \kappa, \tau$) are therefore *underdetermined* within the SM family by the current scan: the result uniquely selects the discrete gauge structure and generation count $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$, but does not pin down the continuous cosmological parameters within that family.

Constraint breakdown at Ψ_{SM} . Table 1 decomposes $D_{\text{SM}} = 1.0667$ by individual constraint. All 13 non- C_4 constraints evaluate to exactly zero at the SM point; the entire dissonance is carried by C_4 (Quarter-Lock).

Table 1: Individual constraint contributions to $D[\Psi_{\text{SM}}] = 1.0667$. $C_i[\Psi_{\text{SM}}]$ is the raw constraint violation; the weighted contribution is $w_i \times C_i[\Psi_{\text{SM}}]$. Only C_4 is non-zero.

#	Constraint	$C_i(\Psi_{\text{SM}})$	Weighted	Note
C_1	Dimensional ($d = 4$)	0.000	0.000	
C_2	SRRG Fixed Point	0.000	0.000	SM-targeted
C_3	SRRG Viability	0.000	0.000	SM-targeted
C_4	Quarter-Lock	0.001067	1.067	UGP-derived; 5% deficit at M_Z
C_5	RG Flow Stability	0.000	0.000	SM-targeted
C_6	Kähler Structure	0.000	0.000	
C_7	Area Law	0.000	0.000	
C_8	Unitary Evolution	0.000	0.000	
C_9	RIET Equivalence	0.000	0.000	SM-targeted (proxy)
C_{10}	Einstein Equations	0.000	0.000	
C_{11}	Coherence Field	0.000	0.000	SM-targeted (proxy)
C_{12}	Information Profit	0.000	0.000	$\rho_{\text{SM}} = 1.13 \geq 1.13$
C_{13}	Necessary Observers	0.000	0.000	
C_{14}	Lambda Relation	0.000	0.000	$\Lambda_{\text{SM}} = 10^{-122}$
Total			1.067	

The C_4 contribution arises because $\sqrt{3}g_1 = 0.6191$ whereas $g_2 = 0.6517$ at M_Z (ratio 0.950), a 5% deficit from the UGP Quarter-Lock prediction. This is not a scan artefact: C_4 encodes a genuine UGP-derived prediction ($k_M = k_G + \frac{1}{4}k_L^2$, `quarterLockLaw` in `ugp-lean`) that the SM approximately but not exactly satisfies at the M_Z renormalization scale. The 5% deviation may reflect renormalization-scale running (the relation may be exact at a higher scale); this is an open question for the programme.

Extended BSM gauge group scan. The canonical scan uses 7 gauge groups. The extended scan adds 5 major BSM candidates: Pati-Salam $\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$, E_6 , G_2 , $\text{SU}(6)$, and $\text{SU}(4)$, for a total of 12 gauge groups and 34,560 universe descriptions (SHA-256: 407078d7..., `results/extended_scan_results.json`).

- All 5 new BSM groups fail the hard PSC filters without exception; their minimum dissonance is $D \approx 2.19 \times 10^6$ (vs. $D_{\text{SM}} = 1.009$ in the extended scan).
- The 12 PSC-passing universes are unchanged: all have $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$, now representing 0.035% of the extended grid.

The failure modes of the new BSM groups are driven by the hard PSC filters C_1 (dimensionality), C_6 (Kähler structure), and C_8 (unitary evolution), not by the SM-targeted constraints $C_2/C_3/C_5$. This confirms that the sieve result is robust to the gauge group choice and covers all major alternatives considered in BSM physics.

UGP-derived coupling predictions. Three new constraints, derived from `ugp-lean` machine-checked rational constants (`g1Sq_bare`, `g2Sq_bare`, `g3Sq_bare`, Zenodo 10.5281/zenodo.19433538), are included in the extended scan:

- C_{15} (UGP g_1^2/g_2^2 ratio): $g_1^2/g_2^2|_{\text{UGP}} = \frac{16/125}{2329/5400} = \frac{86400}{291125} \approx 0.2969$. SM at M_Z : 0.3008. Deviation: 1.34%; weighted contribution: 0.018.
- C_{16} (UGP g_3^2/g_2^2 ratio): $g_3^2/g_2^2|_{\text{UGP}} \approx 3.444$. SM at M_Z : 3.510. Deviation: 1.90%; weighted contribution: 0.036.
- C'_4 (Quarter-Lock exact): $g_1^2/g_2^2 = 1/3$ (exact, at the UGP unification scale). SM at M_Z : 0.301. Deviation: 9.77%; weighted contribution: 0.955.

These predictions are not derived from SM coupling data; they are algebraic consequences of the UGP orbit invariants ($k_M = k_G + \frac{1}{4}k_L^2$, $k_L^2 = 7/512$) and the group-specific discrete invariants (U(1) 3-volume, SU(2) harmonic mean, SU(3) Vandermonde). The 1–2% deviations of C_{15} and C_{16} from the SM values at M_Z are consistent with RG running from the UGP unification scale to M_Z ; the 9.77% deviation of C'_4 is the same physical gap already encoded in C_4 , expressed as a ratio rather than a coupling ratio.

Additional Lean-certified tree-level EW predictions. Two further observables are now Lean-certified at tree level from the same bare coupling triple (C_{17} – C_{18}):

- C_{17} (tree-level $\sin^2\theta_W$): $g_1^2/(g_1^2 + g_2^2) = 3456/15101 \approx 0.2289$. PDG: 0.23121. Deviation: 1.0%; proved as `sin2_theta_W_bare_eq` (zero sorry, `ugp-lean`).
- C_{18} (tree-level $1/\alpha_{\text{EM}}$): $4\pi(1/g_1^2 + 1/g_2^2) = 4\pi \cdot 377525/37264 \approx 127.31$. PDG: 127.952. Deviation: 0.5%; proved as `alpha_em_formula_exact` (zero sorry, `ugp-lean`).

Both are tree-level predictions with no threshold corrections; the sub-1% deviations match the structural ceiling established by C_{15} – C_{16} . Together with the four existing UGP-derived coupling predictions, this gives six independent Lean-certified EW observables from the same three bare rationals, all within $\leq 1.6\%$ of PDG without per-point tuning.

Four theorems about these predictions and the gauge-group labelling are proved (0 sorry) in `UgpLean.TE22.ScanCertificate`:

- `ugp_coupling_predictions_are_independent`: C_{15} , C_{16} , C'_4 are algebraically derived from `ugp-lean` constants.
- `ugp_g1g2_prediction_close_to_SM`: the UGP g_1^2/g_2^2 prediction is within 2% of the SM value at M_Z .
- `SM_gauge_uniquely_selected`: among the 60 (GaugeGroup, Dimension) pairs in the TE2.2 discretisation, exactly $(\text{SU}(3) \times \text{SU}(2) \times \text{U}(1), 4\text{D})$ satisfies the SM gauge predicate (proved by `decide`).
- `isSMGauge_iff`: full logical characterisation of the SM gauge label.
- `psc_enumeration_forces_nngen_3` (`native_decide`): all 34,560 universe descriptions are enumerated as a `Fintype`; exactly 12 pass all Layer I hard filters; all 12 have $N_{\text{gen}} = 3$ and $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ gauge in $d = 4$.
- `psc_12_survivors_card`: the cardinality of Layer I survivors equals 12 (machine-verified).
- `psc_12_survivors_have_nngen_3`: every Layer I survivor has $N_{\text{gen}} = 3$.
- `psc_admissible_forces_sm_gauge`: every Layer I survivor carries $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ gauge.

- `universe_params_card`: the full 34,560-universe parameter space is a finite `Fintype` instance.

The Layer I enumeration is fully machine-certified (Lean-certified, zero `sorry`, commit `a836ba7`): the `Fintype` instance on `UniverseParams` is provided, and 34,560 universes are exhaustively enumerated in Lean 4. A machine-checked proof that among the 12 Layer I survivors the SM carries the unique MDL-minimal description-length score (Layer II: Presentation Invariance selects the minimum- K survivor) remains to be Lean-certified.

SM rank. The Standard Model (as the canonical representative of the $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ family) ranks #1 out of 34,560 candidates in the extended scan (and #1 out of 20,160 in the canonical scan) when ordered by $D[\Psi]$.

Top-ranked structure. The four lowest-dissonance universes all have $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$, differing only in $\rho \in \{1.13, 1.5\}$ and $\tau \in \{\text{flat}, \text{hyperbolic}\}$. All four achieve $D = 1.0667$. Universes with $\Lambda = 0$ (but otherwise SM-like structure) achieve $D = 11.0667$ —a gap of approximately 10 units driven entirely by the Lambda constraint C_{14} . This provides additional numerical evidence that $\Lambda \approx 10^{-122}$ is strongly preferred by the PSC functional, and that the cosmological constant is the dominant parameter distinguishing the top from the second tier of PSC-passing universes.

Hessian stability. The Hessian $\nabla^2 D$ evaluated at Ψ_{SM} with respect to the *continuous* parameters (Λ, ρ, κ) has minimum eigenvalue

$$\lambda_{\min} = 2.0 > 0, \quad (4)$$

establishing that Ψ_{SM} is a strict local minimizer with respect to continuous perturbations of those parameters. Note that the discrete parameters $(d, G, N_{\text{gen}}, N_{\text{obs}}, \tau)$ do not admit a smooth Hessian; for these, the local stability claim rests on the discrete enumeration showing that all immediate grid neighbours have strictly higher $D[\Psi]$. Combined with global rank #1, this confirms that Ψ_{SM} is a stable minimum within the scanned parameter space. (The four co-minimizers at the same $D_{\min}$ are not isolated from each other in parameter space; topology and ρ variation within the SM (d, G, N_{gen}) family produces a degenerate valley.)

PSC rarity. Only 0.06% of the 20,160 enumerated universes pass the hard PSC filters. This “PSC rarity” confirms that the PSC conditions are highly constraining: they eliminate more than 99.9% of the universe space at the level of the discrete enumeration.

The key quantitative results are summarized in Table 2.

Table 2: TE2.2 scan summary statistics

Quantity	Value	Significance
Total universes evaluated	20,160	Exhaustive over grid
PSC-passing universes	12	0.06% of grid
SM rank	#1	Lowest- D family
$D_{\text{SM}} = D_{\text{min}}$	1.0667	Min.; 4 co-minimizers
Hessian λ_{min}	2.0	Stable local min
Scan time	0.14 s	Efficient
All PSC survivors	SM-like	$(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$

3.6 Scope and Interpretation

Several aspects of the TE2.2 result require careful interpretation.

What the scan proves (computational certificate). Within the 20,160-universe discrete parameterization, the $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ family achieves the unique minimum value $D_{\text{min}} = 1.0667$, with four co-minimizers (differing in ρ and τ). No universe with $(d, G, N_{\text{gen}}) \neq (4, G_{\text{SM}}, 3)$ achieves $D[\Psi] \leq D_{\text{min}}$, and all PSC-passing universes are SM-like. This is a *computational fact*, verified by an explicit enumeration that can be independently reproduced (see Appendix B).

What the scan does not prove. The scan does not establish, by itself, that the SM is the unique minimizer over the full infinite-dimensional space of quantum field theories. The extension to the continuum (Phase 3 of the methodology) relies on density and continuity arguments that are analytic and documented in [4] but not machine-checked. The scan also does not resolve the RCC: it confirms that no non-SM universe in the discretization passes the hard PSC filters, but it does not constitute a proof that no such universe exists in continuous theory space.

Relationship to Layer II. The TE2.2 scan implicitly incorporates a Layer II-type argument through

constraints C_{12} (Information Profit) and C_{13} (Necessary Observers). However, the information-theoretic criterion used in the scan ($\rho \geq 1.13$) is a necessary condition derived from the MDL analysis, not the full MDL argument. The numerical $N_{\text{gen}} = 3$ selection in the scan therefore has both a structural component (the SRRG fixed-point constraint C_2 prefers the SM structure) and an information-theoretic component (the profit constraint C_{12} rejects $\rho < 1.13$).

4 Theory–Computation Concor- dance

The formal and computational programmes converge on the same answer within their respective scopes. In this section we articulate why this convergence is non-trivial, what the agreement establishes (and what it does not), and how it should be read given the dependency structure identified in §3.6.

4.1 Methodological Independence: Ex- tent and Limits

The two methods are largely methodologically independent; we characterize both the extent and the limits of that independence precisely.

Formal side. The formal results of Papers 03, 05, 20, and 21 are derived by syntactic deduction from explicitly stated axioms. The key steps are:

- (i) Statement of PSC closure conditions as typed predicates;
- (ii) Algebraic classification of anomaly-free representations;
- (iii) Application of RC and NM* to exclude non-SM structures;
- (iv) MDL argument to select $N_{\text{gen}} = 3$;
- (v) Lean 4 machine-checking of all steps.

The formal side uses *no numerical computation*. Its conclusions hold if and only if its axioms hold.

Computational side. The TE2.2 scan operates by:

- (i) Constructing numerical functions $C_i[\Psi]$;

- (ii) Evaluating these functions over a finite discrete grid;
- (iii) Sorting by $D[\Psi]$ and identifying the minimum;
- (iv) Checking hard filter conditions independently.

The computational side uses *no deductive logic* and derives *no theorems*. Its conclusions are computational facts about a finite set of universe descriptions.

Independence and its limits. The two methods share no common proof steps, deductive rules, code, or numerical procedures. The formal side reasons syntactically; the computational side evaluates numerical functions over a finite grid. However, two of the fourteen TE2.2 constraint terms require careful scrutiny.

We give a complete classification of all 14 constraint terms by their degree of SM-specificity.

Explicitly SM-targeted (3 constraints). Constraint C_2 (SRRG Fixed Point) is implemented as a direct penalty for $G \neq G_{\text{SM}}$ and $N_{\text{gen}} \neq 3$. Constraint C_3 (SRRG Viability) assigns zero violation to SM-like configurations and a constant viability gap to all others. Both constraints encode the SRRG theoretical prediction—itsself derived from the formal SRRG programme [6]—that the SM is the unique RG fixed point; they operationalize this result numerically. C_5 (RG Flow Stability) is implemented as a direct check of `is_sm_like()` (source code: `RGFlowStabilityConstraint.evaluate`) and returns zero violation for SM-like universes and unit violation otherwise. The physical justification is the SRRG TS9 result that non-SM gauge theories are RG-unstable at the 97% rate; however, the implementation uses the SM as a shortcut rather than computing the actual flow for each gauge group. Specifically, C_5 does not evaluate the U(1) Landau pole that technically afflicts the SM in the UV; it passes the SM by definition. This is a limitation of the current implementation that should be addressed in future work.

SM-targeted via MFRR proxy (2 constraints). Constraints C_9 (RIET Equivalence) and C_{11} (Coherence Field Consistency) are MFRR-internal concepts whose formal definitions resolve to SM-like configurations; the scan implements both via `is_sm_like()`. These constraints should be read as “MFRR framework selects configurations that are SM-like,” not as independent physical or mathematical criteria.

Algebraically targeted at observed value (1 constraint). C_{14} (Lambda Relation) uses the formula $\Lambda_{\text{PSC}} = \ln \varphi / \ln(2\pi) \approx 10^{-122}$ as its target value. While presented as a PSC-derived prediction, this formula is an algebraic expression identified by the TE1.E programme via structural fitting to the observed Λ_{obs} ; it has not been derived from the PSC axioms prior to and independently of the observed cosmological constant value. C_{14} should therefore be read as “PSC-conditional” (the formula holds *if* the IPP structural estimator matches the observed scale) rather than PSC-derived in the strictest sense.

UGP-derived prediction (1 constraint). C_4 (Quarter-Lock) encodes $\sqrt{3}g_1 \approx g_2$, derived from the UGP orbit invariant $k_M = k_G + \frac{1}{4}k_L^2$ (`quarterLockLaw`, `ugp-lean`). This is a genuine UGP prediction that the SM approximately satisfies; it is not the Weinberg angle, and it is not SM-targeted.

Genuinely PSC-derived or generic (7 constraints). C_1 (Dimensional, $d = 4$), C_6 (Kähler Structure), C_7 (Area Law), C_8 (Unitary Evolution), C_{10} (Einstein Equations), C_{12} (Information Profit), and C_{13} (Necessary Observers) do not directly reference SM field content in their evaluation criteria.

In summary: **5 of 14 constraints** ($C_2, C_3, C_5, C_9, C_{11}$) are SM-targeted to varying degrees; 1 (C_{14}) uses a structurally-fit value; 1 (C_4) encodes a UGP-derived prediction; and 7 are genuinely generic or PSC-derived.

This fuller accounting does *not* reduce the concordance to a tautology, for the following reasons. First, the 7 genuinely generic constraints converge on the SM as their minimum independently; this is a non-trivial result not pre-loaded into the scan. Second, the hard PSC filters that select the 12 surviving universes do not include C_2, C_3, C_5, C_9 , or C_{11} ; the hard filters are C_1, C_6, C_8, C_{12} , and C_{13} , none of which directly encode $G = G_{\text{SM}}$. Third, the concordance’s primary value is not the absolute independence of every term, but the demonstration that SM structure emerges as the unique minimum across a functionally diverse 14-dimensional penalty landscape.

Constraint ablation study (v2: principled C5/C9/C11). Three constraints previously implemented as `is_sm_like()` shortcuts have been replaced with principled group-theoretic computations:

C_{5p} : One-loop beta-function RG stability — penalises theories with Landau poles below M_{GUT} using

gauge-group-specific b_0 coefficients, with no reference to G_{SM} .

C_{9p} : Gauge and gravitational anomaly cancellation — computes $\sum Y^3 = 0$ and $\sum Y = 0$ in the all-left-handed Weyl basis for each gauge group; U(1) alone fails (gravitational anomaly $A = -n_{\text{gen}} \neq 0$), $SU(3) \times SU(2) \times U(1)$ and $SO(10)$ pass, $SU(5)$ passes.

C_{11p} : Witten global $SU(2)$ anomaly and gravitational chiral balance — counts $SU(2)$ doublets per generation and requires parity; $SU(5)$ with $n_{\text{gen}} = 3$ has $5 \times 3 = 15$ doublets (odd, fails); the SM has $4 \times 3 = 12$ (even, passes); $SO(10)$ has $8 \times 3 = 24$ (even, passes).

A pre-committed ablation study
(`comp_p14_constraint_ablation.json`) runs
the 34,560-universe scan under five conditions:

A0 (legacy): All 14 constraints with `is_sm_like()` shortcuts: SM rank 1.

p (principled): All 14 with $C_{5p}/C_{9p}/C_{11p}$ replacing shortcuts: SM rank 1 (passes).

A1: Remove C_2 and C_3 (SRRG fixed-point, the 2 most explicitly targeted); principled $C_{5p}/C_{9p}/C_{11p}$ remain: **SM falls to rank 193**. Top universe: $SU(2)$ with $n_{\text{gen}} = 4$ at $D = 0$.

A2: Principled-only, 11 constraints, no C_{14} : SM rank 577.

A3: Only 3 genuinely UGP-derived (C_1, C_7, C_8): SM rank 289.

A0p passes (principled all-14: SM is rank-1), but **A1 fails**.

This ablation confirms, and sharpens, the earlier disclosure: the C_2/C_3 SRRG fixed-point constraints are the load-bearing element for SM selection over anomaly-free alternatives ($SO(10)$, E_6 , $SU(2)$ with 4 generations, etc.). Crucially, C_2/C_3 are *not* arbitrary SM-targeted inputs: they encode the SRRG theoretical result that the SM is the unique renormalization-group fixed point among all gauge group structures, derived within the MFRR framework [6]. Their current implementation uses the SM as a shortcut to evaluate this claim; replacing them with a principled computation of the SRRG flow for each gauge group is the identified open task.

The C_{9p} and C_{11p} replacements succeed in discriminating U(1) alone (anomalous) and $SU(5)$ (Witten anomaly for $n_{\text{gen}} = 3$) without any reference to G_{SM} — this closes the most circular part of the previous analysis.

We recommend that future versions implement principled SRRG flow computations for C_2/C_3 (checking convergence to a fixed-point for each gauge group via the SRRG beta function), and that C_{14} be revisited once an analytic derivation of Λ_{PSC} from PSC axioms has been established.

4.2 The Concordance Table

Table 3 presents a structured comparison of the key claims of the two methods.

4.3 Why the Agreement Strengthens the Case

The formal and computational results could *a priori* have diverged in several ways. Even with C_2 – C_3 encoding the SM as the SRRG fixed point, the remaining twelve constraints could have assigned lower dissonance to a non-SM configuration—for instance, by finding that $d \neq 4$ compensates for a gauge mismatch, or that $\Lambda = 0$ dominates. Alternatively, the hard PSC filters (which do not include C_2 or C_3) could have passed non-SM universes. Neither happened: all 12 PSC-passing universes have $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$, and the global minimum is achieved at Ψ_{SM} .

This agreement—across both the C_2/C_3 -influenced soft penalties and the C_2/C_3 -independent hard filters—is evidence that the PSC conditions, whether implemented as typed formal predicates or as numerical violation norms, mutually reinforce each other in pointing toward the SM structure. The multi-constraint convergence is the core of the concordance.

Several specific concordance points deserve emphasis.

The $d = 4$ selection. Dimensionally, both the formal exclusion of $d \neq 4$ via RC (Paper 03) and the numerical dominance of $d = 4$ in the dissonance functional (constraint C_1) agree. The formal argument is deductive within the stated domain (4D renormalizable gauge QFT); the numerical argument finds that any $d \neq 4$ universe incurs a dissonance penalty that pushes it far above D_{SM} .

Table 3: Theory–computation concordance: key claims and their status on each side. $\dagger$ marks computational evidence that involves constraints C_2 or C_3 , which partially encode the SM SRRG fixed-point as a prior (see §4.1); $\ddagger$ marks claims where the Residual Classification was previously a conditional (now a theorem: `PSC.RCCInfiniteFamilies`, zero sorry).

Claim	Formal status	Reference	Computational status	TE2.2 evidence
$d = 4$ is selected	Theorem (RC+NM*)	[9]	Global min at $d = 4$	C_1 (independent)
$G = G_{\text{SM}}$ is selected $\dagger$	Theorem (Layer I) $\ddagger$	[6]	All PSC-survivors have G_{SM}	$C_2^\dagger, C_3^\dagger$; hard filter C_8
$N_{\text{gen}} \geq 3$	Theorem (Layer I)	[6]	All PSC-survivors have $N_{\text{gen}} = 3$	$C_2^\dagger, C_{12}$
$N_{\text{gen}} = 3$ exactly	Theorem (Layer II/MDL)	[6]	Unique min at $N_{\text{gen}} = 3$	$C_2^\dagger, C_{12}$ (independent)
GUT groups excluded	Theorem	[9]	GUT-group universes have high D	$C_2^\dagger, C_3^\dagger, C_5$
Vector-like fermions excluded	Theorem	[9]	Not directly parameterized	(Scope gap)
CP-conserving excluded	Theorem	[9]	Captured by C_8 (hard filter)	C_8 (independent)
$\Lambda \approx 10^{-122}$ preferred	Not yet formal	—	D gap of 10 vs $\Lambda = 0$	C_{14} (independent)
$(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ is unique min family $\ddagger$	Conditional on RCC $\ddagger$	[10]	4 co-minimizers at D_{min} ; all SM-like	Global rank, degeneracy in ρ, τ
PSC is rare filter ($< 0.1\%$)	Implied by exclusions	[6, 9]	12/20,160 = 0.06%	Hard filter count
All PSC-survivors are SM-like $\ddagger$	Conditional on RCC $\ddagger$	[10]	All 12 are SM-like	Direct enumeration

A principled derivation of $d = 4$ from PSC axioms independently of the gauge-theory domain was analysed in the TE₁.Z Minimality Theorem experiment. The TE₁.Z analysis establishes that $d = 4$ minimises the PSC dissonance functional under three independent structural criteria: holographic sufficiency (boundary capacity sufficient to encode bulk states), adjudication connectivity (constraint propagation converges before pile-up), and Kähler/symplectic structure (allowing both massless gauge bosons and stable solitons). Each criterion is violated in $d < 4$ (insufficient boundary capacity, graph diameter too large) and the excess overhead of $d > 4$ (extra degrees of freedom without benefit) raises the dissonance. This does not yet constitute a machine-checked proof that $d = 4$ is forced by PSC axioms independent of the gauge-theory domain restriction; it is an analytic argument documenting that $d = 4$ minimises the reflexive ground problem within the PSC framework. A fully independent derivation of the dimensionality from first principles is an open target of the TE₂ programme.

The G_{SM} selection. Both methods agree that G_{SM} is the PSC-preferred gauge group among those in the scan grid. Formally, GUT groups and non-chiral structures are excluded by Theorem 2.4 via NM*; full uniqueness among all compact semisimple gauge groups follows from the RCC theorem (`PSC.RCCInfiniteFamilies`, zero sorry, [12]). Computationally, all non-SM gauge groups in the scan grid are rejected by the hard PSC filters. We note that while C_2 encodes the SM as the SRRG fixed-point and contributes to non-SM penalties, the

hard filter rejection of non-SM gauge groups is driven by the C_2/C_3 -independent constraints C_8 (unitarity/CP) and the dimensional hard filter C_1 —it is not solely an artefact of C_2 ’s circular definition.

The $N_{\text{gen}} = 3$ selection. Both methods agree on three generations. Formally, the Layer I condition gives $N_{\text{gen}} \geq 3$ and Layer II/MDL gives $N_{\text{gen}} = 3$. Computationally, the SRRG fixed-point constraint C_2 is minimized exactly at $N_{\text{gen}} = 3$ for all PSC-passing universes.

The cosmological constant. Here the two methods reveal complementary information. The formal theory does not yet have a complete derivation of $\Lambda \approx 10^{-122}$ from PSC alone (this is an open problem). The computational scan, however, finds a dramatic dissonance gap: universes with $\Lambda = 0$ have $D \approx 11.07$, while those with $\Lambda = 10^{-122}$ have $D \approx 1.07$ —a factor of ≈ 10 difference driven entirely by constraint C_{14} . The numerical evidence is therefore stronger in this direction than the formal evidence.

PSC rarity. The formal exclusion results imply that PSC is highly constraining, but do not by themselves give a numerical rarity estimate. The computational scan provides a concrete number: only 0.06% of the enumerated universes pass the hard PSC filters. This quantifies the filtering power of PSC in a way that complements the formal exclusions.

4.4 What the Concordance Establishes

The concordance supports the following combined conclusion.

Proposition 4.1 (Concordance Conclusion). *Under the Two-Layer PSC Theorem (now unconditional: `RCC` is a Lean-certified theorem, `PSC.RCCInfiniteFamilies`, zero sorry [12]) and the TE2.2 computational certificate, the Standard Model gauge-structure-and-generation tuple $(d, G, N_{\text{gen}}) = (4, G_{\text{SM}}, 3)$ is the uniquely PSC-compatible structure in the following sense:*

- (i) **Formally:** *it is the unique theory surviving all Layer I structural constraints and the Layer II/MDL generation selection (now unconditional (`RCC` established as theorem)).*
- (ii) **Computationally:** *the (d, G, N_{gen}) tuple $(4, G_{\text{SM}}, 3)$ is the unique co-minimum family of $D[\Psi]$ over the 20,160-universe discrete parameterization; all PSC-passing candidates are SM-like; four co-minimizers at $D_{\text{min}} = 1.0667$ exist, differing only in ρ and τ within the SM family.*
- (iii) **Concordantly:** *both methods, using largely independent implementations, identify the same (d, G, N_{gen}) tuple as uniquely PSC-preferred.*

We emphasize that Proposition 4.1 is a theorem: the formal result is unconditional (`RCC` is a Lean-certified theorem) and the computational result is a finite certificate. Together they provide strong evidence for SM necessity that is now unconditional on the `RCC` side.

5 Residuals and Controlled Tensions

We now characterize the principal residuals—the gaps between current evidence and an unconditional claim of SM necessity—and the controlled tensions between the two methods.

5.1 Residual Classification (`RCC`)

Definition 2.10 is supported by computational certificates and by Lean-certified classification of the infinite classical families [12]. The argument has three components:

1. **TE2.2 finite certificate:** 34,560 universes (12 gauge groups) scanned; all PSC-passing universes carry the SM signature (Lean-certified in `TE22.ScanCertificate`).
2. **Extended computational certificate:** 11 additional gauge groups (`E7`, `E8`, `F4`, `SO(12–18)`, `SU(7–10)`) all fail PSC. `E7`, `E8`, `F4`, `SO(16)` fail Layer I by the Lie-algebraic theorem: no complex representations $\Rightarrow$ no chiral fermions in 4D.
3. **Infinite classical families (Lean-certified):** All four infinite families fail by analytic argument (`PSC.RCCInfiniteFamilies`, zero sorry, [12]):
 - $B_n = \text{SO}(2n+1)$, all n : longest Weyl element $w_0 = -\text{id} \Rightarrow$ all irreps self-dual $\Rightarrow$ Layer I fail.
 - $C_n = \text{Sp}(2n)$, all n : same argument.
 - D_n , n even: $-\text{id} \in W(D_n) \Rightarrow$ same.
 - D_n , n odd ≥ 5 : spinor $\dim = 2^{n-1} \geq 16 \Rightarrow$ Layer II.
 - $A_n = \text{SU}(n+1)$, $n \geq 3$: adj $\dim \geq 15 \Rightarrow$ Layer II.

Cases A_1, A_2, D_3, D_5 (i.e., $\text{SU}(2), \text{SU}(3), \text{SU}(4), \text{SO}(10)$) are covered by the TE2.2 finite scan. The Standard Model gauge group $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ is the *unique* compact simple (or semi-simple rank-3) gauge group that passes both PSC layers in 4D with 3 generations.

5.2 The Discretization Gap

The TE2.2 computational certificate is, by construction, a finite discrete enumeration. The extension to the continuum relies on density, continuity, and compactness arguments that are documented analytically in [4] but not formally machine-checked.

The principal uncertainty is whether the discrete grid is truly representative of the full theory space. Specifically:

- The TE2.2-style universe grid is finite; coverage of non-SM gauge structure is therefore certified at the discretized parameterization level, then extended by enlarged group lists and analytic closure over all compact simple Lie algebras [12] (§5.1).
- The generation count N_{gen} ranges from 1 to 4; the scan provides no direct evidence about $N_{\text{gen}} \geq 5$.

- Continuous parameters (Λ, ρ, κ) are discretized; the behavior between grid points is not directly certified.

These limitations are controlled in the sense that the analytic extension argument provides a basis for extrapolation, but they are not eliminated.

5.3 The Neutrino Sector

Neither the formal programme nor the TE2.2 scan directly addresses the neutrino sector. Both operate at the level of gauge group and generation count; neither derives whether neutrinos are Dirac or Majorana, the mass hierarchy, or the CP-violating phase in the PMNS matrix. This is an explicit and acknowledged scope limitation. The concordance established in this paper is at the level of *gauge structure and generation count*; the neutrino sector represents a further frontier for both methods.

Structural neutrino prediction. A structural prediction of the neutrino mass-squared ratio is given in the companion paper [3]. The Braid Atlas right-handed neutrino b -values $\{5, 11, 19\}$ satisfy $m_{\nu_g} \propto b_g^{29/9}$; the exponent admits three independent structural decompositions:

$$\frac{29}{9} = \frac{N_c^3 + s}{N_c^2} = \frac{4N_c^2 - \delta}{N_c^2} = \frac{d_{45} - d_{16}}{N_c^2},$$

where $s = (N_c^2 - 1)/4 = 2$, $\delta = 7$, and $d_{45} - d_{16} = \dim(45_{\text{SU}(5)}) - \dim(16_{\text{SO}(10)}) = 29$. The formula predicts the mass-squared splitting ratio $\Delta m_{21}^2 / \Delta m_{31}^2 = 0.02936$, matching NuFIT 6.0 [2] (0.02951 ± 0.00098) to 0.16σ (0.52% ; 1.58σ from PDG 2024 world average) with no free parameters. Normal ordering is automatic. With the Dirac scale $E_D = v_H / (4N_c^2 - \delta) = v_H / 29$ (same structural integer as the exponent numerator), the absolute sum is $\sum m_\nu = 56$ meV at $M_R = 2 \times 10^{16}$ GeV (within the Planck bound), and 60 meV at the best-fit $M_R = 1.88 \times 10^{16}$ GeV. The structural neutrino prediction thus covers both the mass-squared ratio and the absolute Planck-window sum; a first-principles Lagrangian derivation combining the SO(10) Yukawa matrix with Braid Atlas flavour structure, and a UGP-internal determination of M_R , remain open.

5.4 Vector-Like Fermion Representations

The formal exclusion of vector-like fermions (Theorem 2.4(ii)) is not directly verified by the TE2.2

scan, because the scan’s parameterization does not include a vector-like vs. chiral parameter. This is a scope gap between the two methods: the formal side makes a claim that the computational side does not independently check. A future TE2.2-type scan that includes fermion chirality as a parameter would close this gap.

5.5 Falsifiability

The concordance result makes concrete falsifiability predictions.

Formal falsifiability. The RCC would be falsified by construction of an explicit gauge theory satisfying all PSC conditions with $G \neq G_{\text{SM}}$ or $N_{\text{gen}} \neq 3$. The Lean 4 formalization [11] provides a machine-checkable sieve: a proposed alternative theory can be submitted to the sieve and its PSC status verified.

Computational falsifiability. The TE2.2 result would be contradicted by:

- A universe description Ψ' with $G \neq G_{\text{SM}}$ or $N_{\text{gen}} \neq 3$ achieving $D[\Psi'] < D_{\text{SM}}$ under the same functional (2);
- A non-SM universe passing all hard PSC filters under the same weights w_i ;
- A finer discretization that reveals non-SM minimizers.

The data file is publicly available [13] and the scan code is reproducible (see Appendix B); any such contradiction would be immediately detectable.

Experimental falsifiability. Both methods would be undermined by experimental evidence for:

- A fourth generation of chiral fermions within the SM gauge structure;
- Observation of a GUT symmetry at accessible energies;
- Detection of vector-like quarks or leptons at the LHC or successor colliders.

These possibilities are physically testable and none has been confirmed to date.

6 Conclusion

What this paper establishes. This paper serves two related functions.

First, it is the computational certificate for the Two-Layer PSC Theorem of Papers 03 and 05 [6, 9]. That theorem *proves* as formal results, within the domain of 4D renormalizable gauge QFT: (i) GUT groups, vector-like fermions, and CP-conserving theories are structurally excluded by PSC; (ii) any remaining self-contained theory must have $G = G_{\text{SM}}$ and $N_{\text{gen}} \geq 3$ (Layer I, forced by consistency); (iii) $N_{\text{gen}} = 3$ is the unique MDL-minimal Layer I survivor (Layer II, optimality). These steps are now unconditional: the Residual Classification (RCC) is established as a Lean-certified theorem (PSC.RCCInfiniteFamilies, zero `sorry`, [12]), with key steps machine-checked in nems-lean [8]. The TE2.2 scan provides the computational analogue: exhaustive evaluation of $D[\Psi]$ over 20,160 discretized theories confirms that the SM is the unique minimizer at the (d, G, N_{gen}) level, with all 12 PSC-passing universes (0.06%) being SM-like.

Second, this paper strengthens the formal case by contributing three new UGP-derived coupling-ratio predictions (C_4, C_{15}, C_{16}) that are satisfied by the SM to within 2–7% after RG running, derived entirely from ugp-lean arithmetic invariants—not from SM coupling data. The entire $D_{\text{SM}} = 1.0667$ residual is carried by C_4 (Quarter-Lock), encoding the 5% deviation of $\sqrt{3}g_1 \approx g_2$ at M_Z from the UGP-predicted ratio; all other constraints evaluate to zero at the SM point. An extended scan over 12 gauge groups—including Pati-Salam $SU(4) \times SU(2) \times SU(2)$ and E_6 —confirms that all major BSM candidates fail the PSC sieve.

The forced vs. selected distinction. A key conceptual contribution of this paper is sharpening the distinction between what PSC *forces* and what it *selects*. $N_{\text{gen}} \geq 3$ is *forced* by the Layer I CP-violation requirement (Remark 2.8). $N_{\text{gen}} = 3$ is *selected* by the Layer II MDL criterion. G_{SM} is forced by chirality and anomaly cancellation. The continuous cosmological parameters $(\Lambda, \rho, \kappa, \tau)$ remain *underdetermined* within the SM family. This taxonomy is important: the PSC programme achieves structural forcing of the gauge group and generation count floor, optimality selection of the exact generation count, and identifies the remaining underdetermined sector as the target

for future work.

Principal residuals. The RCC (established as a Lean-certified theorem, PSC.RCCInfiniteFamilies, zero `sorry` [12]; SM uniqueness is now unconditional), the discretization gap (controlled but not eliminated), the neutrino sector (mass-squared ratio predicted to 0.16σ from NuFIT 6.0 (0.52%); absolute sum within Planck bound via $E_D = v_H/29$; exponent 29/9 identified via three independent structural decompositions; full Lagrangian derivation remains open), the absence of a complete Lean 4 proof of the Phase 3 continuum extension, and the algebraic fit of C_{14} to Λ_{obs} (not yet independently derived), and the vacuum sector (the Strong CP phase $\theta_{\text{QCD}} = 0$ is argued conditionally in [3] but is not established as a PSC-forced result). Each represents a well-defined target for future work.

Working hypothesis. The Standard Model is not merely an empirically successful theory fitted to data, but reflects a privileged self-consistent structure under strong information-theoretic, algebraic, and geometrodynamical closure conditions. Whether this amounts to a proof of SM *necessity* requires an independent analytic derivation of C_{14} ; the RCC has now been resolved (PSC.RCCInfiniteFamilies, zero `sorry`, [12]). The present evidence—formal theorem chain, computational certificate, three new UGP-derived coupling predictions, and extended BSM exclusion scan—constitutes the strongest current case for that hypothesis.

Acknowledgments. The TE2.2 scan was implemented and executed in the ugp-physics repository [13]. The Lean 4 formalization is archived at [11].

A The Fourteen TE2.2 Constraint Terms

We give the analytic definition of each constraint term $C_i[\Psi]$ used in the dissonance functional (2). In all cases $\Psi = (d, G, N_{\text{gen}}, N_{\text{obs}}, \Lambda, \rho, \kappa, \tau)$ is the universe parameter vector.

C_1 : Dimensional. $\|C_1[\Psi]\|^2 = (d - 4)^2$. This penalizes deviation from the PSC-optimal spacetime dimension $d = 4$.

C_2 : SRRG Fixed Point. $\|C_2[\Psi]\|^2 = 1[G \neq G_{\text{SM}}] + (N_{\text{gen}} - 3)^2$. This penalizes departure from the self-renewing renormalization group (SRRG) fixed point $(G_{\text{SM}}, N_{\text{gen}} = 3)$.

C_3 : SRRG Viability. $\|C_3[\Psi]\|^2 = \max(0, F_{\text{max}}[S] - F[S; \Psi])^2$, where $F[S; \Psi]$ is the gauge coupling functional and $F_{\text{max}}[S]$ is its maximum over PSC-compatible structures. This penalizes failure to achieve maximal SRRG viability.

C_4 : Quarter-Lock. $\|C_4[\Psi]\|^2 = (\sqrt{3}g_1 - g_2)^2$, evaluated at the SRRG fixed point. This encodes the gauge-coupling ratio condition required for the SRRG structure to be self-sustaining.

C_5 : RG Flow Stability. $\|C_5[\Psi]\|^2 = \max(0, dc/dt)^2$, where c is the running coupling. This penalizes positive beta functions (growing couplings in the UV).

C_6 : Kähler Structure. $\|C_6[\Psi]\|^2 = \|g_{\text{Fisher}} - \omega_{\text{symp}}\|_F^2$, where g_{Fisher} is the Fisher information metric on coupling-constant space and ω_{symp} is the natural symplectic form. This penalizes failure of the Fisher metric to be Kähler (i.e., compatible with the symplectic structure).

C_7 : Area Law. $\|C_7[\Psi]\|^2 = (S_{\text{EE}} - A/(4\ell_P^2) - \beta \log A)^2$, where S_{EE} is the entanglement entropy of the vacuum state and A is the area of the causal horizon. This penalizes deviation from the Bekenstein–Hawking area law with logarithmic correction.

C_8 : Unitary Evolution. $\|C_8[\Psi]\|^2 = 1[\text{CP-violation absent}]$. A CP-conserving theory cannot maintain unitary time evolution in the PSC framework; this constraint is a hard binary filter.

C_9 : RIET Equivalence. $\|C_9[\Psi]\|^2 = \|\nabla_\mu R - \nabla_\mu T - \nabla_\mu I - \nabla_\mu C\|^2$, penalizing failure of the mutual duality between spacetime curvature (R), energy-momentum (T), entropy (I), and information processing rate (C) that characterizes Reflexive Information-Energy-Topology equivalence.

C_{10} : Einstein Equations. $\|C_{10}[\Psi]\|^2 = \|G_{\mu\nu} - 8\pi G T_{\mu\nu}\|^2$, evaluated on the ground state of Ψ . This penalizes deviation from general-relativistic dynamical equations.

C_{11} : Coherence Field. $\|C_{11}[\Psi]\|^2 = \|\delta S_{\text{coh}}/\delta\Psi\|^2$, where S_{coh} is the coherence field action. This penalizes inconsistent coupling of the coherence field Ψ to the matter sector.

C_{12} : Information Profit. $\|C_{12}[\Psi]\|^2 = \max(0, 1.13 - \rho)^2$. This is a hard threshold: $\rho < 1.13$ (information generation rate less than drain rate) violates the PSC information-viability condition.

C_{13} : Necessary Observers. $\|C_{13}[\Psi]\|^2 = \max(0, 1 - N_{\text{obs}})^2$. PSC requires at least one observer class; universes with $N_{\text{obs}} < 1$ are PSC-inviable.

C_{14} : Lambda Relation. $\|C_{14}[\Psi]\|^2 = (\log_{10} \Lambda - \log_{10}(\ln \varphi / \ln 2\pi))^2$, where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio. The PSC-predicted value is $\Lambda_{\text{PSC}} = \ln \varphi / \ln 2\pi \approx 10^{-122}$ in Planck units; this constraint penalizes deviation from that value on a logarithmic scale.

The weights w_i are set so that each active constraint contributes approximately equally at the PSC boundary. The exact values are documented in the TE2.2 code base [4, 13].

B Reproducibility

Data location. The TE2.2 scan results are archived in the `ugp-physics` repository [13] at path:

```
MFRR/TE_2_Advanced_Explorations/
TE_2_2_Minimal_PSC_Universe/results/
phase2_scan_results.json
```

Data integrity. The canonical data file has SHA-256 hash:

```
f810c1d2b07b598ef301205fee53512310552ea7
8cf8fb7476b3e9058d5fde93
```

Key fields. The JSON file contains the following key fields relevant to the results reported in this paper:

- `total_universes`: 20,160
- `psc_universes`: 12
- `D_sm`: 1.066657903568035
- `D_min`: 1.066657903568035
- `sm_rank`: 1
- `elapsed_seconds`: 0.13975...
- `global_minimizer`:
`{d:4, G:"SU(3)xSU(2)xU(1)", n_gen:3,...}`

Code. The scan code is available in the same repository under `MFR/TE_2_Advanced_Explorations/TE_2_2_Minimal_PSC_Universe/`. The primary modules are:

1. `te2_2_constraint_base.py`
2. `te2_2_dimensional_constraint.py`
3. `te2_2_srrg_constraint.py`
4. `te2_2_remaining_constraints.py`
5. `te2_2_constraint_aggregator.py`
6. `te2_2_universe_enumerator.py`
7. `te2_2_run_scan.py`

Lean formalization. The Lean 4 machine-checked formalization of the formal PSC constraints is available at [11] (DOI: 10.5281/zenodo.19433538).

How to reproduce. To reproduce the TE2.2 scan:

1. Clone `ugp-physics` [13].
2. Install Python 3.9+ and `numpy`.
3. Run `python te2_2_run_scan.py` from the `TE_2_2_Minimal_PSC_Universe/src/phase2_truncation/` directory.
4. Verify the output JSON against the SHA-256 hash above.

Expected runtime: < 1 second on modern hardware.

C Lean Formalization of the PSC Gauge Sieve

The `nems-lean` library (Zenodo 10.5281/zenodo.19429710, GitHub <https://github.com/novaspiavack/nems-lean>) provides a machine-checked formalization of the PSC gauge sieve results cited in this paper. The build command is `lake update && lake build` from the `nems-lean` root. All results are proved with zero `sorry` and zero custom axioms.

NemS.Physics.GaugeTheorySpace.gauge_signature_rigidity (NEMS Paper 20, Zenodo 10.5281/zenodo.19575045). *Statement:* The `ResidualClassificationConjecture` on which this theorem was conditioned is now proved as a theorem (`PSC.RCCInfiniteFamilies`, zero `sorry`, [12]); every PSC-admissible gauge theory satisfies `isSMGaugeGroup` $\wedge$ $N_{\text{gen}} = 3$ unconditionally. *Key predicates:* `RC`, `NM*`, `SA`, `TV`, `AnomalyConsistent`, `SMSignature` are real Lean predicates (not `True` stubs); `ResidualClassification` carries the consolidated certificates of Section 5.1 [12]. *Relationship to this paper:* This theorem provides the machine-checked backing for the concordance claim that the formal PSC sieve selects G_{SM} and $N_{\text{gen}} = 3$. The conditional structure (on `RCC`) has been resolved; §2 and the concordance table (§4.2) mark formerly-conditional claims with ‡ for historical reference.

References

- [1] John C. Baez and John Huerta. The algebra of grand unified theories. *Bulletin of the American Mathematical Society*, 47(3):483–552, 2010. 2013 Levi L. Conant Prize, AMS. Algebraic analysis of GUT and SM gauge group structure.
- [2] I. Esteban, M.C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J.P. Pinheiro, and T. Schwetz. NuFit-6.0: updated global analysis of three-flavor neutrino oscillations. *JHEP*, 12:216, 2024. NuFIT 6.0 global fit (2024), <http://www.nu-fit.org/>.
- [3] N. Spivack. A deterministic number-theoretic framework for the standard model parameter spectrum. Zenodo, 2026. Paper P01 in

the UGP Physics series. <https://doi.org/10.5281/zenodo.20168787>.

- [4] Nova Spivack. The TE2.2 minimal PSC universe scan: Finite exhaustive enumeration of PSC-optimal universe structures. Computational result archived in ugp-physics repository, 2025. 20,160-universe exhaustive enumeration; Standard Model ranks #1; 12 of 20,160 pass hard PSC filters; all 12 are SM-like. See the ugp-physics repository.
- [5] Nova Spivack. From NEMS to MFRR: A machine-checked bridge between semantic closure and reflexive reality. NEMS/PSC Suite, Paper 08, 2026. Lean-verified bridge theorems and diagonal barrier; zero custom axioms.
- [6] Nova Spivack. PSC-optimality of the standard model in 4D gauge theory space. NEMS/PSC Suite, Paper 05, 2026.
- [7] Nova Spivack. Reflexive reality: A survey for physicists. NEMS/PSC Suite, Paper 88, 2026.
- [8] Nova Spivack. Rigidity of the gauge signature under PSC: A sieve theorem. NEMS/PSC Suite, Paper 20, 2026.
- [9] Nova Spivack. Structural stability as a necessity principle in self-contained gauge theories. NEMS/PSC Suite, Paper 03, 2026. <https://doi.org/10.5281/zenodo.19429717>. Derives the NM^* structural stability constraint from PSC; excludes GUT groups, vector-like fermions, and CP-conserving theories. Provides the gauge-theory-space filter used in the TE2.2 scan.
- [10] Nova Spivack. The theorem of existential rigidity: The collapse of contingency. NEMS/PSC Suite, Paper 21, 2026. Proves that if the Residual Classification Conjecture holds, the Standard Model signature is the only ontologically legal foundation for a universe.
- [11] Nova Spivack. ugp-lean: A machine-checked formalization of the universal generative principle in Lean 4. Zenodo, 2026. Formalization paper (latest version): <https://doi.org/10.5281/zenodo.19554688>. Concept DOI (always-latest): <https://doi.org/10.5281/zenodo.19433538>. 117 modules across twelve architectural layers including GTE, Universality, ElegantKernel (UCL unconditional closure), MassRelations (Koide closed form, S_3 -Newton flow, binary cascade, physical mass spectrum), BraidAtlas (ChargeTheorem, CompositeTriples, ChiralitySquaring, ChargeDerivation, CoxeterConductor, CoxeterConductor-TowerLaw, EWBosons), GaloisStructure, PSC RCC infinite-family closure, and TE2.2 scan certification. Zero sorry, one disclosed axiom (`mirror_winding_number_zero` for GTE-P7 dark matter, contained to mirror sector only). Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [12] Nova Spivack. ugp-lean: Lean 4 formalization of UGP and GTE. <https://github.com/novaspivack/ugp-lean>, 2026. 117-module Lean 4 library (module count updated as of 2026-05-09). `lake build to verify` (zero sorry, zero custom axioms, standard Mathlib axiom signature). Includes MassRelations, ElegantKernel unconditional closure, BraidAtlas composites and charge derivation, PSC RCC infinite-family layer, TE2.2 scan certification. New in GTE.GeneralTheorems: E8 cyclotomic universality theorems (all 8 Zamolodchikov E8 mass ratios lie in $\mathbb{Q}(\zeta_{120})$, `e8_all_masses_divisibility`, `e8_cyclotomic_divisibility`). Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [13] Nova Spivack. ugp-physics: Companion code and papers for the UGP physics series. <https://github.com/novaspivack/ugp-physics>, 2026. Includes `computational_concordance/` tables, basin JSON exports, and capstone L^AT_EX.

Acknowledgments

AI coding assistants were used for LaTeX formatting, coding assistance, and adversarial review during manuscript preparation. All mathematical derivations, numerical computations, and Lean 4 certifications were independently verified by deterministic scripts and the Lean 4 kernel.