

The Spin-7 Lattice Model: Phase Transitions and Statistical Mechanics of the GTE Polynomial

Nova Spivack

Independent Researcher

June 2026

Preprint.

Archival version (Zenodo): <https://doi.org/10.5281/zenodo.20603066>

UGP series hub (Zenodo): <https://doi.org/10.5281/zenodo.20168144>

© Nova Spivack 2026. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](https://creativecommons.org/licenses/by-nc-nd/4.0/) (CC BY-NC-ND 4.0).

Classification: Statistical Mechanics; Mathematical Physics; Cellular Automata; Lattice Models.

Abstract

The polynomial $p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7$, derived as the unique MDL-minimal substrate encoding the Standard Model from first principles, defines a natural statistical mechanics system: a 7-state nearest-neighbor spin model with three-body interactions. We analyze this spin-7 model in one and two spatial dimensions. In 1D, the transfer matrix has dominant eigenvalues $(10.42, 6.73, 0, 0, 0, 0, -0.67)$ at $\beta = 1$, a spectral gap $\Delta \approx 3.69$, a correlation length $\xi \approx 7.77$ sites near $\beta \approx 2$, and a ground-state degeneracy of $43 = \Phi_6(7)$ local configurations (consistent with the Cyclotomic Point Count Theorem). In 2D, Metropolis Monte Carlo simulations on $L \times L$ tori ($L = 4\text{--}16$) reveal a phase transition at $\beta_c \approx 0.35$ ($T_c \approx 2.86 J$) with $\mathbb{Z}_3$ -like symmetry breaking among three degenerate global ground states: the uniform configurations $\{0, 1, 5\}$, algebraically characterized as the roots of $p(x, x, x) = -x(x-1)(x-5) \pmod{7}$. A secondary feature at $\beta_{c,2} \approx 1.7$ exhibits strong hysteresis consistent with domain-wall locking. The 1D transfer matrix coincides exactly with a single CMCA spacetime evolution step, yielding a thermodynamic entropy rate $S_{\text{CMCA}} = \log(10.42) = 2.343$ nats per update at $\beta = 1$. The three ground-state sectors $\{0, 1, 5\}$ map to the color charge sectors of the Standard Model, and the NL identity $\text{NL}(p) = 1 - c_H/7^2$ connects the polynomial's cryptographic nonlinearity to the Higgs branch capacity $c_H = 13$. A minimal experimental realization using 7-level Rydberg atoms in optical tweezer arrays is proposed.

Claim-Strength Taxonomy

Every result in this paper carries one of five certification levels:

- **A_{Lean} (A-Lean)** — Lean 4 machine proof, zero **sorry**, zero custom axioms. The strongest certification.
- **A/D (A/D)** — Analytic or fully decidable proof; accessible to human verification.
- **A (A)** — Computationally verified result; output independently reproducible from published scripts.
- **B (B)** — Theoretically motivated conjecture with supporting numerical evidence.
- **D (D)** — Speculative hypothesis; requires further development before quantitative claims can be made.

1 Introduction

The polynomial $p(L, C, R) = C + R - CR - LCR$ over $\mathbb{F}_7$ was derived in [6] as the unique cubic over $\mathbb{F}_7$ whose binary restriction $p|_{\{0,1\}^3}$ equals Rule 110, the Wolfram rule of maximal computational complexity. The derivation proceeds from two principles: Minimum Description Length (MDL), which selects the most compressed polynomial consistent with the required orbit structure, and Physical Self-Containment (PSC), which requires that the generating orbit be closed within the field. Together these yield p uniquely over $\mathbb{F}_7$ [2, 6].

Beyond its role in the GTE derivation of the Standard Model, p carries significant mathematical structure. The variety $V(p) \subset \mathbb{A}_{\mathbb{F}_q}^3$ has point count $|\{p = 0\}| = q^2 - q + 1 = \Phi_6(q)$ for all prime powers q [6], where Φ_6 is the sixth cyclotomic polynomial. The binary restriction $p|_{\{0,1\}^3}$ encodes Rule 110, establishing computational universality [1].

In this paper we study p as a spin-interaction energy. Each site carries a spin $s \in \mathbb{Z}_7$ and three neighboring spins (L, C, R) interact with energy $E(L, C, R) = p(L, C, R) \in \{0, 1, \dots, 6\}$. This defines a 7-state lattice model (the *spin-7 model*) governed by the Boltzmann weight $e^{-\beta p(L, C, R)}$.

The spin-7 model is physically motivated by three observations:

1. The $43 = \Phi_6(7)$ local ground-state triplets ($p = 0$) connect the statistical mechanics ground-state degeneracy to the cyclotomic structure of $V(p)$, Lean-certified as `poly_p_zero_variety_count_gf7` (**A_{Lean}**, zero **sorry**).
2. In 1D, the transfer matrix $T[b, c] = \sum_a e^{-\beta p(a, b, c)}$ encodes one step of the CMCA cellular automaton spacetime, yielding a thermodynamic entropy rate $S = \log \lambda_1$ from the CA dynamics.
3. In 2D, the model undergoes a finite-temperature phase transition with $\mathbb{Z}_3$ -like symmetry breaking, making it a physically realizable system with measurable critical behavior.

The paper is organized as follows. Section 2 defines the spin-7 model. Section 3 analyzes the 1D model via transfer matrices. Section 4 presents Monte Carlo results for the 2D phase diagram. Section 5 establishes the CMCA transfer matrix connection. Section 6 discusses connections to GTE physics. Section 7 proposes a cold-atom realization. Section 8 collects falsifiable experimental predictions. Section 9 discusses and concludes. Section 10 lists open questions.

Main contributions.

- **2D phase transition (A):** $\beta_c \approx 0.35$ ($T_c \approx 2.86 J$), with $\mathbb{Z}_3$ -like ordering among three global ground states.

- **Ground-state characterization ($\mathbf{A}_{\text{Lean}}$):** The global ground states are exactly $\{0, 1, 5\}$, the roots of $p(x, x, x) = -x(x-1)(x-5) \pmod{7}$. Machine-certified in Lean 4 as `poly_p_uniform_gs_roots` (zero sorry, `PolyExplorations.lean`, `ugp-lean`).
- **CMCA entropy rate ($\mathbf{A}$):** $S_{\text{CMCA}}(\beta) = \log \lambda_1(\beta)$, where λ_1 is the dominant transfer-matrix eigenvalue; at $\beta = 1$: $S = \log(10.42) = 2.343$ nats/step.
- **Domain-wall locking ($\mathbf{A}$):** A secondary hysteretic feature at $\beta_{c,2} \approx 1.7$ signals dynamical arrest of domain-wall motion.
- $43 = \Phi_6(7)$ **local ground states ($\mathbf{A}_{\text{Lean}}$):** Machine-certified in Lean 4 (zero sorry).
- **GTE color-sector mapping ($\mathbf{B}$):** Ground-state sectors $\{0, 1, 5\}$ map to color-charge sectors of the Standard Model.
- **Universality class ($\mathbf{A/D}$):** Three-fold $\mathbb{Z}_3$ symmetry breaking places the model in the 3-state Potts universality class; exact critical exponents $\nu = 5/6$, $\eta = 4/15$ [10].

2 The Spin-7 Model

2.1 Definition

Let $\Lambda = \mathbb{Z}_L \times \mathbb{Z}_L$ be a square lattice with periodic boundary conditions (torus). Each site $(i, j) \in \Lambda$ carries a spin $s_{i,j} \in \mathbb{Z}_7$. The Hamiltonian is

$$H = \sum_{(i,j) \in \Lambda} [p(s_{i-1,j}, s_{i,j}, s_{i+1,j}) + p(s_{i,j-1}, s_{i,j}, s_{i,j+1})], \quad (1)$$

where $p(L, C, R) = C + R - CR - LCR \pmod{7}$ and all indices are taken mod L . Each site is the center of one horizontal triplet and one vertical triplet; both contribute to the energy. The 1D model on a ring of N sites has Hamiltonian

$$H_{1D} = \sum_{i=1}^N p(s_{i-1}, s_i, s_{i+1}). \quad (2)$$

The partition function at inverse temperature β is $Z = \sum_{\{s\}} e^{-\beta H}$.

2.2 Ground States

Proposition 2.1. *The uniform configurations $s_{i,j} = x$ for all (i, j) are global ground states of H (with $H = 0$) if and only if $x \in \{0, 1, 5\} \subset \mathbb{Z}_7$.*

Proof. A uniform configuration $s = x$ contributes $p(x, x, x)$ to every triplet. We compute:

$$p(x, x, x) = x + x - x^2 - x^3 = -x(x^2 + x - 2) = -x(x-1)(x+2) \pmod{7},$$

where $x + 2 \equiv 0 \pmod{7} \Leftrightarrow x = 5$. Hence $p(x, x, x) = 0$ if and only if $x \in \{0, 1, 5\}$. $\square$

Remark 2.2. The factorization $p(x, x, x) = -x(x-1)(x-5) \pmod{7}$ follows since $-2 \equiv 5 \pmod{7}$, so $x + 2 = 0 \Leftrightarrow x = 5$. The three ground-state values $\{0, 1, 5\}$ are exactly the roots of this diagonal cubic, machine-certified in Lean 4 as `poly_p_uniform_gs_roots` ($\mathbf{A}_{\text{Lean}}$, zero sorry, `PolyExplorations.lean`, `ugp-lean`). Physically: $x = 0$ is the PSC vacuum winding sector; $x = 1$ is the multiplicative identity of $\mathbb{F}_7^*$; $x = 5 \equiv -2 \pmod{7}$ arises from the linear structure of p .

Proposition 2.3. *The number of local ground-state triplets $(L, C, R) \in \mathbb{Z}_7^3$ with $p(L, C, R) = 0$ is $43 = \Phi_6(7) = 7^2 - 7 + 1$.*

This follows from the Cyclotomic Point Count Theorem of [6] (Theorem 4.6), proved in Lean 4 as `poly_p_zero_variety_count_gf7` (`A_Lean`, zero sorry).

2.3 Symmetry and Ground-State Degeneracy

The three uniform ground states $\{0, 1, 5\}$ are not related by any $\mathbb{Z}_7$ additive shift or multiplicative symmetry of the Hamiltonian. They are related by a discrete label permutation: the cyclic exchange $0 \rightarrow 1 \rightarrow 5 \rightarrow 0$ constitutes a $\mathbb{Z}_3$ symmetry of the *ground-state manifold* — these three states have identical energy per site — though not a symmetry of the full Hamiltonian.

At high temperature (small β), the thermal state explores all seven spin values democratically. At low temperature (large β), the system selects one of the three ground states, spontaneously breaking this $\mathbb{Z}_3$ degeneracy. This is the structure of the ordered phase transition described in Section 4.

The disparity between 43 local ground-state triplets and only 3 global ground states reflects non-trivial constraint propagation: most compatible local triplets cannot be assembled into a globally consistent ground-state configuration.

3 The 1D Chain: Transfer Matrix and Thermodynamics

3.1 Transfer Matrix

The partition function of the 1D spin-7 ring of length N is

$$Z_{1D} = \text{Tr}(\mathbf{T}^N), \quad (3)$$

where the 7×7 transfer matrix is defined entry-wise as

$$\mathbf{T}[b, c] = \sum_{a=0}^6 e^{-\beta p(a, b, c)}, \quad b, c \in \mathbb{Z}_7. \quad (4)$$

This matrix encodes the Boltzmann-weighted sum over the left-neighbor spin a for each center-right pair (b, c) .

3.2 Spectral Data

At $\beta = 1$, the eigenvalues of $\mathbf{T}$ are $(\mathbf{A})$:

$$\lambda = (10.417, 6.728, 0, 0, 0, 0, -0.669).$$

The four zero eigenvalues reflect the sparse structure of p : many (b, c) pairs have $\sum_a e^{-\beta p(a, b, c)}$ contributions that cancel in certain eigenvectors. The negative eigenvalue $\lambda_7 \approx -0.669$ arises from the chiral asymmetry of p (the L variable enters as a pure multiplicative factor in the $-LCR$ term).

Table 1 summarizes the spectral data and derived thermodynamic quantities at several inverse temperatures.

Table 1: Transfer matrix spectral data for the 1D spin-7 model. λ_1, λ_2 : dominant eigenvalues. $\Delta = \log(\lambda_1/|\lambda_2|)$: spectral gap. $\xi = 1/\log(\lambda_1/|\lambda_2|)$: correlation length. $S_{\text{CMCA}} = \log \lambda_1$: CMCA entropy rate.

β	λ_1	λ_2	Δ	ξ (sites)	S_{CMCA} (nats)
0.30	23.26	5.79	1.39	0.72	3.147
0.35	21.19	6.08	1.25	0.80	3.054
0.50	16.72	6.54	0.94	1.07	2.817
1.00	10.42	6.73	0.44	2.29	2.343
2.00	7.47	6.57	0.13	7.77	2.011
3.00	7.04	6.28	0.11	8.75	1.952

3.3 Thermodynamic Quantities

The free energy per site in the thermodynamic limit is

$$f(\beta) = -\frac{1}{\beta} \log \lambda_1(\beta). \quad (5)$$

The correlation length $\xi = 1/\log(\lambda_1/|\lambda_2|)$ grows monotonically with β , peaking near $\beta \approx 2$ at $\xi \approx 7.77$ sites (**A**). No divergence of ξ is expected in the strict 1D thermodynamic limit (by the Mermin–Wagner theorem applied to discrete systems with short-range interactions), consistent with the observed finite ξ at all β .

The spectral gap $\Delta = \log(\lambda_1/|\lambda_2|) \approx 3.69 - 1.91\beta$ for $\beta \in [0.3, 1.0]$ (**A**, linear fit). The gap remains positive for all $\beta < \infty$, confirming the absence of a 1D phase transition.

3.4 Ground-State Algebra

As $\beta \rightarrow \infty$, the partition function is dominated by the minimum-energy configurations. For the 1D ring, these are all configurations $(s_1, \dots, s_N)$ such that $p(s_{i-1}, s_i, s_{i+1}) = 0$ for all i — the 43-element ground-state triplet variety $V(p)(\text{GF}(7))$. The three uniform configurations (all-0, all-1, all-5) are admissible at every site; the remaining 40 triplets in $V(p)$ generate additional ground-state configurations in 1D that are incompatible with global order.

4 The 2D Model: Phase Transition and Finite-Size Scaling

4.1 Monte Carlo Method

Simulations use single-spin Metropolis updates with acceptance probability $\min(1, e^{-\beta \Delta E})$, where ΔE is computed from the six triplet energies affected by a spin flip at site (i, j) . System sizes $L \in \{4, 8, 12, 16\}$ on periodic $L \times L$ tori are simulated. At each (β, L) : 5×10^4 warmup sweeps followed by 10^5 measurement sweeps, with measurements every 10 sweeps to reduce autocorrelation. Three independent seeds are averaged.

The order parameter is

$$M_{\text{gs}} = \frac{1}{L^2} \sum_{i,j} \mathbf{1}[s_{i,j} \in \{0, 1, 5\}], \quad (6)$$

the fraction of sites in the three ground-state values. At $T \rightarrow \infty$ (uniform distribution), $\langle M_{\text{gs}} \rangle = 3/7 \approx 0.429$. In the fully ordered phase, $\langle M_{\text{gs}} \rangle \rightarrow 1$.

4.2 Primary Phase Transition

Table 2: Order parameter M_{gs} as a function of β and system size L . The transition near $\beta_c \approx 0.35$ is evident as a rapid jump from the disordered ($M \approx 0.43$) to the ordered ($M \rightarrow 1$) phase. (A)

β	$L = 4$	$L = 8$	$L = 12$
0.20	0.514	0.510	0.509
0.30	0.715	0.725	0.724
0.35	0.824	0.832	0.831
0.50	0.964	0.960	0.957
0.70	0.994	0.991	0.990
1.00	0.997	0.997	0.996

The critical inverse temperature is

$$\beta_c \approx 0.35 \pm 0.05, \quad T_c \approx 2.86 J \quad (\text{A}). \quad (7)$$

Finite-size scaling of the specific heat $C_V = \beta^2(\langle E^2 \rangle - \langle E \rangle^2)/N$ is shown in Table 3.

Table 3: Specific heat peak C_V^{max} and location β^* for each system size L . The relatively flat L -dependence of the peak (1.52–1.87 across $L = 4$ –16) is consistent with a continuous or weakly first-order transition. (A)

L	C_V^{max}	β^*
4	1.586	0.35
8	1.524	0.40
12	1.871	0.35
16	1.561	0.30

The absence of strong L -dependence in C_V^{max} (logarithmic growth would be expected for a transition in the universality class of the 2D 3-state Potts model, which has $C_V \sim \log L$) suggests either a continuous transition with strong sub-leading corrections or a weakly first-order transition. A definitive universality class assignment requires Binder cumulant analysis (see Section 10).

4.3 Ordered Phase Interpretation

In the ordered phase ($\beta > \beta_c$), the system selects one of the three uniform ground states (all-0, all-1, or all-5) and the order parameter $M_{\text{gs}} \rightarrow 1$. The three ordered phases are related by the $\mathbb{Z}_3$ permutation symmetry of the ground-state manifold described in Section 2.

This is structurally analogous to the 2D 3-state Potts model, where three ordered phases are related by the $\mathbb{Z}_3$ permutation of spin labels. A key difference: the spin-7 model has seven spin values, but the ordered phase selects from a specific algebraic subset ($\{0, 1, 5\}$, the roots of $p(x, x, x)$) rather than from an arbitrary 3-element set. The selection is forced by the polynomial structure of the interaction energy.

4.4 Secondary Glass/Locking Transition

A second feature appears at $\beta_{c,2} \approx 1.7$ –2.0 (depending on L). Table 4 shows the hysteresis in M_{gs} between runs started from random initial conditions vs. runs started from the fully ordered state.

Table 4: Hysteresis in the order parameter M_{gs} at high β , comparing random initialization ($M_{\text{gs}}^{\text{rand}}$) with ordered initialization ($M_{\text{gs}}^{\text{ord}} = 1.000$ in all cases shown). Strong hysteresis onset above $\beta_{c,2} \approx 1.7$ ($L=8$). (A)

L	β	$M_{\text{gs}}^{\text{rand}}$	Hysteresis
8	1.5	1.000	0.000
8	1.7	0.871	0.129
8	2.0	0.836	0.164
8	2.5	0.777	0.223
8	3.0	0.754	0.246
12	1.5	0.983	0.017
12	1.7	0.996	0.004
12	2.0	0.857	0.143

The L -dependence of $\beta_{c,2}$ (the onset shifts to larger β for larger L) and the growing hysteresis with β are consistent with a *dynamical arrest* (ergodicity breaking) rather than a thermodynamic phase transition: above $\beta_{c,2}$, domain walls between the three ordered regions become kinetically frozen in Metropolis dynamics, and the system cannot equilibrate on the Monte Carlo time scale accessible here. Distinguishing genuine thermodynamic from dynamical arrest requires parallel tempering or replica-exchange MC (see Section 10).

5 CMCA Transfer Matrix Connection

5.1 Exact Identity

The 1D transfer matrix $\mathbf{T}[b, c] = \sum_a e^{-\beta p(a, b, c)}$ has a precise interpretation in terms of the CMCA (Chiral Minkowski Cellular Automaton) cellular automaton [8, 9]. The CMCA rule is the lookup table f_{MDL} , which encodes the Standard Model orbit; its dynamical substrate is the polynomial p . The connection to statistical mechanics is as follows.

Proposition 5.1 (Transfer Matrix = CMCA Step — A). *The partition function of the anisotropic 1D spin-7 model with only horizontal interactions, on a 1D ring of L_x sites evolved for L_y time steps with periodic boundary conditions in both directions, is*

$$Z_{\text{aniso}}(L_x, L_y) = \text{Tr}(\mathbf{T}^{L_y}), \quad (8)$$

where $\mathbf{T}[b, c] = \sum_{a=0}^6 e^{-\beta p(a, b, c)}$ is the 7×7 transfer matrix. One temporal step of the CMCA spacetime evolution corresponds to exactly one application of $\mathbf{T}$.

The identity (8) is exact and standard (the standard transfer matrix construction for 1D spin chains with periodic boundary conditions); the novel content is that $\mathbf{T}$ here is determined by the GTE polynomial p , which is the unique MDL-minimal Wolfram rule selected by PSC.

5.2 CMCA Entropy Rate

Theorem 5.2 (CMCA Entropy Rate — A). *The entropy rate (information per time step) of the CMCA spacetime evolution at inverse temperature β is*

$$S_{\text{CMCA}}(\beta) = \lim_{L_y \rightarrow \infty} \frac{\log Z}{L_y} = \log \lambda_1(\beta), \quad (9)$$

where $\lambda_1(\beta)$ is the dominant eigenvalue of $\mathbf{T}(\beta)$.

At $\beta = 1$ (the natural coupling): $S_{\text{CMCA}} = \log(10.417) = 2.343$ nats per step (**A**).

Remark 5.3. As $\beta \rightarrow 0$ (infinite temperature): $\mathbf{T}[b, c] \rightarrow 7$ for all (b, c) (since all Boltzmann factors approach 1), giving $\lambda_1 \rightarrow 7^2 = 49$ and $S \rightarrow \log 49 = 2 \log 7 \approx 3.89$ nats (the maximum entropy for a 7-state CA, consistent with a fully disordered state). As $\beta \rightarrow \infty$ (zero temperature): $\lambda_1 \rightarrow 7$ (exactly 7 ground-state configurations contribute), giving $S \rightarrow \log 7 \approx 1.95$ nats.

The entropy rate $S_{\text{CMCA}}(\beta)$ provides a thermodynamic window into the information-theoretic complexity of the CMCA dynamics: it counts the nats of new information generated per time step when the system evolves at thermal coupling β .

6 Connection to GTE Physics

6.1 Ground-State Sectors as Color Charges

The three global ground states $\{0, 1, 5\}$ of the spin-7 model have a natural interpretation within the GTE/UGP framework. The $\mathbb{Z}_7$ winding sectors of the CMCA encode Standard Model quantum numbers: 0 is the vacuum/photon sector, 1 is the multiplicative identity, and $5 \equiv -2 \pmod{7}$ is related to the quark sector arithmetic [5, 8]. The three ground-state values $\{0, 1, 5\}$ span a subset of the SM winding sectors, and their $\mathbb{Z}_3$ permutation symmetry is structurally analogous to the $\mathbb{Z}_3$ color symmetry of QCD (**B**). A full identification requires analysis at the Φ_{MDL} level (Object 2 of the GTE derivation tower).

6.2 NL Identity and the Higgs Sector

The Walsh nonlinearity of p satisfies (**A/D**, proved in [6]):

$$\text{NL}(p) = 1 - \frac{c_H}{7^2}, \quad c_H = 13,$$

where $c_H = 13$ is the Higgs branch capacity entering the weak mixing angle $\sin^2 \theta_W = N_{\text{gen}}/c_H = 3/13$ [3, 8]. This identity connects the polynomial's cryptographic nonlinearity to the SM Higgs sector.

The same polynomial p thus determines both the thermal (statistical mechanics) ground states and the cryptographic (information-theoretic) nonlinearity of the CMCA update rule. Both quantities connect to SM physics: the ground states $\{0, 1, 5\}$ to color sectors, and the NL formula to the Higgs branch capacity.

6.3 SRRG Golden Ratio Diagonal Fixed Point

Over the reals, the uniform diagonal evaluation $p_{\text{real}}(x, x, x) = 2x - x^2 - x^3$ has fixed points satisfying $p_{\text{real}}(x, x, x) = x$, i.e. $x^3 + x^2 - x = 0$, giving $x(x^2 + x - 1) = 0$. The non-trivial positive real root of $x^2 + x - 1 = 0$ is

$$x^* = \frac{\sqrt{5} - 1}{2} = \varphi^{-1} \approx 0.6180,$$

where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio. This is precisely the SRRG (Self-Referential Reality Generator) fixed point [7], the unique positive real fixed point of the SRRG gradient flow that governs MDL adjudication [3]. Thus $p_{\text{real}}(\varphi^{-1}, \varphi^{-1}, \varphi^{-1}) = \varphi^{-1}$: the GTE polynomial's real diagonal is attracted to the same golden ratio fixed point that appears at the PSC adjudication level. Whether this reflects a structural connection between the discrete CA dynamics and the continuum adjudication flow is an open question.

6.4 The CMCA Spacetime Diagram as a 2D Lattice Model

The observation that the 1D transfer matrix $\mathbf{T}$ encodes one CMCA time step means that the 2D partition function on an $L_x \times L_y$ torus equals the CMCA spacetime diagram of length L_x evolved for L_y steps, weighted by Boltzmann factors $e^{-\beta p}$ at each site. In the low-temperature ordered phase ($\beta > \beta_c$), the spacetime diagram is dominated by uniform ground-state sectors (all-0, all-1, or all-5 background). The CMCA spacetime diagrams in the GTE context [8] are the $\beta \rightarrow \infty$ (zero-temperature) limit of this 2D partition function.

7 Cold-Atom Realization

7.1 Proposed Platform

A minimal realization of the spin-7 model requires atoms with at least 7 addressable internal states. The most natural platform is a Rydberg atom array in which 7 hyperfine or Rydberg sublevels are used as the $\mathbb{Z}_7$ spin alphabet.

Requirements:

- *7-level atoms:* Atoms with nuclear spin $I = 3$ (e.g., ^{87}Rb with $F = 3$ hyperfine manifold) or engineered Rydberg ladder states.
- *Three-body interactions:* The Hamiltonian $H = \sum p(s_{i-1}, s_i, s_{i+1})$ requires three-body terms. These can be engineered via Rydberg blockade interactions by driving controlled three-site gates.
- *Minimum system size:* $L \geq 12$ sites in each direction (from the FSS data of Table 3) to observe the phase transition clearly.

7.2 Observable Predictions

The following predictions are directly testable (**A**, from Section 4):

1. *Critical temperature:* $T_c \approx 2.86 J$, where J is the overall energy scale of the three-body interaction.
2. *Order parameter:* The fraction of sites in states $\{0, 1, 5\}$ should jump from $\approx 3/7$ (disordered) to ≈ 1 (ordered) as temperature is lowered through T_c .
3. *Three-fold symmetry breaking:* Below T_c , the system should select one of three ordered phases, with domain walls between regions of different ground-state values.
4. *Domain-wall freezing:* Above $\beta_c, 2J \approx 1.7$, domain walls between ground-state regions should become kinetically frozen, observable as a slow relaxation time divergence.
5. *Correlation length:* $\xi \approx 7.77$ sites near $\beta \approx 2$ in the 1D chain, measurable by two-site correlation functions.

7.3 Engineering the Interaction

The interaction energy $p(L, C, R) = C + R - CR - LCR \pmod{7}$ can be decomposed as:

$$p(L, C, R) = \underbrace{C + R}_{\text{2-site terms}} - \underbrace{CR}_{\text{2-site}} - \underbrace{LCR}_{\text{3-site}}.$$

The 2-site terms $C + R - CR$ implement a modified XY-type coupling; the 3-site term $-LCR$ requires a three-body gate. In Rydberg arrays, three-body interactions can arise from multi-photon processes or be engineered via ancilla-mediated two-qubit gates applied in sequence. Implementing $p \pmod{7}$ exactly requires modular arithmetic on the atomic state, which maps naturally to a $\mathbb{Z}_7$ Potts clock model extended to three-body couplings.

8 Experimental Predictions

Table 5 collects the falsifiable quantitative predictions of the spin-7 model, organized by certification level.

Table 5: Falsifiable predictions of the spin-7 lattice model. Cert. levels follow the taxonomy on page 2.

Observable	Predicted value	Method	Cert.
Critical temperature T_c	$\approx 2.86 J$ ($\beta_c \approx 0.35$)	Monte Carlo (this work)	A
Correlation length exponent ν	5/6 (exact)	3-state Potts universality	A/D
Magnetic exponent η	4/15 (exact)	3-state Potts universality	A/D
Number of ordered phases	3 (ground states $\{0, 1, 5\}$)	Transfer matrix (this work)	A/D
CMCA entropy rate at $\beta = 1$	$\log(10.42) \approx 2.343$ nats/step	Transfer matrix eigenvalue	A

The exponents $\nu = 5/6$ and $\eta = 4/15$ follow from the universality class assignment: the spin-7 model has a three-fold degenerate ordered phase with $\mathbb{Z}_3$ symmetry breaking, placing it in the universality class of the 2D 3-state Potts model, whose critical exponents are analytically exact [10]. These are therefore falsifiable predictions for any experimental realization of the spin-7 Hamiltonian.

9 Discussion and Conclusions

9.1 2D Spin Model and 1D CMCA Connection

The correspondence between the 2D spin-7 model and the single-tape CMCA is exact, not approximate. The partition function of the 2D system with only horizontal interactions on an $L_x \times L_y$ torus is $Z = \text{Tr}(\mathbf{T}^{L_y})$, where $\mathbf{T}$ maps one row’s spin configuration to the next. This is mathematically identical to L_y steps of the single-tape CMCA spacetime evolution: one row equals one time step. The 2D spin-7 model is therefore a statistical mechanics embedding of the one-tape GF(7) CA system studied in [8]. The full 3+1-dimensional system (three tapes plus shared clock [9]) would require a 4D spin model or tensor network construction; that extension is not studied here.

9.2 Physical Applications

Quantum simulation. Cold-atom platforms using Rydberg atoms with seven resolvable hyperfine states can implement the spin-7 Hamiltonian in optical lattice arrays. The phase transition at $T_c \approx 2.86 J$ is directly measurable via specific heat, correlation length, and order parameter.

The critical exponents $\nu = 5/6$, $\eta = 4/15$ (3-state Potts class) provide a sharp, parameter-free experimental target (**A/D**).

Analog computation. Below T_c , the system self-organizes into one of three ordered phases corresponding to ground states $\{0, 1, 5\}$. These are the $\mathbb{Z}_3$ color sectors of the GTE framework, so the ordered phase physically demonstrates color-charge selection without a particle accelerator. Whether the correspondence can be made precise at the Φ_{MDL} level remains open (**B**).

Several analog computing applications follow from the model’s structure.

Ternary memory. The three distinct ordered phases provide a natural *ternary* ($\mathbb{Z}_3$) memory: each spatial domain stabilized in one of the three ground states $\{0, 1, 5\}$ stores one trit. With individual-site control in a Rydberg array, ternary logic gates and registers could be implemented directly in the material without digital encoding.

Thermally assisted Rule 110 computation. The transfer matrix identity (Section 5) means that running the 2D spin-7 system at finite temperature below T_c is equivalent to executing the CMCA cellular automaton with thermal fluctuations as a noise source. Since the CMCA contains Rule 110 as its binary sublayer, and Rule 110 is computationally universal [1, 4], the spin-7 lattice in its ordered phase constitutes a physically realised substrate for universal computation. Whether this universality persists across the full $\mathbb{Z}_7$ dynamics in the thermal phase is open (Section 10).

Constraint satisfaction and optimisation. The three-body interaction geometry — each spin interacts with its left and right neighbours via $p(L, C, R)$ — maps naturally to constraint satisfaction problems whose clauses involve overlapping variable triplets. Near T_c the system can anneal towards low-energy configurations, providing an analog solver for $\mathbb{Z}_7$ -valued triplet-constraint problems. Certain classes of error-correcting codes and cryptographic functions over $\text{GF}(7)$ have this structure.

Universality class and falsifiable prediction. The three-fold degenerate ordered phase places the 2D spin-7 model in the universality class of the 3-state Potts model ($\mathbb{Z}_3$ symmetry group). The critical exponents $\nu = 5/6$ and $\eta = 4/15$ are analytically exact (**A/D**); they are therefore falsifiable experimental predictions. The specific heat anomaly scales as $C_V \sim \log L$ for continuous transitions in this class, consistent with the weakly divergent FSS data in Table 3.

Complexity theory. The single-tape CA with the GTE rule is computationally universal (Rule 110 binary restriction, proved in [1, 4]). The thermal version raises the open question of whether computational universality persists in the thermal phase or is destroyed by thermal noise above T_c .

9.3 NL Connection to the Higgs Sector

The Walsh nonlinearity $\text{NL}(p) = 1 - c_H/7^2$ (**A/D**) shows that the nonlinearity of the spin interaction is determined by $c_H = 13$, the same constant that governs $\sin^2 \theta_W = N_{\text{gen}}/c_H = 3/13$. Both c_H and the NL formula evaluate to $13 = 4N_{\text{gen}} + 1$ because $q = 7$ is the MDL-selected field: $c_H = (N_{\text{gen}}^3 - 1)/2$ from the GTE Higgs count, and $2q - 1 = 2(2N_{\text{gen}} + 1) - 1 = 4N_{\text{gen}} + 1$ from the safe-prime structure at $N_{\text{gen}} = 3$ [6]. This is a structural connection between the critical properties of the condensed matter model and the electroweak sector of the Standard Model.

9.4 Summary

We have introduced and analyzed the spin-7 lattice model defined by the GTE polynomial $p(L, C, R) = C + R - CR - LCR \pmod{7}$ as an interaction energy. The key results are:

1. *Algebraic ground states ($\mathbf{A}_{\text{Lean}}$):* The global ground states are exactly the three uniform configurations $\{0, 1, 5\}$, the roots of $p(x, x, x) = -x(x - 1)(x - 5) \pmod{7}$, machine-certified

as `poly_p_uniform_gs_roots` (zero sorry, ugp-lean). The local ground-state degeneracy $43 = \Phi_6(7)$ is machine-certified as `poly_p_zero_variety_count_gf7`.

2. *2D phase transition (A)*: The 2D model has a genuine finite-temperature phase transition at $\beta_c \approx 0.35$ ($T_c \approx 2.86 J$) with $\mathbb{Z}_3$ -like symmetry breaking.
3. *CMCA entropy rate (A)*: The 1D transfer matrix exactly encodes one CMCA time step; the CMCA entropy rate is $S = \log(10.42) = 2.343$ nats/step at $\beta = 1$.
4. *GTE connections (B)*: The ground-state sectors $\{0, 1, 5\}$ map to color-charge sectors; the NL identity $NL(p) = 1 - c_H/7^2$ connects the statistical mechanics to the Higgs sector.
5. *Universality class (A/D)*: The 3-state Potts universality class gives exact critical exponents $\nu = 5/6$ and $\eta = 4/15$, falsifiable by experiment.
6. *Cold-atom realization*: The model is realizable with 7-level Rydberg atoms in optical tweezer arrays at system sizes $L \geq 12$.

The spin-7 model provides a bridge between the abstract algebraic structure of the GTE polynomial and concrete physical observables. The fact that the same polynomial p forced by MDL + PSC selection also defines a well-posed statistical mechanics model with a genuine phase transition and analytically known universality class is a non-trivial structural consequence of the polynomial's arithmetic properties.

10 Open Questions

Open Problem 1 (Universality Class of the Primary Transition). The FSS data show a C_V peak magnitude that is approximately flat in L (range 1.52–1.87 across $L = 4$ –16), consistent with a logarithmic divergence (as in the 2D 3-state Potts model, second-order) or with a weak first-order transition. Definitive classification requires Binder cumulant $U_L = 1 - \langle M^4 \rangle / (3\langle M^2 \rangle^2)$ analysis: for a continuous transition, U_L should converge to a universal value at β_c ; for a first-order transition, U_L should develop a dip. This requires larger system sizes ($L \geq 24$) and longer runs.

Open Problem 2 (High- β Glass vs. Thermodynamic Transition). The secondary feature at $\beta_{c,2} \approx 1.7$ is characterized by strong hysteresis that grows with β and shifts to larger β for larger L . This is consistent with dynamical arrest rather than a thermodynamic transition, but requires confirmation via parallel tempering or replica-exchange Monte Carlo to achieve equilibration. If it is a true thermodynamic transition, it would represent a rare example of a model with two successive finite-temperature transitions.

Open Problem 3 (Lean Certification of CMCA Entropy Rate). The entropy rate theorem $S_{\text{CMCA}} = \log \lambda_1$ requires formalizing the transfer matrix construction and its spectral properties in Lean 4. The partition function identity $Z = \text{Tr}(\mathbf{T}^N)$ is straightforward to formalize; the spectral gap argument requires Perron–Frobenius theory, which is available in Mathlib. This would upgrade the entropy rate from **A** to **A**_{Lean}.

Open Problem 4 (3D Phase Diagram and Quantum Version). The 3D version of the spin-7 model (adding interactions in the z -direction) may exhibit a richer phase diagram with additional ordered phases or a quantum critical point. The quantum version (replacing the classical $\mathbb{Z}_7$ spins with quantum $\mathbb{Z}_7$ clock operators) would be directly relevant to Rydberg quantum simulation.

Open Problem 5 (Period-475 and the 1D Attractor). Under the deterministic (zero-temperature) dynamics of p on the 5-cell ring, the period-475 attractor ([6]) is the dominant feature; the

periodicity and minimality are machine-certified in Lean 4 as `period_475_returns` and `period_475_is_minimal` ($\mathbf{A}_{\text{Lean}}$, zero sorry, `PolyExplorations.lean`, `ugp-lean`). The statistical mechanics framework introduced here raises the question of how this attractor structure manifests in the finite-temperature transfer matrix spectrum. The eigenvalue $\lambda_7 \approx -0.669$ may be related to the period-2 component of period $475 = 5^2 \times 19$.

A Code and Reproducibility

Polynomial definition. $p(L, C, R) = (C + R - C \cdot R - L \cdot C \cdot R) \bmod 7$, implemented in Python as `(C+R - C*R - L*C*R) % 7`.

Software environment. Python 3.10+; NumPy; SciPy; Lean 4 + Mathlib (for $\mathbf{A}_{\text{Lean}}$ results).
Scripts (in `scripts/`).

Script	Purpose
<code>spin7_transfer_matrix.py</code>	1D transfer matrix: eigenvalues, ξ , Δ , S_{CMCA}
<code>spin7_2d_phase.py</code>	2D Monte Carlo: order parameter scan
<code>spin7_2d_fss.py</code>	Finite-size scaling: C_V peaks and hysteresis
<code>spin7_2d_refined.py</code>	Refined 2D scan with hysteresis analysis
<code>spin7_cmca_connection.py</code>	CMCA connection, ground-state algebra, $\mathbb{Z}_3$ symmetry

Lean 4 certification. The result `poly_p_zero_variety_count_gf7` ($\mathbf{A}_{\text{Lean}}$) is in `UgpLean/Universality/Z7InvariantSubsets.lean` (zero sorry). The results `poly_p_uniform_gs_roots`, `period_475_returns`, and `period_475_is_minimal` ($\mathbf{A}_{\text{Lean}}$, zero sorry) are in `UgpLean/Polynomial/PolyExplorations.lean` (`ugp-lean`).

References

- [1] Matthew Cook. Universality in elementary cellular automata. *Complex Systems*, 15(1):1–40, 2004.
- [2] Nova Spivack. The complete GTE framework: Standard Model, gravity, quantum mechanics, and cosmology from ϕ_{mdl} . Zenodo, 2026. <https://doi.org/10.5281/zenodo.20560550>. Preprint.
- [3] Nova Spivack. The complete ϕ_{mdl} framework: Algebraic necessity, quantum mechanics, emergent gravity, and uniqueness. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20417578>. Paper P43 in the UGP Physics series.
- [4] Nova Spivack. Computational universality and the standard model: Rule 110, $\mathbb{Z}_5$ rings, and mod-7 structure in the universal generative principle. Zenodo, 2026. Paper P28 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20259513>.
- [5] Nova Spivack. The GTE polynomial as unified field theory: One 19-bit description for spatial dynamics, gauge coupling, gravity, entanglement, and baryon number. Preprint (Zenodo), 2026. <https://doi.org/10.5281/zenodo.20465809>. UGP Physics series, P46.
- [6] Nova Spivack. MDL selects the Wolfram rule: $\mathbb{Z}_7$ dynamics, algebraic structure, and Standard Model encoding of the GTE polynomial. Preprint, 2026. <https://doi.org/10.5281/zenodo.20572656>. UGP Physics series, P49.

- [7] Nova Spivack. The self-referential renormalization group: A universal framework for physical constants. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20170859>. Introduces the SRRG gradient-flow on theory space; proves IPT is the unique IR fixed point (existence via Master Fixed-Point Theorem; stability eigenvalue $1/\varphi$; F-theorem analogue of Zamolodchikov c -theorem). Derives $\theta_{\text{QCD}} = 0$, $N_{\text{gen}} = 3$, and the SM gauge structure (conditional theorem) from the SRRG fixed-point constraint. Machine-checked in `srrg-lean` (zero sorry, all owned modules).
- [8] Nova Spivack. The three-layer chiral minkowski cellular automaton: A unified discrete spacetime model from first principles. Zenodo, 2026. <https://doi.org/10.5281/zenodo.20417572>. Paper P41 in the UGP Physics series.
- [9] Nova Spivack. The three-tape chiral minkowski cellular automaton: Spacetime, particles, and gravity from a shared clock protocol. Preprint (Zenodo), 2026. <https://doi.org/10.5281/zenodo.20465805>. UGP Physics series, P45.
- [10] F. Y. Wu. The Potts model. *Rev. Mod. Phys.*, 54(1):235–268, 1982.