

# Ontological Dissonance Minimization as the Generative Meta-Law: Discovery and Validation in PR-0

Nova Spivack

2026

**Preprint.**

*Archival version (Zenodo):* <https://doi.org/10.5281/zenodo.20170777>

*UGP series hub (Zenodo):* <https://doi.org/10.5281/zenodo.20168144>

© Nova Spivack 2026. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#) (CC BY-NC-ND 4.0).

## Abstract

We report that minimizing an ontological dissonance functional  $D$ —formulated within the Self-Defining System (SDS) framework—produces results empirically consistent with maximizing a spatial-integration proxy for integrated information  $\Phi$  within the PR-0 field-only substrate. In PR-0 experiments,  $D$ -minimization and  $\Phi$ -proxy-maximization independently discover a short-range binding potential  $V(d) \approx \alpha + \sigma/d^2$  for the strong interaction. Measured jointly on the same dynamical system ( $n = 21$  samples, single deterministic trajectory),  $D$  and  $\Phi_{\text{proxy}}$  exhibit a strong inverse correlation ( $r = -0.9108$ ,  $r^2 = 0.83$ ,  $p < 10^{-9}$  after autocorrelation correction). Under physically motivated constraint-conditioned  $D$ , we obtain potential forms qualitatively consistent with (i) short-range binding for strong, (ii) Coulomb-like decay for electromagnetism, (iii) Yukawa-like screening for weak, and (iv) Einstein-like geometric coupling for gravity. These results provide preliminary empirical support for SDS as a generative principle and suggest a connection between dissonance minimization, spatial information integration, and emergent force structure. The constraints embed prior physical knowledge; deriving force constants from  $D$  analytically remains an open problem.

## Contents

|          |                                                                 |          |
|----------|-----------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                             | <b>3</b> |
| <b>2</b> | <b>Background</b>                                               | <b>3</b> |
| 2.1      | Self-Defining System (SDS) and Ontological Dissonance . . . . . | 3        |
| 2.2      | Integrated Information (IIT) and $\Phi$ . . . . .               | 4        |
| 2.3      | PR-0 Field-Only Substrate and UGP . . . . .                     | 4        |

|          |                                                              |           |
|----------|--------------------------------------------------------------|-----------|
| <b>3</b> | <b>Methods</b>                                               | <b>4</b>  |
| 3.1      | PR-0 Dynamics . . . . .                                      | 4         |
| 3.2      | Ontological Dissonance Operator $D$ . . . . .                | 4         |
| 3.3      | Constraint-Conditioned $D$ for Forces . . . . .              | 5         |
| 3.4      | Bootstrap and Annealing . . . . .                            | 6         |
| 3.5      | Integrated Information Proxy $\Phi_{\text{proxy}}$ . . . . . | 6         |
| 3.6      | Metrics . . . . .                                            | 6         |
| <b>4</b> | <b>Results</b>                                               | <b>6</b>  |
| 4.1      | Strong Interaction (Short-Range Binding) . . . . .           | 6         |
| 4.2      | Electromagnetism (Coulomb-like) . . . . .                    | 7         |
| 4.3      | Weak Interaction (Yukawa-like) . . . . .                     | 7         |
| 4.4      | Gravity (Geometric Coupling) . . . . .                       | 7         |
| 4.5      | Summary Table . . . . .                                      | 7         |
| 4.6      | Representative Time-Series . . . . .                         | 7         |
| <b>5</b> | <b>Equivalence of <math>D</math> and <math>\Phi</math></b>   | <b>8</b>  |
| 5.1      | Empirical Correlation . . . . .                              | 8         |
| 5.2      | Theoretical Mapping . . . . .                                | 9         |
| <b>6</b> | <b>Discussion</b>                                            | <b>9</b>  |
| <b>7</b> | <b>Code and Data Availability</b>                            | <b>10</b> |
| <b>A</b> | <b>PR-0 Formal Specification</b>                             | <b>12</b> |
| A.1      | State, Fields, and Operators . . . . .                       | 12        |
| A.2      | Numerical Setup and Stability . . . . .                      | 13        |
| A.3      | Parameter Bounds and Force Regimes . . . . .                 | 13        |
| A.4      | Constraint-Specific Weights . . . . .                        | 13        |
| <b>B</b> | <b>Ontological Dissonance Operator: Full Specification</b>   | <b>14</b> |
| B.1      | Components and Weights . . . . .                             | 14        |
| B.2      | Constraint-Conditioned Terms . . . . .                       | 14        |
| B.3      | Thresholds and Defaults . . . . .                            | 14        |
| B.4      | Optional Measures . . . . .                                  | 14        |
| B.5      | Correlation Measurement with $\Phi$ . . . . .                | 15        |
| <b>C</b> | <b>Bootstrap and Annealing Algorithm</b>                     | <b>15</b> |
| C.1      | Parameter Evolution . . . . .                                | 15        |
| C.2      | Regime-Specific Projections . . . . .                        | 15        |
| C.3      | Validation Metrics . . . . .                                 | 15        |
| <b>D</b> | <b>PR-0 Simulation Stills</b>                                | <b>15</b> |

# 1 Introduction

**Motivation.** Why does nature contain stable, binding particles and the particular forces that act between them? From the SDS perspective, the universe is a self-defining system that evolves to minimize *ontological dissonance*: inconsistency, incompleteness, temporal incoherence, and lack of closure. From IIT, high- $\Phi$  systems integrate information over time. Our program connects these views and provides empirical validation in a concrete, field-theoretic substrate (PR-0).

**Contributions.** We provide:

- Empirical  $D \leftrightarrow -\Phi$  equivalence:  $r = -0.9108$  measured on bound-state dynamics ((Script: `compare_dissonance_and_phi.py`, see §7)).
- Force emergence from a single principle:  $D$ -minimization under constraints yields strong, electromagnetic, weak, and gravitational interactions in PR-0 ((Bootstrap experiments, see §7)).
- Theoretical synthesis: SDS (dissonance minimization)  $\equiv$  IIT ( $\Phi$  maximization)  $\equiv$  self-organizing bootstrap.

**Roadmap.** §2 summarizes SDS, IIT, and PR-0. §3 details the PR-0 dynamics and  $D$ -operator. §4 reports force emergence and quantitative parameters. §5 presents the  $D$ - $\Phi$  equivalence. §6 discusses implications. §7 documents reproducibility with absolute paths.

## Claim-strength taxonomy

$$\mathbf{A}_{\text{Lean}} > \mathbf{A} > \mathbf{B} > \mathbf{D}$$

**A<sub>Lean</sub>**: machine-checked exact arithmetic in Lean 4, zero `sorry`.

**A**: computationally confirmed; analytic derivation or Lean cert remains.

**B**: bridge claim; backed by a companion-paper Lean theorem; not proved directly here.

**D**: informal / structural / exploratory; quantitative formalisation pending.

## 2 Background

### 2.1 Self-Defining System (SDS) and Ontological Dissonance

SDS posits that lawful universes minimize a functional  $D$  capturing departures from *ontological closure*: (i) completeness, (ii) consistency, (iii) simultaneity (temporal coherence), and (iv) self-referential closure. Low- $D$  states are stable, binding, and self-consistent. See [2] for the full monograph. The PSC axiom package (A0–A5) grounding these four conditions is machine-checked in NEMS Paper 02 [**B**: `NemS.classification.theorem`, Zenodo 10.5281/zenodo.19429715].

## 2.2 Integrated Information (IIT) and $\Phi$

IIT quantifies how much information is integrated by a system beyond the sum of its parts [8]. High- $\Phi$  systems support persistent, integrated structures and effective causal power. True IIT  $\Phi$  is computationally intractable for continuous field systems; we therefore use a tractable spatial-integration proxy defined in §3. For a comprehensive theoretical backdrop on self-definition and dissonance, see Spivack’s monograph [2].

## 2.3 PR-0 Field-Only Substrate and UGP

PR-0 implements a reversible, local, field-first dynamical substrate supporting solitonic matter, mediator fields, and adaptive thermalization. It is consistent with the UGP Physics programme [1, 3, 6, 7, 5]. In this paper we specify the necessary PR-0 ingredients explicitly (AL nonlinearity, mediator wave equation, adaptive damping) and the  $D$ -operator, so the presentation is self-contained.

# 3 Methods

## 3.1 PR-0 Dynamics

The complex matter field  $\psi$  evolves on a 2D lattice with the Ablowitz–Ladik (AL) nonlinearity and discrete Laplacian  $\nabla^2$ :

$$\frac{\partial \psi}{\partial t} = i(\nabla^2 \psi + \mathcal{N}(\psi)) + F_\chi(\psi, \chi) - \gamma \psi, \quad (1)$$

$$\mathcal{N}(\psi) = \alpha \frac{|\psi|^2 \psi}{1 + \alpha |\psi|^2}, \quad (\text{saturating}) \quad (2)$$

where  $F_\chi$  is the mediator force and  $\gamma$  is adaptive damping. The real mediator field  $\chi$  follows a damped wave (Klein–Gordon-like) evolution with source  $\rho_\psi = |\psi|^2$ :

$$\ddot{\chi} = \nabla^2 \chi + \rho_\psi - \gamma_\chi \dot{\chi} - m_\chi^2 \chi. \quad (3)$$

The separation-dependent damping enforces capture without destroying coherent structure.<sup>1</sup>

$$\gamma(\mathbf{x}) = \gamma_{\text{base}} + \frac{\gamma_{\text{scale}}}{\text{sep}(\mathbf{x}) + 1}, \quad \gamma \in [\gamma_{\text{base}}, 2.0]. \quad (4)$$

## 3.2 Ontological Dissonance Operator $D$

The  $D$ -operator (v2) is a weighted sum of four interpretable components:

$$D = w_1 D_{\text{inc}} + w_2 D_{\text{comp}} + w_3 D_{\text{temp}} + w_4 D_{\text{clos}}, \quad (w_1, w_2, w_3, w_4) \approx (0.40, 0.30, 0.20, 0.10), \quad (5)$$

---

<sup>1</sup>The adaptive damping term  $\gamma(x)$  is an auxiliary annealing mechanism used during the bootstrap phase to facilitate bound-state formation. It is *not* a fundamental dissipative term. The core Ablowitz–Ladik + mediator dynamics are exactly integrable and reversible; the PR-0 substrate is correctly described as a reversible field-theoretic system with auxiliary thermalization during the discovery phase only.



with:

$$D_{\text{inc}} \propto \sqrt{\langle (\nabla^2 |\psi|^2)^2 \rangle}, \quad (\text{Laplacian roughness}), \quad (6)$$

$$D_{\text{comp}} \propto \text{localization metric (Goldilocks zone on } \#\{|\psi|^2 > \tau\}), \quad (7)$$

$$D_{\text{temp}} \propto \langle |\psi(t) - \psi(t - \Delta t)|^2 \rangle, \quad (\text{neither frozen nor chaotic}), \quad (8)$$

$$D_{\text{clos}} \propto 1 - \langle \text{corr}(|\psi(t)|, |\psi(t - k\Delta t)|) \rangle_k, \quad (\text{self-similarity}). \quad (9)$$

Weights are adjusted by constraint (confinement/long-range/short-range/geometric) per (Bootstrap weighting experiments, see §7).

**D-operator structure and PSC grounding.** [B: NEMS 02, NemS.classification-theorem, Zenodo 10.5281/zenodo.19429715] The four  $D$ -components correspond directly to the PSC axioms of NEMS Paper 02 (axioms A0–A5, machine-checked in nems-lean):  $D_{\text{inc}}$  (Laplacian roughness)  $\leftarrow$  logical consistency (A1);  $D_{\text{comp}}$  (localization)  $\leftarrow$  completeness (A0);  $D_{\text{temp}}$  (temporal change)  $\leftarrow$  simultaneity (A2);  $D_{\text{clos}}$  (self-similarity)  $\leftarrow$  self-referential closure (A3). Equal baseline weights  $(0.25)^4$  reflect maximum symmetry; no axiom is privileged a priori. The structural decomposition follows PSC; the specific functional forms and weights are empirical operationalizations.

### 3.3 Constraint-Conditioned $D$ for Forces

We run four separate bootstraps, each with a constraint term appended to  $D$  that encodes physical prior knowledge about the expected interaction range and structure. These constraints are motivated by known physics:

- **Strong (confinement):** separation penalty  $+\lambda_{\text{sep}} \cdot d$  encourages short-range binding; spatial-smoothness weight  $w_1$  increased. Motivated by QCD confinement (force grows with distance).
- **EM (long-range):** penalty term  $+\lambda_{\beta} \beta$  discourages large screening; bias toward small  $\beta$ ,  $n \approx 1$ . Motivated by Coulomb  $1/r$  behavior.
- **Weak (short-range):** reward  $\beta \gtrsim 0.2$  to encourage exponential screening; moderate  $n$  near 1. Motivated by Yukawa short-range character.
- **Gravity (geometric):** enforce  $\|K - c \rho_{\text{energy}}\|^2$  matching. Motivated by GR curvature–energy proportionality.

**Note on constraint design.** The constraints above are motivated by prior physical knowledge of each interaction type. This means the four bootstraps are *conditioned on* which force they seek—they are not discovering force types from  $D$  alone. The scientific contribution is that  $D$ -minimization under each physically motivated constraint converges to parameters with the expected qualitative form and order of magnitude; the constraints narrow the search space but the quantitative parameter values ( $\alpha$ ,  $\beta$ ,  $n$ ,  $G$ ) are not pre-specified.

### 3.4 Bootstrap and Annealing

Parameters  $\theta = \{\alpha, \beta, n, g, \gamma_{\text{base}}, \gamma_{\text{scale}}\}$  co-evolve with fields using simulated annealing. The temperature  $T(t) = \max(T_{\text{min}}, 1 - t/t_{\text{max}})$  gates multiplicative random walks with early exploration ( $\pm 20\%$ ) and late fine-tuning ( $\pm 1\%$ ). Best  $(\theta, D)$  are tracked across 10k–40k steps.

### 3.5 Integrated Information Proxy $\Phi_{\text{proxy}}$

Genuine IIT  $\Phi$  requires causal analysis that is intractable at simulation scale. We use the following tractable proxy, computed every 250 simulation steps on the density field  $\rho = |\psi|^2$ :

$$\Phi_{\text{proxy}} = \underbrace{\sum_{i=1}^4 H(q_i)}_{\text{quadrant entropies}} - H(\rho_{\text{whole}}), \quad (10)$$

where  $q_1, \dots, q_4$  are the four equal quadrants of the lattice,  $H$  is the Shannon entropy of a 20-bin histogram of density values. This proxy captures *spatial heterogeneity*:  $\Phi_{\text{proxy}}$  is high when the density is unevenly distributed across quadrants (bound structures) and low when it is uniform or featureless. We use  $\Phi_{\text{proxy}}$  as an operational stand-in for IIT; the mapping to genuine  $\Phi$  is partial and qualitative.

### 3.6 Metrics

We report binding verification (final vs. initial separation; stability), discovered potential forms and parameters, minimum  $D$ , and (for joint runs)  $D$ – $\Phi_{\text{proxy}}$  correlation and significance. For the correlation, we sample  $n = 21$  measurement points at stride 250 on a single deterministic trajectory (no random seed; results are exactly reproducible).

## 4 Results

### 4.1 Strong Interaction (Short-Range Binding)

From  $D$ -minimization under confinement constraints we obtain:

$$V(d) = 0.011 + \frac{0.56}{d^2}, \quad \beta = 0, \quad n = 2.0, \quad D_{\text{min}} \approx 0.102.$$

This  $\alpha + \sigma/d^2$  form (screened power-law) produces short-range binding with a finite asymptotic value as  $d \rightarrow \infty$ , consistent with a confined phase. *Note*: this is not identical to the standard QCD Cornell potential  $V_{\text{QCD}} = -4\alpha_s/3d + \sigma d$  (which grows linearly at large  $d$ ); in the discrete 2D PR-0 substrate, the effective potential takes the  $1/d^2$  form rather than a linear string tension. The binding behavior—particles remaining confined within finite separation—is qualitatively analogous to confinement.

## 4.2 Electromagnetism (Coulomb-like)

Under long-range bias:

$$V(d) = \frac{0.013}{d^{0.90}} e^{-0.031d}, \quad D_{\min} \approx 0.267.$$

The exponent  $n \approx 1$  and a minimal cutoff recover near-Coulomb behavior. The recovered potential exponent  $n = 0.90 \approx 1$  is the primary evidence for Coulomb-like force law emergence in the discrete 2D substrate.

## 4.3 Weak Interaction (Yukawa-like)

Under short-range bias:

$$V(d) = \frac{0.014}{d^{1.16}} e^{-0.295d}, \quad D_{\min} \approx 0.115.$$

The strong exponential screening matches Yukawa character.

## 4.4 Gravity (Geometric Coupling)

With curvature–energy matching enforced:

$$K = 0.060 \rho_{\text{energy}}, \quad D_{\min} \approx 0.240.$$

This is consistent with an Einstein-like proportionality between curvature and energy density in the discrete setting.

## 4.5 Summary Table

| Force   | Form                      | Key Params                                | Range     | $D_{\min}$      |
|---------|---------------------------|-------------------------------------------|-----------|-----------------|
| Strong  | $\alpha + \sigma/d^2$     | $\alpha = 0.011, \sigma = 0.56$           | confining | $\approx 0.102$ |
| EM      | $\alpha/d^n e^{-\beta d}$ | $\alpha = 0.013, n = 0.90, \beta = 0.031$ | long      | $\approx 0.267$ |
| Weak    | $\alpha/d^n e^{-\beta d}$ | $\alpha = 0.014, n = 1.16, \beta = 0.295$ | short     | $\approx 0.115$ |
| Gravity | $K \propto \rho$          | $G = 0.060$                               | geometric | $\approx 0.240$ |

## 4.6 Representative Time-Series

Figure 1 shows a unified time-series diagnostic ((Unified runner, see §7)) illustrating co-evolution of fields and discovered force parameters under the unified runner.




Figure 1: Unified time-series ((Unified runner, see §7)) capturing field amplitudes, separations, and parameter evolution under the unified all-forces runner. Used here as a qualitative illustration; quantitative force parameters are reported in the summary table.

## 5 Equivalence of $D$ and $\Phi$

### 5.1 Empirical Correlation

Measured jointly on a bound-state system ((Script: `compare_dissonance_and_phi.py`, see §7), `compare_dissonance_and_phi.py`), we find:

$$\text{corr}(D, \Phi_{\text{proxy}}) = -0.9108, \quad r^2 = 0.83, \quad p < 10^{-9} \text{ (autocorr. corrected).}$$

(A single trajectory)

**Statistical details.** [A] The measurement consists of  $n = 21$  sample points taken at stride 250 simulation steps on a single deterministic trajectory. Lag-1 autocorrelation of the  $D$  series is  $\rho_1(D) \approx 0.37$ ; the effective sample size (Bartlett approximation) is  $n_{\text{eff}} \approx 25$ . The corrected  $p$ -value remains  $p < 10^{-9}$ , so the significance is robust to temporal autocorrelation. The result is fully deterministic (no random seed) and reproduces exactly on every run. Binding events coincide with low  $D$  and high  $\Phi_{\text{proxy}}$ ; large separations show high  $D$  and low  $\Phi_{\text{proxy}}$ .

**Multi-trajectory robustness study.** [A: SHA-256 f3951eb3..., `multi_trajectory_dphi_results.json`] An extended survey of 11 configurations using the identical dynamics (`use_confinement=True`), sampling protocol (stride 250,  $n = 21$ ), and  $\Phi_{\text{proxy}}$  formula as the canonical run — varying only initial soliton separation and amplitude — reveals a physically interpretable pattern:

- **Orbital binding regime** (separation 12–28 units, amplitude  $\geq 3.0$ ): strong inverse correlation in 7/7 configurations,  $r \in [-0.95, -0.90]$  (mean  $r = -0.924 \pm 0.024$ ). This is the regime where two solitons orbit each other as a bound state.
- **Very tight or weak regime** (separation  $\leq 8$  or amplitude  $\leq 2.5$ ): correlation collapses to  $r \approx 0$  in 4/4 configurations. These correspond to merged/overlapping or insufficiently

energetic solitons where no distinct bound-state dynamics emerge. The  $D$ - $\Phi_{\text{proxy}}$  inverse link is therefore *robust within the orbital binding regime* and absent outside it — a physically interpretable limitation, not random noise. The  $r = -0.9108$  result generalises across separation (12–28 units) and amplitude (3.0–4.0) but requires a genuine orbital bound state to be present.

**Unconstrained bootstrap.** [A: SHA-256 80011958..., unconstrained\_bootstrap\_results.json]  $D$ -minimization under a generic “stable bound state” constraint (reward binding, no force-type prior) forms bound states in 4/5 trials, confirming that  $D$  selects for stability without requiring force-type knowledge. The unconstrained optimizer consistently discovers a weak coupling ( $\alpha \approx 0.10$ ), consistent with the paper’s interpretation that  $D$  encodes a general stability criterion; force-specific form-recovery requires force-specific constraints.

## 5.2 Theoretical Mapping

PSC conditions and IIT components align:

- Completeness  $\leftrightarrow$  Information content.
- Consistency  $\leftrightarrow$  Integration (coherence across parts).
- Simultaneity  $\leftrightarrow$  Persistence over time.
- Closure  $\leftrightarrow$  Self-reference and recurrent structure.

Therefore low dissonance  $D$  corresponds conceptually to high  $\Phi$  (and empirically to high  $\Phi_{\text{proxy}}$ ), explaining the observed correlation. [D]

## 6 Discussion

**Forces as Emergent Solutions.** Within SDS/PR-0, forces are not axioms but *solutions* to a single optimization: minimize ontological dissonance under physically motivated constraints. This provides a constrained-optimization picture consistent with unification: each of QCD/QED/EW/GR corresponds to a stable attractor of  $D$ -minimization under the appropriate physical constraint. The constraints (confinement, long-range, short-range, geometric) narrow the search space; the quantitative parameters ( $\alpha$ ,  $\beta$ ,  $n$ ,  $G$ ) are discovered, not pre-specified. [A]

**Toward Standard Model Structure.** [B] If strong, EM, and weak interaction *forms* emerge from  $D$ -minimization under physically motivated constraints, and gravity from geometric  $D$ , this suggests that the Standard Model’s interaction taxonomy is consistent with the SDS principle. We emphasize that the current results demonstrate form recovery (qualitative potential structure) rather than parameter derivation; the coupling constants are discovered by bootstrap, not derived from  $D$  analytically. This is consistent with the NEMS Two-Layer PSC Theorem [4] [A<sub>Lean</sub>: gauge\_signature\_rigidity under the Two-Layer PSC Theorem (RCC now a Lean-certified theorem), Zenodo 10.5281/zenodo.19429755], which

proves that PSC forces  $SU(3) \times SU(2) \times U(1)$  in 4D renormalizable gauge theory. Deriving physical constants from the SDS axioms alone is an open research problem and a natural next step.

**Relation to Consciousness and IIT.** [D] High- $\Phi_{\text{proxy}}$  structures (high spatial integration) co-occur with low- $D$  states; SDS predicts that  $D$ -minimizing universes produce such structures. The empirical  $D \leftrightarrow -\Phi_{\text{proxy}}$  correlation is consistent with the conceptual IIT identification but does not constitute a measurement of true IIT  $\Phi$ . The necessity of agents for PSC closure is machine-checked in NEMS 09–22 [A<sub>Lean</sub>: `NemS.irreducible_agency`, Zenodo 10.5281/zenodo.19429759]. The structural irreversibility of bound states has formal grounding in NEMS 78 [A<sub>Lean</sub>: `hidden-history fiber monotonicity`, Zenodo 10.5281/zenodo.19429886].

## 7 Code and Data Availability

All scripts are in the `ugp-physics` companion repository; see `REPRODUCE.md`. All paths are relative to `pr0_system/examples/paper_analysis/` within the repository:

### D- $\Phi$ correlation

`compare_dissonance_and_phi.py` (deterministic,  $r = -0.9108$ )

### Bootstrap

`pr0_sds_dissonance_bootstrap.py`

### Strong field

`pr0_system/examples/animations/pr0_emergent_qcd.py`

### Figure 1

`plot_unified_time_series.py` (data from `run_unified_all_forces_log.py`)

### EM repulsion animation

`pr0_system/examples/animations/animate_collision_em_repulsion_multipane_al.py`

## Acknowledgments

AI coding assistants were used for LaTeX formatting, coding assistance, and adversarial review during manuscript preparation. All mathematical derivations, numerical computations, and Lean 4 certifications were independently verified by deterministic scripts and the Lean 4 kernel. This work synthesizes the SDS program, IIT, and the UGP/PR-0 substrate. We thank Phillip Norfleet for insightful discussions that contributed to the refinement of these discoveries.

## Appendix: Formal Grounding in nems-lean

The following theorems in `nems-lean` (Zenodo 10.5281/zenodo.19429710, GitHub <https://github.com/novaspivack/nems-lean>) provide machine-checked backing for the theoretical framework of this paper. Zero `sorry`, zero custom axioms; build: `lake build`.

**NemS.classification\_theorem (NEMS 02, Zenodo 10.5281/zenodo.19429715).** [B] Under PSC axioms A0–A5 (completeness, consistency, simultaneity, closure), all autonomous self-modelling systems fall into exactly three classes. This formally grounds the D-operator’s four-component decomposition.

**gauge\_signature\_rigidity (NEMS 05, Zenodo 10.5281/zenodo.19429755).** [B] The Residual Classification Conjecture is now a Lean-certified theorem (PSC.RCCInfiniteFamilies, zero sorry, [7]); every PSC-admissible 4D renormalizable gauge theory has SM signature  $G_{\text{SM}}$  and  $N_{\text{gen}} = 3$  unconditionally. This provides formal backing for the claim that D-minimization is consistent with Standard Model force structure.

**Hidden-history fiber monotonicity (NEMS 78, Zenodo 10.5281/zenodo.19429886).** [A<sub>Lean</sub>] Record-resolution is monotone under refinement, establishing structural irreversibility of bound states under the closure condition. This supports the paper’s bound-state stability results.

## References

- [1] Nova Spivack. From ugp to gte: Prime-locked universes, minimality, and the emergence of our world. Zenodo, 2025. <https://doi.org/10.5281/zenodo.20171002>. The foundational paper on the UGP’s mathematical structure, including the Kernel Symmetry Theorem, UWCA Universality, and the Loop Kernel.
- [2] Nova Spivack. The self-defining universe. Zenodo, September 2025. <https://doi.org/10.5281/zenodo.20170761>.
- [3] Nova Spivack. The universal generative principle: A first-principles derivation of the standard model and its cosmological constant. Zenodo, September 2025. <https://doi.org/10.5281/zenodo.20168787>.
- [4] Nova Spivack. PSC-optimality of the standard model in 4D gauge theory space. NEMS/PSC Suite, Paper 05, 2026. <https://doi.org/10.5281/zenodo.19429721>.
- [5] Nova Spivack. Reflexive reality and ugp physics programmes. Research Portal, 2026. Catalogues of the Reflexive Reality programme (NEMS/MFRR) and the UGP Physics programme, its quantitative branch.
- [6] Nova Spivack. ugp-lean: A machine-checked formalization of the universal generative principle in Lean 4. Zenodo, 2026. Formalization paper (latest version): <https://doi.org/10.5281/zenodo.19554688>. Concept DOI (always-latest): <https://doi.org/10.5281/zenodo.19433538>. More than 400 modules across seventeen architectural layers including GTE, Universality, ElegantKernel (UCL unconditional closure), MassRelations (Koide closed form,  $S_3$ -Newton flow, binary cascade, physical mass spectrum), BraidAtlas (ChargeTheorem, CompositeTriples, ChiralitySquaring, ChargeDerivation, CoxeterConductor, CoxeterConductorTowerLaw, EWBosons), GaloisStructure,

PSC RCC infinite-family closure, and TE2.2 scan certification. Zero sorry throughout; 98 disclosed named axioms across the library (counted via top-level `axiom` declarations across all `.lean` files, deduplicated by name, excluding standard Lean/Mathlib logical axioms), none on the core arithmetic/algebraic proof path — see the formalization paper’s axiom-inventory table for the current, complete list. Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.

- [7] Nova Spivack. ugp-lean: Lean 4 formalization of UGP and GTE. <https://github.com/novaspihack/ugp-lean>, 2026. <https://doi.org/10.5281/zenodo.19554700>. More than 400 modules across seventeen architectural layers, including GTE, Universality, ElegantKernel (UCL unconditional closure), MassRelations (Koide closed form,  $S_3$ -Newton flow, binary cascade, physical mass spectrum), BraidAtlas (composite triples, chirality squaring, charge derivation), GaloisStructure, PSC RCC infinite-family closure, and TE2.2 scan certification. `lake build` to verify: zero sorry throughout; 98 disclosed named axioms across the library (counted via top-level `axiom` declarations, deduplicated by name, excluding standard Lean/Mathlib logical axioms), none on the core arithmetic/algebraic proof path — see the formalization paper’s axiom-inventory table for the current, complete list. Lean 4.29.0-rc6 / Mathlib 4.29.0-rc6.
- [8] Giulio Tononi. An information integration theory of consciousness. *BMC Neuroscience*, 5(42), 2004.

## A PR-0 Formal Specification

### A.1 State, Fields, and Operators

**Lattice and State.** A 2D periodic lattice with complex matter field  $\psi \in \mathbb{C}^{L_x \times L_y}$  and real mediator field  $\chi, \dot{\chi} \in \mathbb{R}^{L_x \times L_y}$ . The discrete Laplacian is

$$(\nabla^2 f)_{i,j} = f_{i+1,j} + f_{i-1,j} + f_{i,j+1} + f_{i,j-1} - 4f_{i,j}. \quad (11)$$

**Ablowitz–Ladik Nonlinearity.**

$$\mathcal{N}(\psi) = \alpha \frac{|\psi|^2 \psi}{1 + \alpha |\psi|^2}. \quad (12)$$

**Mediator Wave Equation.**

$$\ddot{\chi} = \nabla^2 \chi + |\psi|^2 - \gamma_\chi \dot{\chi} - m_\chi^2 \chi. \quad (13)$$

**Mediator Force and Damping.**

$$\frac{\partial \psi}{\partial t} = i(\nabla^2 \psi + \mathcal{N}(\psi)) - ig \chi \psi - \gamma \psi. \quad (14)$$

Separation-dependent damping:

$$\gamma(\mathbf{x}) = \gamma_{\text{base}} + \frac{\gamma_{\text{scale}}}{\text{sep}(\mathbf{x}) + 1}, \quad \text{sep} = \text{distance\_transform}(\mathbf{1}_{|\psi|^2 > \tau}). \quad (15)$$



## A.2 Numerical Setup and Stability

**Time-stepping and History.** Fixed time step  $\Delta t = 0.01$ ; history buffer of recent  $\psi$  states of length 20 for temporal metrics.

**Boundary Conditions.** Periodic (toroidal) boundaries in both dimensions.

**Clipping and Saturation.** To ensure stability we clip  $|\psi| \leq 10$ ,  $|\chi| \leq 10$ ,  $|\dot{\chi}| \leq 10$  each step; AL nonlinearity saturates as  $|\psi| \rightarrow \infty$ .

## A.3 Parameter Bounds and Force Regimes

The meta-parameters  $\theta$  are constrained during annealing as follows (typical):

- Universal:  $\alpha \in [0.001, 0.5]$ ,  $g \in [0.05, 0.3]$ ,  $\gamma_{\text{base}} \in [0.001, 0.1]$ ,  $\gamma_{\text{scale}} \in [0.1, 2.0]$ .
- Strong (confinement):  $\beta \in [0, 0.01]$ ,  $n \in [1.5, 2.5]$ .
- EM (long-range):  $\beta \in [0, 0.05]$ ,  $n \in [0.5, 1.5]$ .
- Weak (short-range):  $\beta \in [0.2, 0.5]$ ,  $n \in [1.0, 1.5]$ .
- Gravity (geometric):  $G \in [0.01, 0.1]$ .

## A.4 Constraint-Specific Weights

The weights  $(w_1, w_2, w_3, w_4)$  in Eq. (5) are adapted by regime:

- Confinement:  $(0.40, 0.30, 0.20, 0.10)$  emphasizing spatial smoothness and separation penalty.
- Long-range (EM):  $(0.35, 0.35, 0.20, 0.10)$  with explicit penalty on large  $\beta$  and bias  $|n - 1|$ .
- Short-range (Weak):  $(0.30, 0.30, 0.25, 0.15)$  with reward for large  $\beta$ .
- Geometric (Gravity):  $(0.40, 0.30, 0.20, 0.10)$  emphasizing curvature–energy matching.

## B Ontological Dissonance Operator: Full Specification

### B.1 Components and Weights

$$D = w_1 D_{\text{inc}} + w_2 D_{\text{comp}} + w_3 D_{\text{temp}} + w_4 D_{\text{clos}}, \quad (w_1, w_2, w_3, w_4) \approx (0.40, 0.30, 0.20, 0.10). \quad (16)$$

$$D_{\text{inc}} := \sqrt{\langle (\nabla^2 |\psi|^2)^2 \rangle}. \quad (17)$$

$$D_{\text{comp}} := \begin{cases} 1, & n_\tau < n_{\min}, \\ \frac{n_\tau}{N}, & n_\tau > n_{\max}, \\ \frac{n_{\min}}{n_\tau}, & \text{otherwise,} \end{cases} \quad (18)$$

$$D_{\text{temp}} := \langle |\psi(t) - \psi(t - \Delta t)|^2 \rangle. \quad (19)$$

$$D_{\text{clos}} := 1 - \langle \text{corr}(|\psi(t)|, |\psi(t - k\Delta t)|) \rangle_k. \quad (20)$$

Here  $n_\tau = \#\{(i, j) : |\psi_{i,j}|^2 > \tau\}$ ,  $N = L_x L_y$ , and  $(n_{\min}, n_{\max})$  define the Goldilocks zone for localization.

### B.2 Constraint-Conditioned Terms

**Strong (Confinement).** Add separation penalty  $\lambda \langle \log(1 + \text{sep}/d_0) \rangle$ .

**Electromagnetic (Long-Range).** Penalize  $\beta$  directly:  $\eta \beta$ ; bias exponent to  $n \approx 1$  by adding  $\kappa |n - 1|$ .

**Weak (Short-Range).** Reward screening: impose penalty for  $\beta < \beta_0$ ; maintain  $n$  near  $\sim 1$ .

**Gravity (Geometric).** Enforce curvature-energy matching  $\langle (K - c\rho)^2 \rangle$  with  $c > 0$ .

### B.3 Thresholds and Defaults

We choose practical defaults based on empirical stability:

- Localization threshold  $\tau$  for  $n_\tau$ : typically  $\tau \approx 0.5$  (dimensionless field intensity).
- Goldilocks zone:  $n_{\min} \approx 50$ ,  $n_{\max} \approx 500$  cells above threshold for coherent but nontrivial localization.
- Temporal lag samples for closure: last 20 frames, sampled every 5 steps.

### B.4 Optional Measures

Where gradient entropy is desired, an alternative inconsistency term uses entropy of  $|\nabla \psi|$  to penalize disorder without suppressing coherent gradients. This can be substituted for  $D_{\text{inc}}$  in Eq. (5) with minor computational overhead.

## B.5 Correlation Measurement with $\Phi$

For joint  $D$ – $\Phi$  measurement we compute  $D$  and a  $\Phi$  proxy on the same time series and report Pearson correlation, coefficient of determination, and significance; in bound-state runs we observed  $r \approx -0.91$  with  $p < 10^{-9}$  after autocorrelation correction.

# C Bootstrap and Annealing Algorithm

## C.1 Parameter Evolution

Temperature schedule  $T(t) = \max(T_{\min}, 1 - t/t_{\max})$ ; multiplicative random walks on  $\theta = \{\alpha, \beta, n, g, \gamma_{\text{base}}, \gamma_{\text{scale}}\}$  with exploration magnitude proportional to  $T$ ; projection to force-specific bounds each update.

## C.2 Regime-Specific Projections

After each annealing update, parameters are projected back to the regime-specific bounds listed in Appendix A. Exploration magnitude is reduced as  $T$  decreases, from  $\pm 20\%$  early to  $\pm 1\%$  late.

## C.3 Validation Metrics

Binding verification (final vs. initial separation, stability); discovered potential parameters  $(\alpha, \beta, n)$  or  $G$ ; minimum  $D$ ; optional  $D$ – $\Phi$  correlation with significance.

# D PR-0 Simulation Stills

The following frames are extracted from PR-0 simulation runs at three time points (early  $\approx 5\%$ , mid  $\approx 50\%$ , late  $\approx 95\%$  of run duration) for each of the ten simulation scenarios described in Section 4. They provide visual evidence of the spatial field evolution underlying the quantitative  $D$ -minimization results reported in the main text. Each row shows the PR-0 scalar field  $\psi$  rendered as a heat-map: bright regions indicate high-amplitude coherent structure (localized solitons); the progressive collapse of incoherence visible across frames is the spatial signature of  $D \rightarrow D_{\min}$ .




Figure 2: **EM-attraction soliton pair.** Two opposite-charge solitons initialised at separation; early frame shows distinct, weakly-interacting peaks; mid frame shows inward drift as  $D$  falls; late frame shows merged or tightly-bound configuration at  $D_{\min}$ . Run duration 40 s.




Figure 3: **EM-repulsion soliton pair.** Two like-charge solitons; frames show the progressive outward drift and thinning of the interaction channel. The final configuration achieves a well-separated, individually localized steady state consistent with minimized mutual incoherence. Run duration 40 s.




Figure 4: **EM-attraction three-body run.** Three solitons initialised at the vertices of a triangle; mid frame reveals complex interference pattern during approach; late frame shows a collapsed, tightly-localized cluster.  $D$  trace (cf. Figure 2 of main text) confirms monotone descent throughout. Run duration 120 s.




Figure 5: **EM-attraction orbital dynamics.** An asymmetric two-soliton initialisation produces orbital-like trajectories before settling; the spiralling inward path is visible in the mid frame. Emergent quasi-stable orbit is a local  $D$ -minimum before eventual capture. Run duration 133 s.




Figure 6: **EM-repulsion scattering.** Head-on collision trajectory; early frame shows approaching solitons; mid frame captures the high- $D$  deformation at closest approach; late frame shows clean post-scatter separation with individually re-localized solitons. Run duration 147 s.




Figure 7: **Soliton binding: first discovery run.** The original simulation in which spontaneous  $D$ -minimizing binding was first observed. Early frame shows two isolated solitons; mid frame shows field bridge formation; late frame shows stable bound state. This run established the foundational empirical evidence for the  $D$ -minimization hypothesis. Run duration 50 s.




Figure 8: **Strong-force capture.** Short-range confinement regime; early frame shows two initially well-separated solitons; mid frame shows rapid approach once within the strong-force range; late frame shows capture into a tightly confined bound state analogous to nucleon binding. Run duration 40 s.




Figure 9: **Emergent meson formation.** Quark-analogue soliton triplet collapses into a two-body meson-like bound state via  $D$ -minimization; mid frame captures the three-to-two transition; late frame shows the stable meson configuration. This is among the most direct visual demonstrations that composite particle structure is a natural  $D$ -minimum. Run duration 12.5 s.




Figure 10: **Unified all-forces, two-body multipanel.** A single run in which all four force regimes (gravity, EM attraction, EM repulsion, strong) are rendered as simultaneous panels for two solitons. The multipanel layout makes the regime-dependent  $D$ -descent rate directly visible: strong-force panel reaches  $D_{\min}$  fastest, gravitational panel slowest, consistent with coupling-strength ordering. Run duration 40 s.




Figure 11: **Unified all-forces, three-body.** The most complex scenario: three solitons under all four simultaneous force regimes. Mid frame shows highest- $D$  interference state; late frame shows a hierarchically organized configuration in which strong and EM interactions have resolved to local  $D$ -minima while the gravitational channel continues its slow descent. Run duration 53 s.