

The PSL(2,7) Algebraic Structure of the Generative Triple Evolution Framework

Nova Spivack

Independent Researcher

June 2026

<https://doi.org/10.5281/zenodo.20680887>

UGP series hub: <https://doi.org/10.5281/zenodo.20168144>

Classification: Mathematical Physics; Group Theory; Algebraic Number Theory; Standard Model Structure; Neutrino Flavor Symmetry.

Abstract

The Generative Triple Evolution (GTE) framework is built on three algebraic pillars: the Fibonacci–Möbius action on $\mathbb{P}^1(\mathbb{F}_7)$, the order-21 metacyclic group $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ (the Borel stabilizer of ∞ in $\mathrm{PSL}(2, 7)$), and the Eisenstein integers $\mathbb{Z}[\omega]$ ($\omega = e^{2\pi i/3}$, primitive cube root of unity). This paper proves that these three pillars are unified through $\mathrm{PSL}(2, 7)$, the unique simple group of order 168.

We establish five main results, all machine-certified in Lean 4 with zero `sorry`. First, F_{21} is the stabilizer of $\infty \in \mathbb{P}^1(\mathbb{F}_7)$ in $\mathrm{PSL}(2, 7)$ (the Borel subgroup of order 21), and together with the Fibonacci–Möbius generator, these two elements generate $\mathrm{PGL}(2, 7)$ of order 336 (`A_Lean; f21_is_borel_psl27, pgl27_generated_by_singer_and_borel`). Second, $\mathrm{PSL}(2, 7)$ is the unique simple group of order 168 and equals $\mathrm{Aut}(\mathcal{F})$ for the Fano plane $\mathcal{F}$, and F_{21} acts simply transitively on all 21 incidence flags of $\mathcal{F}$ (`A_Lean; psl27_is_aut_fano, f21_regular_on_fano_flags`). Third, the Eisenstein integers yield F_{21} and A_4 from the same functor: 7 splits as $7\mathbb{Z}[\omega] = \pi\bar{\pi}$ (norm-7 primes), giving $\mathbb{Z}[\omega]/(\pi) \cong \mathrm{GF}(7)$ and F_{21} from the unit group; 2 is inert ($2\mathbb{Z}[\omega]$ prime), giving $\mathbb{Z}[\omega]/(2) \cong \mathrm{GF}(4)$, whose additive group is $V_4 \cong \mathbb{Z}_2^2$ and whose multiplicative group is $\mathbb{Z}_3 = \mu_3$, assembling to $A_4 = V_4 \rtimes \mu_3$ (`A_Lean; eisenstein_a4_from_inert_2`). Fourth, the Klein quartic — the unique Hurwitz surface saturating $84(g-1)$ automorphisms for genus g — has genus equal to $N_{\mathrm{gen}} = 3$ (the number of Standard Model generations), arising from the $(2, 3, 7)$ triangle group shared with $\mathrm{PSL}(2, 7)$ (`A_Lean; klein_quartic_genus_eq_n_gen`). Fifth, the manifest GTE neutrino flavor symmetries — the generation cycling μ_3 and the μ – τ exchange $\mathbb{Z}_2$ — generate S_3 of order 6, which embeds in $A_4 = V_4 \rtimes \mu_3$ as a complement to the Klein normal subgroup supplied by the Eisenstein residue $\mathbb{Z}[\omega]/(2)$ (`A_Lean; gte_manifest_flavor_is_s3_in_a4`).

These five results establish a coherent group-theoretic skeleton: $\mathrm{PSL}(2, 7)$ contains F_{21} as its Borel subgroup, acts as the automorphism group of the Fano plane (with F_{21} as its regular flag-transitive subgroup), and is the automorphism group of the genus-3 Klein quartic. The shared μ_3 factor links F_{21} to A_4 via the Eisenstein functor, and the manifest $S_3 \leq A_4$ embedding (result 5) completes the leading-order neutrino flavor mixing group (**A/D**).

The physical interpretation — F_{21} as gauge skeleton group, A_4 as the GTE leading-order neutrino flavor symmetry yielding tri-bimaximal mixing angles, and the Klein quartic as the geometric reason for three generations — is argued at level **A/D** or **D** as stated.

Contents

1	Introduction	3
2	Mathematical Setup: $\text{PSL}(2, 7)$ and F_{21}	4
2.1	The projective line $\mathbb{P}^1(\mathbb{F}_7)$	4
2.2	The Fibonacci–Möbius generator	4
2.3	The metacyclic group F_{21}	5
3	The Fano Plane and the Regular Flag Action	5
3.1	The Fano plane	5
3.2	Regular action of F_{21} on Fano flags	5
4	The Eisenstein Functor: F_{21} and A_4 from $\mathbb{Z}[\omega]$	6
4.1	Eisenstein integers	6
4.2	The split prime 7 and F_{21}	6
4.3	The inert prime 2 and A_4	6
4.4	Amalgamated structure and flavor embedding	6
4.5	GTE manifest flavor symmetry and A_4	7
5	Klein Quartic Geometry and Three Generations	7
5.1	Hurwitz’s theorem and the $(2, 3, 7)$ triangle group	7
5.2	The Klein quartic	7
6	Physical Interpretation	8
6.1	F_{21} as gauge skeleton $(\mathbf{A}/\mathbf{D})$	8
6.2	Fano flags as a coordinate system $(\mathbf{A}/\mathbf{D})$	8
6.3	A_4 and neutrino flavor symmetry $(\mathbf{A}/\mathbf{D})$	8
6.4	Klein quartic and the geometry of three generations $(\mathbf{D})$	8
7	Conclusion	8
	Appendices	9
A	Lean Certification Details	9
B	The $\text{PSL}(2, 7)$ Subgroup Lattice (Summary)	10

1 Introduction

The Generative Triple Evolution framework derives the Standard Model generation structure from a cellular automaton over $\text{GF}(7)$ governed by the polynomial $p(L, C, R) = C + R - CR - LCR$. The algebraic architecture of GTE is built on three recurring objects:

- the **Fibonacci–Möbius map** $\mu : x \mapsto (x + 1)/x$ acting on $\mathbb{P}^1(\mathbb{F}_7) = \mathbb{F}_7 \cup \{\infty\}$, whose cyclic orbit structure encodes the GTE generation sequence;
- the **metacyclic group** $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ of order 21, identified in prior work [2] as the GTE gauge skeleton group; and
- the **Eisenstein integers** $\mathbb{Z}[\omega] = \mathbb{Z}[e^{2\pi i/3}]$, which provide the arithmetic bridge between the GTE polynomial’s $\mathbb{Z}_7$ structure and the Standard Model flavor groups.

The algebraic structure of the GTE $\mathbb{Z}_7$ polynomial and its encoding of Standard Model quantum numbers is established in [3]. The full two-level framework — Level 1 algebraic certificate (p) and Level 2 continuum field ($\mathcal{L}_{\Phi_{\text{MDL}}}$) — is derived in P48 [1]; the present paper operates entirely at Level 1 (the algebraic skeleton) and does not derive the field law.

This paper shows that these three pillars are not independent constructions but aspects of a single algebraic object: $\text{PSL}(2, 7)$, the unique simple group of order 168. The group F_{21} is the Borel subgroup of $\text{PSL}(2, 7)$ — the stabilizer of ∞ in the action on $\mathbb{P}^1(\mathbb{F}_7)$. The Fibonacci–Möbius map together with F_{21} generate the full $\text{PGL}(2, 7)$ of order 336. The Eisenstein functor $\mathbb{Z}[\omega] \rightarrow \text{Res}_q$ applied at $q = 7$ (split prime) yields $\text{GF}(7)$ and F_{21} ; applied at $q = 2$ (inert prime) yields $\text{GF}(4)$ and A_4 , completing the GTE neutrino flavor symmetry to tri-bimaximal mixing. The same $(2, 3, 7)$ triangle group underlying $\text{PSL}(2, 7)$ gives the Klein quartic — the unique Hurwitz surface with $\text{Aut} = \text{PSL}(2, 7)$ — whose genus equals $N_{\text{gen}} = 3$.

The key theorem names and certification levels are collected in Table 1 for reference.

Claim-Strength Taxonomy

Every result in this paper carries one of five certification levels:

- **A_{Lean} (A-Lean)** — Lean 4 machine proof, zero **sorry**, zero custom axioms. Conditional **A_{Lean}** uses a named physics axiom (always stated explicitly).
- **A/D (A/D)** — Analytic or fully decidable proof; human-verifiable.
- **A (A)** — Computationally verified; reproducible from published scripts.
- **B (B)** — Theoretically motivated conjecture with numerical support.
- **D (D)** — Speculative hypothesis; further development needed.

Ordering: **A_{Lean} > A/D > A > B > D**. All Lean certifications use the canonical **ugp-lean** repository; module paths are given in Appendix A.

Table 1: Main results of this paper.

Result	Level	Lean theorem
F_{21} is Borel subgroup of $\mathrm{PSL}(2, 7)$, order 21	$\mathbf{A}_{\text{Lean}}$	f21_is_borel_psl27
$\langle \mu, F_{21} \rangle = \mathrm{PGL}(2, 7)$, order 336	$\mathbf{A}_{\text{Lean}}$	pgl27_generated_by_singer_and_borel
$ \mathrm{PSL}(2, 7) = 168$, unique simple group of this order	$\mathbf{A}_{\text{Lean}}$	psl27_is_aut_fano
F_{21} acts regularly on all 21 Fano flags	$\mathbf{A}_{\text{Lean}}$	f21_regular_on_fano_flags
$A_4 = V_4 \rtimes \mathbb{Z}_3$ from $\mathbb{Z}[\omega]/(2)$	$\mathbf{A}_{\text{Lean}}$	eisenstein_a4_from_inert_2
Manifest GTE flavor $S_3 \leq A_4 = V_4 \rtimes \mu_3$	$\mathbf{A}_{\text{Lean}}$	gte_manifest_flavor_is_s3_in_a4
genus(Klein quartic) = $N_{\text{gen}} = 3$	$\mathbf{A}_{\text{Lean}}$	klein_quartic_genus_eq_n_gen
$G_{84} = F_{21} \times_{\mathbb{Z}_3} A_4 \not\hookrightarrow \mathrm{GL}(2, 7)$	$\mathbf{A}$	(computational)
Klein quartic orbifold formula $\chi = 168 \times (-1/42) = -4$	$\mathbf{A}$	(computational)
A_4 as GTE leading-order neutrino flavor symmetry (TBM)	$\mathbf{A}/\mathbf{D}$	see §6
Klein quartic genus = N_{gen} from Gauss–Bonnet	$\mathbf{A}/\mathbf{D}$	see §5

2 Mathematical Setup: $\mathrm{PSL}(2, 7)$ and F_{21}

2.1 The projective line $\mathbb{P}^1(\mathbb{F}_7)$

Let $\mathbb{F}_7 = \mathrm{GF}(7) = \{0, 1, 2, 3, 4, 5, 6\}$ with arithmetic modulo 7. The projective line $\mathbb{P}^1(\mathbb{F}_7) = \mathbb{F}_7 \cup \{\infty\}$ has $|\mathbb{P}^1(\mathbb{F}_7)| = 8$ points. The group $\mathrm{PGL}(2, 7)$ acts on $\mathbb{P}^1(\mathbb{F}_7)$ by Möbius transformations:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot x = \frac{ax + b}{cx + d}, \quad ad - bc \neq 0 \pmod{7}, \quad (1)$$

with $\infty \mapsto a/c$ when $c \neq 0$, and $\infty \mapsto \infty$ when $c = 0$. The subgroup $\mathrm{PSL}(2, 7) \leq \mathrm{PGL}(2, 7)$ consists of matrices with $\det = 1$ modulo identification $M \sim -M$, giving $|\mathrm{PSL}(2, 7)| = |\mathrm{SL}(2, 7)|/2 = 168$.

2.2 The Fibonacci–Möbius generator

The **Fibonacci–Möbius map** is the transformation $\mu : x \mapsto (x + 1)/x$, represented by the matrix

$$M_\mu = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in \mathrm{GL}(2, 7). \quad (2)$$

The sequence $\mu^n(\infty)$ reproduces the Fibonacci sequence modulo 7: since $\mu(\infty) = 1$, $\mu^2(\infty) = 2$, $\mu^3(\infty) = 3/2$, etc., and the periods are governed by the Pisano period $\pi(7) = 16$. In the GTE framework, μ is the map whose orbit on $\mathbb{P}^1(\mathbb{F}_7)$ encodes the generation-changing dynamics [1, 2].

2.3 The metacyclic group F_{21}

The **affine group** $\text{AGL}(1, 7) = \mathbb{F}_7 \rtimes \mathbb{F}_7^\times$ has order $7 \times 6 = 42$ and acts on $\mathbb{F}_7$ by $x \mapsto ax + b$ ($a \neq 0$). Its Sylow-3 subgroup is the **metacyclic group**

$$F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3, \quad |F_{21}| = 21, \quad (3)$$

generated by the translation $\tau : x \mapsto x + 1$ and the multiplication $\sigma : x \mapsto 2x$ (note $2^3 = 8 \equiv 1 \pmod{7}$, so σ has order 3). This group is identified in the GTE framework as the gauge skeleton group whose orbit structure encodes the $N_{\text{gen}} = 3$ generations [2].

Theorem 2.1 (`ALean — f21_is_borel_psl27`). *The stabilizer of $\infty \in \mathbb{P}^1(\mathbb{F}_7)$ in $\text{PSL}(2, 7)$ consists of all upper-triangular matrices $\begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}$ with $a^3 = 1$ in $\mathbb{F}_7^\times$, and this subgroup is isomorphic to $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ of order 21.*

Theorem 2.2 (`ALean — pgl27_generated_by_singer_and_borel`). *Let μ denote the Fibonacci–Möbius action on $\mathbb{P}^1(\mathbb{F}_7)$ and let F_{21} be generated by the translation $x \mapsto x + 1$ and multiplication $x \mapsto 2x$. Then $\langle \mu, F_{21} \rangle = \text{PGL}(2, 7)$, a group of order 336.*

Remark 2.3. Theorem 2.2 means that the GTE’s two fundamental building blocks — the Fibonacci–Möbius dynamics and the F_{21} gauge skeleton — are not independent; together they generate the full symmetry group of $\mathbb{P}^1(\mathbb{F}_7)$. The index $[\text{PGL}(2, 7) : \text{PSL}(2, 7)] = 2$ reflects the outer automorphism of $\text{PSL}(2, 7)$ corresponding to field conjugation.

3 The Fano Plane and the Regular Flag Action

3.1 The Fano plane

The **Fano plane** $\mathcal{F} = \text{PG}(2, \mathbb{F}_2)$ is the unique projective plane of order 2, with 7 points and 7 lines, each line containing 3 points and each point on 3 lines. An **incidence flag** is a pair (p, L) where p is a point of $\mathcal{F}$ and L is a line incident to p . The Fano plane has $7 \times 3 = 21$ incidence flags.

Theorem 3.1 (`ALean — psl27_is_aut_fano`). *$\text{PSL}(2, 7)$ is the unique simple group of order 168, and it acts as the full automorphism group of the Fano plane: $\text{Aut}(\mathcal{F}) \cong \text{PSL}(2, 7)$.*

3.2 Regular action of F_{21} on Fano flags

Theorem 3.2 (`ALean — f21_regular_on_fano_flags`). *Encode the Fano plane points as $\mathbb{Z}_7$, lines as translates of the Singer difference set $\{0, 1, 3\}$, and flags as incidence pairs (p, L) . Define the F_{21} action by $\sigma_{(a,k)} : x \mapsto 2^k x + a \pmod{7}$ for $(a, k) \in \mathbb{Z}_7 \times \mathbb{Z}_3$. Then F_{21} acts simply transitively (regularly) on all 21 Fano incidence flags: for any two flags f, f' , there is a unique $g \in F_{21}$ with $g \cdot f = f'$.*

Remark 3.3. Simple transitivity means $|F_{21}| = 21$ equals the number of flags 21 and the action is both transitive (orbit = all flags) and free (no element fixes a flag). Geometrically, F_{21} is a Singer cycle complement that provides a *coordinate system* on the Fano plane: every flag is reachable from any reference flag by a unique gauge transformation. In GTE, this identifies F_{21} as the group of gauge configurations over the $\mathbb{Z}_7$ field.

4 The Eisenstein Functor: F_{21} and A_4 from $\mathbb{Z}[\omega]$

4.1 Eisenstein integers

Let $\omega = e^{2\pi i/3}$ be a primitive cube root of unity. The **Eisenstein integers** $\mathbb{Z}[\omega] = \mathbb{Z}[\omega]$ form a principal ideal domain with unit group $\{\pm 1, \pm\omega, \pm\omega^2\} \cong \mathbb{Z}_6$ and norm $N(a + b\omega) = a^2 - ab + b^2$. The factorization of a rational prime q in $\mathbb{Z}[\omega]$ depends on $q \pmod{3}$:

- $q \equiv 1 \pmod{3}$: **splits** as $q\mathbb{Z}[\omega] = \pi\bar{\pi}$ (two distinct conjugate primes of norm q).
- $q \equiv 2 \pmod{3}$: **inert** (stays prime in $\mathbb{Z}[\omega]$, giving residue field $\text{GF}(q^2)$).
- $q = 3$: **ramifies** as $3\mathbb{Z}[\omega] = (1 - \omega)^2 \cdot \epsilon$.

4.2 The split prime 7 and F_{21}

Since $7 \equiv 1 \pmod{3}$, the prime 7 splits in $\mathbb{Z}[\omega]$ as $7\mathbb{Z}[\omega] = \pi\bar{\pi}$ for a norm-7 Eisenstein prime π . The residue ring $\mathbb{Z}[\omega]/(\pi) \cong \text{GF}(7)$, and the unit group $(\mathbb{Z}[\omega]/(\pi))^\times = \text{GF}(7)^\times \cong \mathbb{Z}_6$. The subgroup of cube roots of unity $\mu_3 = \{1, \omega, \omega^2\}$ reduces to the unique index-2 subgroup $\mathbb{Z}_3 \leq \text{GF}(7)^\times$, generated by $2 \in \text{GF}(7)^\times$ (the element of order 3). Combined with the additive group $\text{GF}(7)^+ = \mathbb{Z}_7$, this gives the affine action of $F_{21} = \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ on $\text{GF}(7)$.

4.3 The inert prime 2 and A_4

Since $2 \equiv 2 \pmod{3}$, the prime 2 is inert in $\mathbb{Z}[\omega]$: $2\mathbb{Z}[\omega]$ remains a prime ideal, and the residue ring $\mathbb{Z}[\omega]/(2) \cong \text{GF}(4)$, the field of four elements.

Theorem 4.1 (**A**_{Lean} — `eisenstein_a4_from_inert_2`). *The Eisenstein residue at the inert prime 2 gives:*

- (i) $\mathbb{Z}[\omega]/(2) \cong \text{GF}(4)$; the additive group $(\text{GF}(4), +) \cong V_4 = \mathbb{Z}_2^2$ and the multiplicative group $\text{GF}(4)^\times \cong \mathbb{Z}_3 = \mu_3$.
- (ii) The semidirect product $V_4 \rtimes \mu_3 \cong A_4$ (the alternating group on 4 letters, order 12).

Remark 4.2. The $\mu_3 = \mathbb{Z}_3$ factor is shared between F_{21} (from the split prime 7) and A_4 (from the inert prime 2). The physical identification of this shared cyclic subgroup with $\text{SU}(2)_L^{\text{cyc}}$ as the bridge between the GTE gauge skeleton and the flavor sector is argued at level **D**; see §6.

4.4 Amalgamated structure and flavor embedding

The two Eisenstein residues combine into an amalgamated product:

$$G_{84} = F_{21} \times_{\mu_3} A_4 = (\mathbb{Z}_7 \times V_4) \rtimes \mathbb{Z}_3, \quad |G_{84}| = 84. \quad (4)$$

The value $|G_{84}| = 84 = |\text{PSL}(2, 7)|/2$ is half the automorphism count of the Klein quartic. A computational check (**A**) confirms that G_{84} does not embed in $\text{GL}(2, 7)$ or $\text{PGL}(2, 7)$, so the flavor-gauge amalgamation is genuinely a larger structure than the gauge group alone.

4.5 GTE manifest flavor symmetry and A_4

Theorem 4.3 ($\mathbf{A}_{\text{Lean}} - \text{gte_manifest_flavor_is_s3_in_a4}$). *The GTE manifest neutrino flavor symmetries — the generation cycling $\mu_3 = \langle (123) \rangle$ and the μ - τ exchange $U = (23)$ — generate the symmetric group S_3 of order 6. The group S_3 embeds in $A_4 = V_4 \rtimes \mu_3$ as a complement to the Klein normal subgroup V_4 , which is supplied by the Eisenstein residue $\mathbb{Z}[\omega]/(2)$.*

Remark 4.4. Theorem 4.3 states a group-theoretic fact ($\mathbf{A}_{\text{Lean}}$). The identification of A_4 as the GTE leading-order neutrino flavor symmetry yielding tri-bimaximal (TBM) mixing angles $\sin^2 \theta_{12} = 1/3$, $\sin^2 \theta_{23} = 1/2$, $\theta_{13} = 0$ is an analytic derivation ($\mathbf{A/D}$) and is discussed in §6.

5 Klein Quartic Geometry and Three Generations

5.1 Hurwitz's theorem and the $(2, 3, 7)$ triangle group

Hurwitz's theorem states that a compact Riemann surface of genus $g \geq 2$ has at most $84(g - 1)$ conformal automorphisms. The bound is achieved if and only if the surface is a quotient of the hyperbolic plane $\mathbb{H}$ by a torsion-free normal subgroup of the $(2, 3, 7)$ triangle group $\Delta(2, 3, 7) = \langle x, y, z \mid x^2 = y^3 = z^7 = xyz = 1 \rangle$.

The **orbifold Euler characteristic** of $\Delta(2, 3, 7)$ is

$$\chi_{\text{orb}} = 1 - \frac{1}{2} - \frac{1}{3} - \frac{1}{7} = 1 - \frac{21}{42} - \frac{14}{42} - \frac{6}{42} = -\frac{1}{42}. \quad (5)$$

This is the minimum possible orbifold area $\pi/21$ for a hyperbolic orbifold, by Siegel's theorem.

5.2 The Klein quartic

The **Klein quartic** is the unique compact Riemann surface of genus 3 that attains the Hurwitz bound $84(g - 1) = 168$ automorphisms, with $\text{Aut}(\text{Klein quartic}) \cong \text{PSL}(2, 7)$. It can be defined as the smooth projective curve $x^3y + y^3z + z^3x = 0$ in $\mathbb{P}^2(\mathbb{C})$.

Theorem 5.1 ($\mathbf{A}_{\text{Lean}} - \text{klein_quartic_genus_eq_n_gen}$). *The orbifold covering gives:*

- (i) *Orbifold Euler characteristic:* $\chi_{\text{orb}}(\Delta(2, 3, 7)) = -1/42$ (machine-verified by **norm_num**).
- (ii) *Surface Euler characteristic:* $\chi(\text{Klein}) = |\text{PSL}(2, 7)| \times \chi_{\text{orb}} = 168 \times (-1/42) = -4$ (machine-verified by **norm_num**).
- (iii) *Genus formula:* $g = (2 - \chi)/2 = (2 - (-4))/2 = 3 = N_{\text{gen}}$ (machine-verified by **norm_num**).

The Klein quartic is a Hurwitz surface with $|\text{Aut}| = 84(g - 1) = 168 = |\text{PSL}(2, 7)|$ (saturation verified).

Corollary 5.2 ($\mathbf{A/D}$). *The genus of the unique Hurwitz surface whose automorphism group contains $\text{PSL}(2, 7)$ equals the number of Standard Model generations, $N_{\text{gen}} = 3$. This is a consequence of Theorems 3.1 and 5.1 together with Hurwitz's theorem: $\text{PSL}(2, 7)$ is characterized by both $\text{Aut}(\mathcal{F}) = \text{PSL}(2, 7)$ (Fano plane automorphisms) and $\text{Aut}(\text{Klein quartic}) = \text{PSL}(2, 7)$, and the genus-3 surface is the unique Hurwitz surface in genus 2 or higher with an automorphism group isomorphic to $\text{PSL}(2, 7)$.*

Remark 5.3. Corollary 5.2 does not in itself explain *why* $N_{\text{gen}} = 3$ physically — it shows that N_{gen} and the genus of the Klein quartic are the same number (both equal to 3) and that this equality follows from the same algebraic structure ($\text{PSL}(2, 7)$ and the $(2, 3, 7)$ triangle group) that underlies the GTE framework. The physical derivation of $N_{\text{gen}} = 3$ from GTE first principles is given in [1, 2].

6 Physical Interpretation

This section describes the proposed physical meaning of the algebraic results proved above. The group-theoretic statements are $\mathbf{A}_{\text{Lean}}$; the physical interpretations carry level $\mathbf{A/D}$ or $\mathbf{D}$ as stated.

6.1 F_{21} as gauge skeleton ($\mathbf{A/D}$)

In GTE, the cellular automaton update map on $\text{GF}(7)^{\mathbb{Z}}$ generates orbit families indexed by $\mathbb{Z}_7$ sector values. The F_{21} action on $\text{GF}(7)$ by $x \mapsto 2^k x + a$ is the gauge freedom of the CMCA tape: it maps one valid tape configuration to another without changing the physical content. The identification $F_{21} = \text{Stab}_{\text{PSL}(2,7)}(\infty)$ (Theorem 2.1) places this gauge group naturally inside the full $\text{PSL}(2,7)$ symmetry.

6.2 Fano flags as a coordinate system ($\mathbf{A/D}$)

Theorem 3.2 shows that F_{21} acts simply transitively on the 21 Fano flags. Since $|F_{21}| = 21$ equals the number of flags, F_{21} provides a *bijection* between the abstract gauge group and the set of all Fano incidence data. This means that every consistent assignment of $\mathbb{Z}_7$ field values to the three GTE roles (Left, Center, Right) corresponds to a unique flag, and the 21-element gauge orbit is the same as the 21-element flag set.

6.3 A_4 and neutrino flavor symmetry ($\mathbf{A/D}$)

The Majorana neutrino mass matrix in GTE is constructed from the orbit of the $\text{GF}(7)$ polynomial under the combined Eisenstein residue symmetry. The leading-order mixing is governed by $A_4 = V_4 \rtimes \mu_3$, where μ_3 is the generation cycling symmetry (already manifest in GTE) and V_4 is the additional discrete symmetry supplied by the Eisenstein residue at the inert prime 2. The A_4 symmetry forces the **tri-bimaximal mixing** (TBM) pattern:

$$\sin^2 \theta_{12} = \frac{1}{3}, \quad \sin^2 \theta_{23} = \frac{1}{2}, \quad \theta_{13} = 0 \quad (6)$$

at leading order [1]. Corrections from breaking of $A_4 \rightarrow S_3$ (the manifest part retained in GTE) account for the observed nonzero θ_{13} ($\mathbf{D}$; open problem).

6.4 Klein quartic and the geometry of three generations ($\mathbf{D}$)

The identification $\text{genus}(\text{Klein quartic}) = N_{\text{gen}} = 3$ (Theorem 5.1) suggests a geometric interpretation: the three Standard Model generations correspond to the three handles of the genus-3 Riemann surface that appears when the $(2, 3, 7)$ Hurwitz structure is realized as a surface. Whether this geometric correspondence has a direct physical realization in the GTE CMCA dynamics is currently open ($\mathbf{D}$).

7 Conclusion

We have proved that the three principal algebraic structures of the GTE framework — the Fibonacci–Möbius map, the metacyclic gauge skeleton group F_{21} , and the Eisenstein integer arithmetic — are all facets of a single algebraic object: $\text{PSL}(2, 7)$, the unique simple group of order 168.

The certified results ($\mathbf{A}_{\text{Lean}}$) are:

1. F_{21} is the Borel subgroup of $\mathrm{PSL}(2, 7)$, and $\mathrm{PSL}(2, 7)$ itself is generated by F_{21} together with the Fibonacci–Möbius map (Theorems 2.1 and 2.2).
2. $\mathrm{PSL}(2, 7) \cong \mathrm{Aut}(\mathcal{F})$ and F_{21} acts simply transitively on the 21 incidence flags of the Fano plane (Theorems 3.1 and 3.2).
3. The Eisenstein functor yields F_{21} and A_4 from the single ring $\mathbb{Z}[\omega]$, with a shared μ_3 bridge (Theorem 4.1).
4. The manifest GTE flavor symmetries μ_3 and the μ – τ exchange generate $S_3 \leq A_4$ (Theorem 4.3).
5. $\mathrm{genus}(\text{Klein quartic}) = N_{\mathrm{gen}} = 3$ (Theorem 5.1).

The analytic corollaries (**A/D**) establish that the A_4 residue symmetry from the Eisenstein functor is the leading-order GTE neutrino flavor group completing manifest S_3 to the TBM-generating A_4 , and that the Klein quartic genus equals N_{gen} as a consequence of the $(2, 3, 7)$ covering structure.

Open Problem 1. Derive the magnitude of the corrections to the TBM mixing pattern from the breaking $A_4 \rightarrow S_3$ within the GTE Eisenstein framework.

Open Problem 2. Determine whether the embedding $G_{84} = F_{21} \times_{\mu_3} A_4$ can be realized as a physical symmetry group of the CMCA tape dynamics, or whether it appears only as a structural artifact of the Eisenstein functor.

Open Problem 3. Establish whether the Klein quartic has a direct realization in the GTE geometric structure — for instance, as the moduli space of PSC-admissible field configurations in genus 3, or as the quotient $\mathbb{H}/\Gamma$ for some natural lattice Γ arising from the $\mathrm{GF}(7)$ dynamics.

A Lean Certification Details

All theorems labeled **A_{Lean}** in this paper are machine-verified in Lean 4 using the Mathlib4 library, with zero `sorry` and zero custom axioms. The certifications reside in the canonical public `ugp-lean` repository.

Table 2: Lean theorem locations.

Theorem name	Module	Cat
<code>f21_is_borel_psl27</code>	<code>Polynomial/PSL27Unification.lean</code>	A_{Lean}
<code>pgl27_generated_by_singer_and_borel</code>	<code>Polynomial/PSL27Unification.lean</code>	A_{Lean}
<code>psl27_is_aut_fano</code>	<code>Polynomial/PSL27Unification.lean</code>	A_{Lean}
<code>f21_regular_on_fano_flags</code>	<code>Algebra/FanoRegularAction.lean</code>	A_{Lean}
<code>eisenstein_a4_from_inert_2</code>	<code>Algebra/EisensteinFunctor.lean</code>	A_{Lean}
<code>gte_manifest_flavor_is_s3_in_a4</code>	<code>Algebra/FlavorGroupStructure.lean</code>	A_{Lean}
<code>klein_quartic_genus_eq_n_gen</code>	<code>Cosmology/CCBracketHurwitz.lean</code>	A_{Lean}

To verify, build the `ugp-lean` repository:

```
cd /path/to/ugp-lean
lake build UgpLean
```

A clean build with no errors and no `sorry` warnings confirms all certifications.

B The $\text{PSL}(2,7)$ Subgroup Lattice (Summary)

For reference, the key subgroup relations used in this paper are:

$$\mathbb{Z}_7 \leq F_{21} \leq \text{PSL}(2, 7) \leq \text{PGL}(2, 7) \quad (7)$$

with $|\mathbb{Z}_7| = 7$, $|F_{21}| = 21$, $|\text{PSL}(2, 7)| = 168$, $|\text{PGL}(2, 7)| = 336$, and $[\text{PGL}(2, 7) : \text{PSL}(2, 7)] = 2$. The Eisenstein residues yield:

$$\mu_3 \leq F_{21} \leq \text{PSL}(2, 7), \quad \mu_3 \leq A_4, \quad S_3 \leq A_4, \quad (8)$$

where $|\mu_3| = 3$, $|S_3| = 6$, $|A_4| = 12$, and A_4 is the leading-order GTE neutrino flavor group ($\mathbf{A/D}$).

References

- [1] Nova Spivack. The complete GTE framework: Standard Model, gravity, quantum mechanics, and cosmology from ϕ_{mdl} . Zenodo, 2026. <https://doi.org/10.5281/zenodo.20560550>. Preprint.
- [2] Nova Spivack. The GTE-Möbius substrate: Arithmetic unification of computation and transputation in the universal generative principle. Zenodo, 2026. Paper P34 in the UGP Physics series. <https://doi.org/10.5281/zenodo.20319419>.
- [3] Nova Spivack. MDL selects the Wolfram rule: $\mathbb{Z}_7$ dynamics, algebraic structure, and Standard Model encoding of the GTE polynomial. Preprint, 2026. <https://doi.org/10.5281/zenodo.20572656>. UGP Physics series, P49.