



American Diabetes Association
Professional Practice Committee*

10. Cardiovascular Disease and Risk Management: Standards of Care in Diabetes—2024

Diabetes Care 2024;47(Suppl. 1):S179–S218 | <https://doi.org/10.2337/dc24-S010>

The American Diabetes Association (ADA) “Standards of Care in Diabetes” includes the ADA’s current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. Members of the ADA Professional Practice Committee, an interprofessional expert committee, are responsible for updating the Standards of Care annually, or more frequently as warranted. For a detailed description of ADA standards, statements, and reports, as well as the evidence-grading system for ADA’s clinical practice recommendations and a full list of Professional Practice Committee members, please refer to **Introduction and Methodology**. Readers who wish to comment on the Standards of Care are invited to do so at professional.diabetes.org/SOC.

For prevention and management of diabetes complications in children and adolescents, please refer to Section 14, “Children and Adolescents.”

Atherosclerotic cardiovascular disease (ASCVD)—defined as coronary heart disease (CHD), cerebrovascular disease, or peripheral artery disease (PAD) presumed to be of atherosclerotic origin—is the leading cause of morbidity and mortality for individuals with diabetes and results in an estimated \$39.4 billion in cardiovascular-related spending per year associated with diabetes (1). Common conditions coexisting with type 2 diabetes (e.g., hypertension and dyslipidemia) are clear risk factors for ASCVD, and diabetes itself confers independent risk. Numerous studies have shown the efficacy of controlling individual cardiovascular risk factors in preventing or slowing ASCVD in people with diabetes. Furthermore, large benefits are seen when multiple cardiovascular risk factors (glycemic, blood pressure, and lipid control) are addressed simultaneously, with evidence for legacy benefits (2–4). Under the current paradigm of aggressive risk factor modification in people with diabetes, there is evidence that measures of 10-year CHD risk among U.S. adults with diabetes have improved significantly over the past decade (5) and that ASCVD morbidity and mortality have decreased (3,6).

Heart failure is another major cause of morbidity and mortality from cardiovascular disease. The American Diabetes Association (ADA) has developed a consensus report to summarize guidance for the screening, diagnosis, and treatment of people with diabetes (7). Recent studies have found that rates of incident heart failure hospitalization (adjusted for age and sex) were twofold higher in people with diabetes compared with those without (8,9). People with diabetes may present with a wide spectrum of heart failure, including heart failure with preserved ejection fraction (HFpEF), heart failure with mildly reduced ejection fraction (HFrEF), or heart failure with reduced ejection fraction (HFrEF). Hypertension is often a precursor of heart failure of either type, and ASCVD can coexist with either type of heart failure (10), whereas prior myocardial infarction (MI) is often a major factor in HFrEF. Recent trials including people with type 2 diabetes, most of whom also had ASCVD, have shown

*A complete list of members of the American Diabetes Association Professional Practice Committee can be found at <https://doi.org/10.2337/dc24-SINT>.

Duality of interest information for each author is available at <https://doi.org/10.2337/dc24-SDIS>.

This section has received endorsement from the American College of Cardiology.

Suggested citation: American Diabetes Association Professional Practice Committee. 10. Cardiovascular disease and risk management: Standards of Care in Diabetes—2024. *Diabetes Care* 2024;47(Suppl. 1):S179–S218

© 2023 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

that rates of heart failure hospitalization significantly decreased with use of sodium–glucose cotransporter 2 (SGLT2) inhibitors (11–14).

A recent meta-analysis indicated that SGLT2 inhibitors reduce the risk of heart failure hospitalization, cardiovascular mortality, and all-cause mortality in people with (secondary prevention) and without (primary prevention) cardiovascular disease (15).

For prevention and management of both ASCVD and heart failure, cardiovascular risk factors should be systematically assessed at least annually in all people with diabetes. These risk factors include duration of diabetes, obesity/overweight, hypertension, dyslipidemia, smoking, a family history of premature coronary disease, chronic kidney disease (CKD), and the presence of albuminuria. Modifiable abnormal risk factors should be treated as described in these guidelines. Notably, the majority of evidence supporting interventions to reduce cardiovascular risk in diabetes comes from trials of people with type 2 diabetes. No randomized trials have been specifically designed to assess the impact of cardiovascular risk reduction strategies in people with type 1 diabetes. Therefore, the recommendations for cardiovascular risk

factor modification for people with type 1 diabetes are extrapolated from data obtained in people with type 2 diabetes and are similar to those for people with type 2 diabetes.

As depicted in Fig. 10.1, a comprehensive approach to the reduction in risk of diabetes-related complications is recommended. Therapy that includes multiple, concurrent evidence-based approaches to care will provide complementary reduction in the risks of microvascular outcomes, including kidney, retinopathy, neurologic, and cardiovascular complications. Management of glycemia, blood pressure, and lipids and the incorporation of specific therapies with cardiovascular and kidney outcomes benefit (as individually appropriate) are considered fundamental elements of global risk reduction in diabetes.

THE RISK CALCULATOR

The American College of Cardiology ASCVD risk calculator (Risk Estimator Plus) is generally a useful tool to estimate 10-year risk of a first ASCVD event (available online at tools.acc.org/ASCVD-Risk-Estimator-Plus). The calculator was developed to stratify cardiovascular risk and identify those people who will benefit

most from statin therapy and from treatment with antihypertensive medications (16). The calculator includes diabetes as a risk factor, since diabetes itself confers increased risk for ASCVD, although it should be acknowledged that these risk calculators do not account for the duration of diabetes or the presence of diabetes complications, such as albuminuria. In addition, the majority of people with diabetes should be treated with statin therapy, and hypertension should be promptly treated. As we will discuss below, comprehensive management of hypertension, hyperlipidemia, and hyperglycemia using management approaches with established benefit are important strategies to reduce cardiovascular risk.

HYPERTENSION/BLOOD PRESSURE CONTROL

An elevated blood pressure is defined as a systolic blood pressure 120–129 mmHg and a diastolic blood pressure <80 mmHg (17). Hypertension is defined as a systolic blood pressure ≥130 mmHg or a diastolic blood pressure ≥80 mmHg (17). This is in agreement with the definition of hypertension by the American College of Cardiology and American Heart Association (17). Hypertension is common among people with either type 1 or type 2 diabetes. Hypertension is a major risk factor for ASCVD, heart failure, and microvascular complications. Moreover, numerous studies have shown that antihypertensive therapy reduces ASCVD events, heart failure, and microvascular complications. Please refer to the ADA position statement “Diabetes and Hypertension” for a detailed review of the epidemiology, diagnosis, and treatment of hypertension (18) and recent updated hypertension guideline recommendations (17,19,20).




Figure 10.1—Multifactorial approach to reduction in risk of diabetes complications. *Risk reduction interventions to be applied as individually appropriate.

Screening and Diagnosis

Recommendations

- 10.1 Blood pressure should be measured at every routine clinical visit. When possible, individuals found to have elevated blood pressure (systolic blood pressure 120–129 mmHg and diastolic <80 mmHg) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. **A** Hypertension is defined as a systolic blood pressure ≥130 mmHg or a diastolic blood pressure ≥80 mmHg based

on an average of two or more measurements obtained on two or more occasions. **A** Individuals with blood pressure $\geq 180/110$ mmHg and cardiovascular disease could be diagnosed with hypertension at a single visit. **E**

10.2 All people with hypertension and diabetes should be counseled to monitor their blood pressure at home after appropriate education. **A**

Blood pressure should be measured at every routine clinical visit by a trained individual and should follow the guidelines established for the general population: measurement in the seated position, with feet on the floor and arm supported at heart level, after 5 min of rest. Cuff size should be appropriate for the upper-arm circumference (21). Elevated values should preferably be confirmed on a separate day; however, in individuals with cardiovascular disease and blood pressure $\geq 180/110$ mmHg, it is reasonable to diagnose hypertension at a single visit (19). Postural changes in blood pressure and pulse may be evidence of autonomic neuropathy and therefore require adjustment of blood pressure targets. Orthostatic blood pressure measurements should be checked on initial visit and as indicated.

Home blood pressure self-monitoring and 24-h ambulatory blood pressure monitoring may provide evidence of white coat hypertension, masked hypertension, or other discrepancies between office and “true” blood pressure (22,23). In addition to confirming or refuting a diagnosis of hypertension, home blood pressure assessment may be useful to monitor antihypertensive treatment. Studies of individuals without diabetes found that home measurements may better correlate with ASCVD risk than office measurements (22,23). Moreover, home blood pressure monitoring may improve medication-taking behavior and thus help reduce cardiovascular risk (24).

Treatment Goals

Recommendations

10.3 For people with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects

of antihypertensive medications, and individual preferences. **B**

10.4 The on-treatment target blood pressure goal is $<130/80$ mmHg, if it can be safely attained. **A**

10.5 In pregnant individuals with diabetes and chronic hypertension, a blood pressure threshold of 140/90 mmHg for initiation or titration of therapy is associated with better pregnancy outcomes than reserving treatment for severe hypertension, with no increase in risk of small-for-gestational-age birth weight. **A** There are limited data on the optimal lower limit, but therapy should be deintensified for blood pressure $<90/60$ mmHg. **E** A blood pressure target of 110–135/85 mmHg is suggested in the interest of reducing the risk for accelerated maternal hypertension. **A**

Randomized clinical trials have demonstrated unequivocally that treatment of hypertension reduces cardiovascular events as well as microvascular complications (25–31). There has been controversy on the recommendation of a specific blood pressure goal in people with diabetes. The committee recognizes that there has been no randomized controlled trial to specifically demonstrate a decreased incidence of cardiovascular events in people with diabetes by targeting a blood pressure $<130/80$ mmHg. The recommendation to support a blood pressure goal of $<130/80$ mmHg in people with diabetes is consistent with guidelines from the American College of Cardiology and American Heart Association (18), the International Society of Hypertension (19), and the European Society of Cardiology (20). The committee’s recommendation for the blood pressure target of $<130/80$ mmHg derives primarily from the collective evidence of the following randomized controlled trials. The Systolic Blood Pressure Intervention Trial (SPRINT) demonstrated that treatment to a target systolic blood pressure of <120 mmHg decreases cardiovascular event rates by 25% in high-risk individuals, although people with diabetes were excluded from this trial (32). The recently completed Strategy of Blood Pressure Intervention in the Elderly Hypertensive Patients (STEP) trial included nearly 20% of people with diabetes and noted decreased cardiovascular events with treatment of hypertension to a blood pressure target of <130 mmHg

(33). While the ACCORD (Action to Control Cardiovascular Risk in Diabetes) blood pressure trial (ACCORD BP) did not confirm that targeting a systolic blood pressure of <120 mmHg in people with diabetes results in decreased cardiovascular event rates, the prespecified secondary outcome of stroke was reduced by 41% with intensive treatment (34). The Action in Diabetes and Vascular Disease: Preterax and Diamicro MR Controlled Evaluation (ADVANCE) trial revealed that treatment with perindopril/indapamide to an achieved systolic blood pressure of ~ 135 mmHg significantly decreased cardiovascular event rates compared with a placebo treatment with an achieved blood pressure of 140 mmHg (35). Therefore, it is recommended that people with diabetes who have hypertension should be treated to blood pressure targets of $<130/80$ mmHg. Notably, there is an absence of high-quality data available to guide blood pressure targets in people with type 1 diabetes, but a similar blood pressure target of $<130/80$ mmHg is recommended in people with type 1 diabetes. As discussed below, treatment should be individualized, and treatment should not be targeted to $<120/80$ mmHg, as a mean achieved blood pressure of $<120/80$ mmHg is associated with adverse events.

Randomized Controlled Trials of Intensive Versus Standard Blood Pressure Control

SPRINT provides the strongest evidence to support lower blood pressure goals in individuals at increased cardiovascular risk, although this trial excluded people with diabetes (32). The trial enrolled 9,361 individuals with a systolic blood pressure of ≥ 130 mmHg and increased cardiovascular risk and treated to a systolic blood pressure target of <120 mmHg (intensive treatment) versus a target of <140 mmHg (standard treatment). The primary composite outcome of MI, coronary syndromes, stroke, heart failure, or death from cardiovascular causes was reduced by 25% in the intensive treatment group. The achieved systolic blood pressures in the trial were 121 mmHg and 136 mmHg in the intensive versus standard treatment group, respectively. Adverse outcomes, including hypotension, syncope, electrolyte abnormality, and acute kidney injury (AKI), were more common in the intensive treatment arm; risk of adverse outcomes needs to be weighed against the cardiovascular benefit of more intensive blood pressure lowering.

ACCORD BP provides the strongest direct assessment of the benefits and risks of intensive blood pressure control in people with type 2 diabetes (34). In the study, a total of 4,733 individuals with type 2 diabetes were assigned to intensive therapy (targeting a systolic blood pressure <120 mmHg) or standard therapy (targeting a systolic blood pressure <140 mmHg). The mean achieved systolic blood pressures were 119 mmHg and 133 mmHg in the intensive versus standard group, respectively. The primary composite outcome of nonfatal MI, nonfatal stroke, or death from cardiovascular causes was not significantly reduced in the intensive treatment group. The prespecified secondary outcome of stroke was significantly reduced by 41% in the intensive treatment group. Adverse events attributed to blood pressure treatment, including hypotension, syncope, bradycardia, hyperkalemia, and elevations in serum creatinine, occurred more frequently in the intensive treatment arm than in the standard therapy arm (**Table 10.1**).

Of note, the ACCORD BP and SPRINT trials targeted a similar systolic blood pressure <120 mmHg, but in contrast to SPRINT, the primary composite cardiovascular end point was nonsignificantly reduced in ACCORD BP. The results have been interpreted to be generally consistent between both trials, but ACCORD BP was viewed as underpowered due to the composite primary end point being less sensitive to blood pressure regulation (32).

The more recent STEP trial assigned 8,511 individuals aged 60–80 years with hypertension to a systolic blood pressure target of 110 to <130 mmHg (intensive treatment) or a target of 130 to <150 mmHg (33). In this trial, the primary composite outcome of stroke, acute coronary syndrome, acute decompensated heart failure, coronary revascularization, atrial fibrillation, or death from cardiovascular causes occurred in 3.5% of individuals in the intensive treatment group versus 4.6% in the standard treatment group (hazard ratio [HR] 0.74 [95% CI 0.60–0.92]; $P = 0.007$). In this trial, 18.9% of individuals in the intensive treatment arm and 19.4% in the standard treatment arm had a diagnosis of type 2 diabetes. Hypotension occurred more frequently in the intensive treatment group (3.4%) compared with the standard treatment group (2.6%), without significant differences in other adverse

events, including dizziness, syncope, or fractures.

In ADVANCE, 11,140 people with type 2 diabetes were randomized to receive either treatment with a fixed combination of perindopril/indapamide or matching placebo (35). The primary end point, a composite of cardiovascular death, nonfatal stroke infarction, or worsening renal or diabetic eye disease, was reduced by 9% in the combination treatment. The achieved systolic blood pressure was ~ 135 mmHg in the treatment group and 140 mmHg in the placebo group.

The Hypertension Optimal Treatment (HOT) trial enrolled 18,790 individuals and targeted diastolic blood pressure <90 mmHg, <85 mmHg, or <80 mmHg (36). The cardiovascular event rates, defined as fatal or nonfatal MI, fatal and nonfatal strokes, and all other cardiovascular events, were not significantly different between diastolic blood pressure targets (≤ 90 mmHg, ≤ 85 mmHg, and ≤ 80 mmHg), although the lowest incidence of cardiovascular events occurred with an achieved diastolic blood pressure of 82 mmHg. However, in people with diabetes, there was a significant 51% reduction in the treatment group with a target diastolic blood pressure of <80 mmHg compared with a target diastolic blood pressure of <90 mmHg.

Meta-analyses of Trials

To clarify optimal blood pressure targets in people with diabetes, multiple meta-analyses have been performed. One of the largest meta-analyses included 73,913 people with diabetes. Compared with a less tight blood pressure control, allocation to a tighter blood pressure control significantly reduced the risk of stroke by 31% but did not reduce the risk of MI (37). Another meta-analysis of 19 trials that included 44,989 individuals showed that a mean blood pressure of 133/76 mmHg is associated with a 14% risk reduction for major cardiovascular events compared with a mean blood pressure of 140/81 mmHg (31). This benefit was greatest in people with diabetes. An analysis of trials including people with type 2 diabetes and impaired glucose tolerance with achieved systolic blood pressures of <135 mmHg in the intensive blood pressure treatment group and <140 mmHg in the standard treatment group revealed a 10% reduction in all-cause mortality and

a 17% reduction in stroke (29). More intensive reduction to <130 mmHg was associated with a further reduction in stroke but not other cardiovascular events.

Several meta-analyses stratified clinical trials by mean baseline blood pressure or mean blood pressure attained in the intervention (or intensive treatment) arm. Based on these analyses, antihypertensive treatment appears to be most beneficial when mean baseline blood pressure is $\geq 140/90$ mmHg (17,25,26,28–30). Among trials with lower baseline or attained blood pressure, antihypertensive treatment reduced the risk of stroke, retinopathy, and albuminuria, but effects on other ASCVD outcomes and heart failure were not evident.

Individualization of Treatment Targets

People with diabetes and clinicians should engage in a shared decision-making process to determine individual blood pressure targets (17). This approach acknowledges that the benefits and risks of intensive blood pressure targets are uncertain and may vary across individuals and is consistent with a person-focused approach to care that values individual priorities and health care professional judgment (38). Secondary analyses of ACCORD BP and SPRINT suggest that clinical factors can help determine individuals more likely to benefit and less likely to be harmed by intensive blood pressure control (39,40).

Absolute benefit from blood pressure reduction correlated with absolute baseline cardiovascular risk in SPRINT and in earlier clinical trials conducted at higher baseline blood pressure levels (40,41). Extrapolation of these studies suggests that people with diabetes may also be more likely to benefit from intensive blood pressure control when they have high absolute cardiovascular risk. This approach is consistent with guidelines from the American College of Cardiology and American Heart Association, which also advocate a blood pressure target of $<130/80$ mmHg for all people, with or without diabetes (18).

Potential adverse effects of antihypertensive therapy (e.g., hypotension, syncope, falls, AKI, and electrolyte abnormalities) should also be taken into account (32,34,42,43). Individuals with older age, CKD, and frailty have been shown to be at higher risk of adverse effects of intensive blood pressure control (42). In addition, individuals with orthostatic hypotension, substantial comorbidity, functional limitations,

Table 10.1—Randomized controlled trials of intensive versus standard hypertension treatment strategies

Clinical trial	Population	Intensive	Standard	Outcomes
ACCORD BP (34)	4,733 participants with T2D aged 40–79 years with prior evidence of CVD or multiple cardiovascular risk factors	SBP target: ≤120 mmHg Achieved (mean) SBP/DBP: 119.3/64.4 mmHg	SBP target: 130–140 mmHg Achieved (mean) SBP/DBP: 135/70.5 mmHg	<ul style="list-style-type: none"> No benefit in primary end point: composite of nonfatal MI, nonfatal stroke, and CVD death Stroke risk reduced 41% with intensive control, not sustained through follow-up beyond the period of active treatment Adverse events more common in intensive group, particularly elevated serum creatinine and electrolyte abnormalities
ADVANCE (35)	11,140 participants with T2D aged ≥55 years with prior evidence of CVD or multiple cardiovascular risk factors	Intervention: a single-pill, fixed-dose combination of perindopril and indapamide Achieved (mean) SBP/DBP: 136/73 mmHg	Control: placebo Achieved (mean) SBP/DBP: 141.6/75.2 mmHg	<ul style="list-style-type: none"> Intervention reduced risk of primary composite end point of major macrovascular and microvascular events (9%), death from any cause (14%), and death from CVD (18%) 6-year observational follow-up found reduction in risk of death in intervention group attenuated but still significant (310)
HOT (36)	18,790 participants, including 1,501 with diabetes	DBP target: ≤80 mmHg Achieved (mean): 81.1 mmHg, ≤80 group; 85.2 mmHg, ≤90 group	DBP target: ≤90 mmHg	<ul style="list-style-type: none"> In the overall trial, there was no cardiovascular benefit with more intensive targets In the subpopulation with diabetes, an intensive DBP target was associated with a significantly reduced risk (51%) of CVD events
SPRINT (42)	9,361 participants without diabetes	SBP target: ≤120 mmHg Achieved (mean): 121.4 mmHg	SBP target: ≤140 mmHg Achieved (mean): 136.2 mmHg	<ul style="list-style-type: none"> Intensive SBP target lowered risk of the primary composite outcome 25% (MI, ACS, stroke, heart failure, and death due to CVD) Intensive target reduced risk of death 27% Intensive therapy increased risks of electrolyte abnormalities and AKI
STEP (33)	8,511 participants aged 60–80 years, including 1,627 with diabetes	SBP target: ≤130 mmHg Achieved (mean): 127.5 mmHg	SBP target: ≤150 mmHg Achieved (mean): 135.3 mmHg	<ul style="list-style-type: none"> Intensive SBP target lowered risk of the primary composite outcome 26% (stroke, ACS [acute MI and hospitalization for unstable angina], acute decompensated heart failure, coronary revascularization, atrial fibrillation, or death from cardiovascular causes) Intensive target reduced risk of cardiovascular death 28% Intensive therapy increased risks of hypotension

ACCORD BP, Action to Control Cardiovascular Risk in Diabetes Blood Pressure trial; ACS, acute coronary syndrome; ADVANCE, Action in Diabetes and Vascular Disease: Preterax and Diamicron MR Controlled Evaluation; AKI, acute kidney injury; CVD, cardiovascular disease; DBP, diastolic blood pressure; HOT, Hypertension Optimal Treatment trial; MI, myocardial infarction; SBP, systolic blood pressure; SPRINT, Systolic Blood Pressure Intervention Trial; STEP, Strategy of Blood Pressure Intervention in the Elderly Hypertensive Patients; T2D, type 2 diabetes.

or polypharmacy may be at high risk of adverse effects, and some individuals may prefer higher blood pressure targets to enhance quality of life. However, ACCORD BP demonstrated that intensive blood pressure lowering decreased the risk of cardiovascular events irrespective of baseline diastolic blood pressure in individuals who also received standard glycemic control

(44). Therefore, the presence of low diastolic blood pressure is not necessarily a contraindication to more intensive blood pressure management in the context of otherwise standard care.

Pregnancy and Antihypertensive Medications
There are few randomized controlled trials of antihypertensive therapy in pregnant

individuals with diabetes. A 2018 Cochrane systematic review of antihypertensive therapy for mild to moderate chronic hypertension included 63 trials and over 5,909 women and suggested that antihypertensive therapy probably reduces the risk of developing severe hypertension but may not affect the risk of fetal or neonatal death, small-for-gestational-age babies, or

preterm birth (45). The Control of Hypertension in Pregnancy Study (CHIPS) (46) enrolled mostly women with chronic hypertension. In CHIPS, targeting a diastolic blood pressure of 85 mmHg during pregnancy was associated with reduced likelihood of developing accelerated maternal hypertension and no demonstrable adverse outcome for infants compared with targeting a higher diastolic blood pressure. The mean systolic blood pressure achieved in the more intensively treated group was 133.1 ± 0.5 mmHg, and the mean diastolic blood pressure achieved in that group was 85.3 ± 0.3 mmHg. A similar approach is supported by the International Society for the Study of Hypertension in Pregnancy, which specifically recommends use of antihypertensive therapy to maintain systolic blood pressure between 110 and 140 mmHg and diastolic blood pressure between 80 and 85 mmHg (47).

The more recent Chronic Hypertension and Pregnancy (CHAP) trial assigned pregnant individuals with mild chronic hypertension to antihypertensive medications to target a blood pressure goal of $<140/90$ mmHg (active treatment group) or to control treatment, in which antihypertensive therapy was withheld unless severe hypertension (systolic pressure ≥ 160 mmHg or diastolic pressure ≥ 105 mmHg) developed (control group) (48). The primary outcome, a composite of preeclampsia with severe features, medically indicated preterm birth at <35 weeks of gestation, placental abruption, or fetal/neonatal death, occurred in 30.2% of female participants in the active treatment group versus 37.0% in the control group ($P < 0.001$). The mean systolic blood pressure between randomization and delivery was 129.5 mmHg in the active treatment group and 132.6 mmHg in the control group.

Current evidence supports controlling blood pressure to 110–135/85 mmHg to reduce the risk of accelerated maternal hypertension but also to minimize impairment of fetal growth. During pregnancy, treatment with ACE inhibitors, angiotensin receptor blockers (ARBs), direct renin inhibitors, and spironolactone are contraindicated, as they may cause fetal damage. Special consideration should be taken for individuals of childbearing potential, and people intending to become pregnant should switch from an ACE inhibitor/ARB or spironolactone to an alternative antihypertensive medication approved during pregnancy. Antihypertensive drugs known

to be effective and safe in pregnancy include methyldopa, labetalol, and long-acting nifedipine, while hydralazine may be considered in the acute management of hypertension in pregnancy or severe pre-eclampsia (49). Diuretics are not recommended for blood pressure control in pregnancy but may be used during late-stage pregnancy if needed for volume control (49,50). The American College of Obstetricians and Gynecologists also recommends that postpartum individuals with gestational hypertension, preeclampsia, and superimposed preeclampsia have their blood pressures observed for 72 h in the hospital and 7–10 days postpartum. Long-term follow-up is recommended for these individuals, as they have increased lifetime cardiovascular risk (51). See Section 15, “Management of Diabetes in Pregnancy,” for additional information.

Treatment Strategies

Lifestyle Intervention

Recommendation

10.6 For people with blood pressure $>120/80$ mmHg, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, smoking cessation, and increased physical activity. **A**

Lifestyle management is an important component of hypertension treatment because it lowers blood pressure, enhances the effectiveness of some antihypertensive medications, promotes other aspects of metabolic and vascular health, and generally leads to few adverse effects. Lifestyle therapy consists of reducing excess body weight through caloric restriction (see Section 8, “Obesity and Weight Management for the Prevention and Treatment of Type 2 Diabetes”), at least 150 min of moderate-intensity aerobic activity per week (see Section 3, “Prevention or Delay of Diabetes and Associated Comorbidities”), restricting sodium intake ($<2,300$ mg/day), increasing consumption of fruits and vegetables (8–10 servings per day) and low-fat dairy products (2–3 servings per day), avoiding excessive alcohol consumption (no more than 2 servings per day in men and no more than 1 serving per day in

women) (52), and increasing activity levels (53) (see Section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes”).

These lifestyle interventions are reasonable for individuals with diabetes and mildly elevated blood pressure (systolic >120 mmHg or diastolic >80 mmHg) and should be initiated along with pharmacologic therapy when hypertension is diagnosed (Fig. 10.2) (53). A lifestyle therapy plan should be developed in collaboration with the person with diabetes and discussed as part of diabetes management. Use of internet or mobile-based digital platforms to reinforce healthy behaviors may be considered as a component of care, as these interventions have been found to enhance the efficacy of medical therapy for hypertension (54,55).

Pharmacologic Interventions

Recommendations

10.7 Individuals with confirmed office-based blood pressure $\geq 130/80$ mmHg qualify for initiation and titration of pharmacologic therapy to achieve the recommended blood pressure goal of $<130/80$ mmHg. **A**

10.8 Individuals with confirmed office-based blood pressure $\geq 150/90$ mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single-pill combination of drugs demonstrated to reduce cardiovascular events in people with diabetes. **A**

10.9 Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in people with diabetes. **A** ACE inhibitors or angiotensin receptor blockers (ARBs) are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease. **A**

10.10 Multiple-drug therapy is generally required to achieve blood pressure targets. However, combinations of ACE inhibitors and ARBs and combinations of ACE inhibitors or ARBs (including ARBs/neprilysin inhibitors) with direct renin inhibitors should not be used. **A**

10.11 An ACE inhibitor or ARB, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in people with diabetes and urinary albumin-to-creatinine ratio ≥ 300 mg/g

Recommendations for the Treatment of Confirmed Hypertension in Nonpregnant People With Diabetes




Figure 10.2—Recommendations for the treatment of confirmed hypertension in nonpregnant people with diabetes. *An ACE inhibitor (ACEi) or angiotensin receptor blocker (ARB) is suggested to treat hypertension for people with coronary artery disease (CAD) or urine albumin-to-creatinine ratio 30–299 mg/g creatinine and strongly recommended for individuals with urine albumin-to-creatinine ratio ≥ 300 mg/g creatinine. †Dihydropyridine calcium channel blocker (CCB). ‡Thiazide-like diuretic; long-acting agents shown to reduce cardiovascular events, such as chlorothalidone and indapamide, are preferred. BP, blood pressure. Adapted from de Boer et al. (18).

creatinine A or 30–299 mg/g creatinine. B If one class is not tolerated, the other should be substituted. B

10.12 For adults treated with an ACE inhibitor, ARB, mineralocorticoid receptor antagonist (MRA), or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels

should be monitored within 7–14 days after initiation of therapy and at least annually. B

Initial Number of Antihypertensive Medications. Initial treatment for people with diabetes depends on the severity of hypertension (Fig. 10.2). Those with blood

pressure between 130/80 mmHg and 150/90 mmHg may begin with a single drug. For individuals with blood pressure $\geq 150/90$ mmHg, initial pharmacologic treatment with two antihypertensive medications is recommended in order to more effectively achieve adequate blood pressure control (56–58). Single-pill antihypertensive combinations may improve

medication taking in some individuals (59).

Classes of Antihypertensive Medications. Initial treatment for hypertension should include any of the drug classes demonstrated to reduce cardiovascular events in people with diabetes: ACE inhibitors (60,61), ARBs (60,61), thiazide-like diuretics (62), or dihydropyridine calcium channel blockers (63). In people with diabetes and established coronary artery disease, ACE inhibitors or ARBs are recommended first-line therapy for hypertension (64–66). For individuals with albuminuria (urine albumin-to-creatinine ratio [UACR] ≥ 30 mg/g), initial treatment should include an ACE inhibitor or ARB to reduce the risk of progressive kidney disease (18) (Fig. 10.2). In individuals receiving ACE inhibitor or ARB therapy, continuation of those medications as kidney function declines to estimated glomerular filtration rate (eGFR) < 30 mL/min/ 1.73 m^2 may provide cardiovascular benefit without significantly increasing the risk of end-stage kidney disease (67). In the absence of albuminuria, risk of progressive kidney disease is low, and ACE inhibitors and ARBs have not been found to afford superior cardioprotection compared with thiazide-like diuretics or dihydropyridine calcium channel blockers (68). β -Blockers are indicated in the setting of prior MI, active angina, or HFrEF but have not been shown to reduce mortality as blood pressure–lowering agents in the absence of these conditions (27,69,70).

Multiple-Drug Therapy. Multiple-drug therapy is often required to achieve blood pressure targets (Fig. 10.2), particularly in the setting of diabetic kidney disease. However, the use of both ACE inhibitors and ARBs in combination, or the combination of an ACE inhibitor or ARB and a direct renin inhibitor, is contraindicated given the lack of added ASCVD benefit and increased rate of adverse events—namely, hyperkalemia, syncope, and AKI (71–73). Titration of and/or addition of further blood pressure medications should be made in a timely fashion to overcome therapeutic inertia in achieving blood pressure targets.

Bedtime Dosing. Although prior analyses of randomized clinical trials found a benefit to evening versus morning dosing of antihypertensive medications (74,75), these results have not been reproduced in subsequent trials. Therefore, preferential use

of antihypertensives at bedtime is not recommended (76).

Hyperkalemia and Acute Kidney Injury. Treatment with ACE inhibitors/ARBs or mineralocorticoid receptor antagonists (MRAs) can cause AKI and hyperkalemia, while diuretics can cause AKI and either hypokalemia or hyperkalemia (depending on mechanism of action) (77,78). Detection and management of these abnormalities is important because AKI and hyperkalemia each increase the risks of cardiovascular events and death (79). Therefore, serum creatinine and potassium should be monitored after initiation of treatment with an ACE inhibitor/ARB, MRA, or diuretic and monitored during treatment and following up titration of these medications, particularly among individuals with reduced glomerular filtration who are at increased risk of hyperkalemia and AKI (77,78,80).

Resistant Hypertension

Recommendation

10.13 Individuals with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for MRA therapy. **A**

Resistant hypertension is defined as blood pressure $\geq 140/90$ mmHg despite a therapeutic strategy that includes appropriate lifestyle management plus a diuretic and two other antihypertensive drugs with complementary mechanisms of action at adequate doses. Prior to diagnosing resistant hypertension, a number of other conditions should be excluded, including missed doses of antihypertensive medications, white coat hypertension, and secondary hypertension. People with diabetes and confirmed resistant hypertension should be evaluated for secondary causes of hypertension, including primary hyperaldosteronism, renal artery stenosis, diabetic kidney disease, and obstructive sleep apnea. In general, barriers to medication taking (such as cost and side effects) should be identified and addressed (Fig. 10.2).

MRAs, including spironolactone and eplerenone, are effective for management of resistant hypertension in people with type 2 diabetes when added to existing treatment with an ACE inhibitor or ARB, thiazide-like diuretic, or dihydropyridine calcium channel blocker (81). In addition,

MRAs reduce albuminuria in people with diabetic nephropathy (82–84). However, adding an MRA to a treatment plan that includes an ACE inhibitor or ARB may increase the risk for hyperkalemia, emphasizing the importance of regular monitoring for serum creatinine and potassium in these individuals, and long-term outcome studies are needed to better evaluate the role of MRAs in blood pressure management.

LIPID MANAGEMENT

Lifestyle Intervention

Recommendations

10.14 Lifestyle modification focusing on weight loss (if indicated); application of a Mediterranean or DASH eating pattern; reduction of saturated fat and *trans* fat; increase of dietary n-3 fatty acids, viscous fiber, and plant stanol/sterol intake; and increased physical activity should be recommended to improve the lipid profile and reduce the risk of developing atherosclerotic cardiovascular disease (ASCVD) in people with diabetes. **A**

10.15 Intensify lifestyle therapy and optimize glycemic control for people with diabetes with elevated triglyceride levels (≥ 150 mg/dL [≥ 1.7 mmol/L]) and/or low HDL cholesterol (< 40 mg/dL [< 1.0 mmol/L] for men and < 50 mg/dL [< 1.3 mmol/L] for women). **C**

Lifestyle intervention, including weight loss in people with overweight or obesity (when appropriate) (85), increased physical activity, and medical nutrition therapy, allows some individuals to reduce ASCVD risk factors. Nutrition intervention should be tailored according to each person's age, pharmacologic treatment, lipid levels, and medical conditions.

Recommendations should focus on application of a Mediterranean (83) or Dietary Approaches to Stop Hypertension (DASH) eating pattern, reducing saturated and *trans* fat intake, and increasing plant stanol/sterol, n-3 fatty acid, and viscous fiber (such as in oats, legumes, and citrus) intake (86,87). Glycemic control may also beneficially modify plasma lipid levels, particularly in people with very high triglycerides and poor glycemic control. See Section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes,” for additional nutrition information.

Ongoing Therapy and Monitoring With Lipid Panel

Recommendations

10.16 In adults with prediabetes or diabetes not taking statins or other lipid-lowering therapy, it is reasonable to obtain a lipid profile at the time of diagnosis, at an initial medical evaluation, annually thereafter, or more frequently if indicated. **E**

10.17 Obtain a lipid profile at initiation of statins or other lipid-lowering therapy, 4–12 weeks after initiation or a change in dose, and annually thereafter, as it may help to monitor the response to therapy and inform medication taking. **A**

In adults with diabetes, it is reasonable to obtain a lipid profile (total cholesterol, LDL cholesterol, HDL cholesterol, and triglycerides) at the time of diagnosis, at the initial medical evaluation, and at least every 5 years thereafter in individuals <40 years of age. In younger people with longer duration of disease (such as those with youth-onset type 1 diabetes), more frequent lipid profiles may be reasonable. A lipid panel should also be obtained immediately before initiating statin therapy. Once an individual is taking a statin, LDL cholesterol levels should be assessed 4–12 weeks after initiation of statin therapy, after any change in dose, and annually (e.g., to monitor for medication taking and efficacy). Monitoring lipid profiles after initiation of statin therapy and during therapy increases dose titration and statin adherence (88–90). If LDL cholesterol levels are not responding in spite of medication taking, clinical judgment is recommended to determine the need for and timing of lipid panels. In individual patients, the highly variable LDL cholesterol-lowering response seen with statins is poorly understood (91). Clinicians should attempt to find a dose or alternative statin that is tolerable if side effects occur. There is evidence for benefit from even extremely low, less than daily statin doses (92).

STATIN TREATMENT

Primary Prevention

Recommendations

10.18 For people with diabetes aged 40–75 years without ASCVD, use moderate-intensity statin therapy in addition to lifestyle therapy. **A**

10.19 For people with diabetes aged 20–39 years with additional ASCVD risk factors, it may be reasonable to initiate statin therapy in addition to lifestyle therapy. **C**

10.20 For people with diabetes aged 40–75 years at higher cardiovascular risk, including those with one or more ASCVD risk factors, it is recommended to use high-intensity statin therapy to reduce LDL cholesterol by $\geq 50\%$ of baseline and to target an LDL cholesterol goal of <70 mg/dL (<1.8 mmol/L). **A**

10.21 For people with diabetes aged 40–75 years at higher cardiovascular risk, especially those with multiple ASCVD risk factors and an LDL cholesterol ≥ 70 mg/dL (≥ 1.8 mmol/L), it may be reasonable to add ezetimibe or a PCSK9 inhibitor to maximum tolerated statin therapy. **B**

10.22 In adults with diabetes aged >75 years already on statin therapy, it is reasonable to continue statin treatment. **B**

10.23 In adults with diabetes aged >75 years, it may be reasonable to initiate moderate-intensity statin therapy after discussion of potential benefits and risks. **C**

10.24 In people with diabetes intolerant to statin therapy, treatment with bempedoic acid is recommended to reduce cardiovascular event rates as an alternative cholesterol-lowering plan. **A**

10.25 Statin therapy is contraindicated in pregnancy. **B**

10.28a For individuals who do not tolerate the intended statin intensity, the maximum tolerated statin dose should be used. **E**

10.28b For people with diabetes and ASCVD intolerant to statin therapy, PCSK9 inhibitor therapy with monoclonal antibody treatment, A bempedoic acid therapy, A or PCSK9 inhibitor therapy with inclisiran siRNA E should be considered as an alternative cholesterol-lowering therapy.

Initiating Statin Therapy Based on Risk

People with type 2 diabetes have an increased prevalence of lipid abnormalities, contributing to their high risk of ASCVD. Multiple clinical trials have demonstrated the beneficial effects of statin therapy on ASCVD outcomes in subjects with and without CHD (93,94). Subgroup analyses of people with diabetes in larger trials (95–99) and trials in people with diabetes (100,101) showed significant primary and secondary prevention of ASCVD events and CHD death in people with diabetes. Meta-analyses including data from over 18,000 people with diabetes from 14 randomized trials of statin therapy (mean follow-up 4.3 years) demonstrated a 9% proportional reduction in all-cause mortality and 13% reduction in vascular mortality for each 1 mmol/L (39 mg/dL) reduction in LDL cholesterol (102). The cardiovascular benefit in this large meta-analysis did not depend on baseline LDL cholesterol levels and was linearly related to the LDL cholesterol reduction without a low threshold beyond which there was no benefit observed (102).

Accordingly, statins are the drugs of choice for LDL cholesterol lowering and cardioprotection. **Table 10.2** shows the two statin dosing intensities that are recommended for use in clinical practice: high-intensity statin therapy will achieve an approximately $\geq 50\%$ reduction in LDL cholesterol, and moderate-intensity statin plans achieve 30–49% reductions in LDL cholesterol. Low-dose statin therapy is generally not recommended in people with diabetes but is sometimes the only dose of statin that an individual can tolerate. For individuals who do not tolerate the intended intensity of statin, the maximum tolerated statin dose should be used.

Table 10.2—High-intensity and moderate-intensity statin therapy

High-intensity statin therapy (lowers LDL cholesterol by $\geq 50\%$)	Moderate-intensity statin therapy (lowers LDL cholesterol by 30–49%)
Atorvastatin 40–80 mg	Atorvastatin 10–20 mg
Rosuvastatin 20–40 mg	Rosuvastatin 5–10 mg
	Simvastatin 20–40 mg
	Pravastatin 40–80 mg
	Lovastatin 40 mg
	Fluvastatin XL 80 mg
	Pitavastatin 1–4 mg

Once-daily dosing. XL, extended release.

As in those without diabetes, absolute reductions in ASCVD outcomes (CHD death and nonfatal MI) are greatest in people with high baseline ASCVD risk (known ASCVD and/or very high LDL cholesterol levels), but the overall benefits of statin therapy in people with diabetes at moderate or even low risk for ASCVD are convincing (103,104). The relative benefit of lipid-lowering therapy has been uniform across most subgroups tested (94,102), including subgroups that varied with respect to age and other risk factors.

Primary Prevention (People Without ASCVD) For primary prevention, moderate-dose statin therapy is recommended for those aged ≥ 40 years (96,103,104), although high-intensity therapy should be considered in the context of additional ASCVD risk factors. The evidence is strong for people with diabetes aged 40–75 years, an age-group well represented in statin trials showing benefit. Since cardiovascular risk is enhanced in people with diabetes, as noted above, individuals who also have multiple other coronary risk factors have increased risk, equivalent to that of those with ASCVD. Therefore, current guidelines recommend that in people with diabetes who are at higher cardiovascular risk, especially those with one or more ASCVD risk factors, high-intensity statin therapy should be prescribed to reduce LDL cholesterol by $\geq 50\%$ from baseline and to target an LDL cholesterol of < 70 mg/dL (< 1.8 mmol/L) (105–107). Since, in clinical practice, it is frequently difficult to ascertain the baseline LDL cholesterol level prior to statin therapy initiation, in those individuals, a focus on an LDL cholesterol target level of < 70 mg/dL (< 1.8 mmol/L) rather than the percent reduction in LDL cholesterol is recommended. In those individuals, it

may also be reasonable to add ezetimibe or proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor therapy to maximum tolerated statin therapy if needed to reduce LDL cholesterol levels by $\geq 50\%$ and to achieve the recommended LDL cholesterol target of < 70 mg/dL (< 1.8 mmol/L) (108). While there are no randomized controlled trials specifically assessing cardiovascular outcomes of adding ezetimibe or PCSK9 inhibitors to statin therapy in primary prevention, the Open-Label Study of Long-term Evaluation Against LDL Cholesterol (OSLER) study included $\sim 80\%$ of study participants without established cardiovascular disease. In the treatment group, LDL cholesterol was reduced by 61%, and, although exploratory and not a primary outcome, cardiovascular events were reduced by 18% in the standard-therapy group to 0.95% in the evolocumab group (HR in the evolocumab group 0.47 [95% CI 0.28–0.78]; $P = 0.003$) (109). Similarly, the Long-term Safety and Tolerability of Alirocumab in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Lipid-Modifying Therapy (ODYSSEY LONG TERM) included $\sim 30\%$ of study participants without established cardiovascular disease and 33% of people with diabetes. Alirocumab added to statin therapy reduced LDL cholesterol by 62% and major adverse cardiovascular events in a post hoc analysis (1.7% vs. 3.3% compared with placebo; HR 0.52 [95% CI 0.31–0.90]; nominal $P = 0.02$) (110). In addition, a meta-analysis suggests that there is a cardiovascular benefit of adding ezetimibe or PCSK9 inhibitors to treatment for high-risk people (111). The evidence is lesser for individuals aged > 75 years; relatively few older people with diabetes have been enrolled in primary prevention trials. However, heterogeneity by age has

not been seen in the relative benefit of lipid-lowering therapy in trials that included older participants (94,101,102), and because older age confers higher risk, the absolute benefits are actually greater (94,112). Moderate-intensity statin therapy is recommended in people with diabetes who are ≥ 75 years of age. However, the risk-benefit profile should be routinely evaluated in this population, with downward titration of dose performed as needed. See Section 13, “Older Adults,” for more details on clinical considerations for this population.

Age < 40 Years and/or Type 1 Diabetes. Very little clinical trial evidence exists for people with type 2 diabetes under the age of 40 years or for people with type 1 diabetes of any age. For pediatric recommendations, see Section 14, “Children and Adolescents.” In the Heart Protection Study (lower age limit 40 years), the subgroup of ~ 600 people with type 1 diabetes had a reduction in risk proportionately similar, although not statistically significant, to that in people with type 2 diabetes (96). Even though the data are not definitive, similar statin treatment approaches should be considered for people with type 1 or type 2 diabetes, particularly in the presence of other cardiovascular risk factors. Individuals < 40 years of age have lower risk of developing a cardiovascular event over a 10-year horizon; however, their lifetime risk of developing cardiovascular disease and suffering an MI, stroke, or cardiovascular death is high. For people who are < 40 years of age and/or have type 1 diabetes with other ASCVD risk factors, it is recommended that the individual and health care professional discuss the relative benefits and risks and consider the use of moderate-intensity statin therapy. Please refer to “Type 1 Diabetes Mellitus and Cardiovascular Disease: A Scientific Statement From the American Heart Association and American Diabetes Association” (113) for additional discussion.

Secondary Prevention (People With ASCVD) Because cardiovascular event rates are increased in people with diabetes and established ASCVD, intensive therapy is indicated and has been shown to be of benefit in multiple large meta-analyses and randomized cardiovascular outcomes trials (94,102,112,114,115). High-intensity statin therapy is recommended for all people with diabetes and ASCVD to target an

LDL cholesterol reduction of $\geq 50\%$ from baseline and an LDL cholesterol goal of $< 55 \text{ mg/dL}$ ($< 1.4 \text{ mmol/L}$). Based on the evidence discussed below, addition of ezetimibe or a PCSK9 inhibitor is recommended if this goal is not achieved on maximum tolerated statin therapy. These recommendations are based on the observation that high-intensity versus moderate-intensity statin therapy reduces cardiovascular event rates in high-risk individuals with established cardiovascular disease in randomized trials (98,114). In addition, the Cholesterol Treatment Trialists' Collaboration, involving 26 statin trials, of which 5 compared high-intensity versus moderate-intensity statins (102), showed a 21% reduction in major cardiovascular events in people with diabetes for every 39 mg/dL (1 mmol/L) of LDL cholesterol lowering, irrespective of baseline LDL cholesterol or individual characteristics (102). However, the best evidence to support lower LDL cholesterol targets in people with diabetes and established cardiovascular disease derives from multiple large randomized trials investigating the benefits of adding nonstatin agents to statin therapy. As discussed in detail below, these include combination treatment with statins and ezetimibe (112,116) or PCSK9 inhibitors (115,117–119). Each trial found a significant benefit in the reduction of ASCVD events that was directly related to the degree of further LDL cholesterol lowering. These large trials included a significant number of participants with diabetes and prespecified analyses on cardiovascular outcomes in people with and without diabetes (116,118,119). The decision to add a nonstatin agent should be made following a discussion between a clinician and a person with diabetes about the net benefit, safety, and cost of combination therapy.

Combination Therapy for LDL Cholesterol Lowering

Statins and Ezetimibe

The IMProved Reduction of Outcomes: Vytorin Efficacy International Trial (IMPROVE-IT) was a randomized controlled trial in 18,144 individuals comparing the addition of ezetimibe to simvastatin therapy versus simvastatin alone (112). Individuals were ≥ 50 years of age, had experienced a recent acute coronary syndrome, and were treated for an average of 6 years. Overall, the addition of ezetimibe led to a 6.4% relative benefit and a 2% absolute

reduction in major adverse cardiovascular events (atherosclerotic cardiovascular events), with the degree of benefit being directly proportional to the change in LDL cholesterol, which was 70 mg/dL (1.8 mmol/L) in the statin group on average and 54 mg/dL (1.4 mmol/L) in the combination group (112). In those with diabetes (27% of participants), the combination of moderate-intensity simvastatin (40 mg) and ezetimibe (10 mg) showed a significant reduction of major adverse cardiovascular events with an absolute risk reduction of 5% (40% vs. 45% cumulative incidence at 7 years) and a relative risk reduction of 14% (HR 0.86 [95% CI 0.78–0.94]) over moderate-intensity simvastatin (40 mg) alone (116).

Statins and PCSK9 Inhibitors

Placebo-controlled trials evaluating the addition of the PCSK9 inhibitors evolocumab and alirocumab to maximum tolerated doses of statin therapy in participants who were at high risk for ASCVD demonstrated an average reduction in LDL cholesterol ranging from 36 to 59%. These agents have been approved as adjunctive therapy for individuals with ASCVD or familial hypercholesterolemia who are receiving maximum tolerated statin therapy but require additional lowering of LDL cholesterol (120,121). No cardiovascular outcome trials have been performed to assess whether PCSK9 inhibitor therapy reduces ASCVD event rates in individuals without established cardiovascular disease (primary prevention).

The effects of PCSK9 inhibition on ASCVD outcomes were investigated in the Further Cardiovascular Outcomes Research With PCSK9 Inhibition in Subjects With Elevated Risk (FOURIER) trial, which enrolled 27,564 individuals with prior ASCVD and an additional high-risk feature who were receiving their maximum tolerated statin therapy (two-thirds were on high-intensity statin) but who still had LDL cholesterol $\geq 70 \text{ mg/dL}$ ($\geq 1.8 \text{ mmol/L}$) or non-HDL cholesterol $\geq 100 \text{ mg/dL}$ ($\geq 2.6 \text{ mmol/L}$) (115). Individuals were randomized to receive subcutaneous injections of evolocumab (either 140 mg every 2 weeks or 420 mg every month based on individual preference) versus placebo. Evolocumab reduced LDL cholesterol by 59% from a median of 92 down to 30 mg/dL in the treatment arm.

During the median follow-up of 2.2 years, the composite outcome of cardiovascular death, MI, stroke, hospitalization for angina, or revascularization occurred in 11.3% vs. 9.8% of the placebo and evolocumab groups, respectively, representing a 15% relative risk reduction ($P < 0.001$). The combined end point of cardiovascular death, MI, or stroke was reduced by 20%, from 7.4 to 5.9% ($P < 0.001$). Evolocumab therapy also significantly reduced all strokes (1.5% vs. 1.9%; HR 0.79 [95% CI 0.66–0.95]; $P = 0.01$) and ischemic stroke (1.2% vs. 1.6%; HR 0.75 [95% CI 0.62–0.92]; $P = 0.005$) in the total population, with findings being consistent in individuals with or without a history of ischemic stroke at baseline (122). Importantly, similar benefits were seen in a prespecified subgroup of people with diabetes, comprising 11,031 individuals (40% of the trial) (119).

In the ODYSSEY OUTCOMES trial (Evaluation of Cardiovascular Outcomes After an Acute Coronary Syndrome During Treatment With Alirocumab), 18,924 individuals (28.8% of whom had diabetes) with recent acute coronary syndrome were randomized to the PCSK9 inhibitor alirocumab or placebo every 2 weeks in addition to maximum tolerated statin therapy, with alirocumab dosing titrated between 75 and 150 mg to achieve LDL cholesterol levels between 25 and 50 mg/dL (117). Over a median follow-up of 2.8 years, a composite primary end point (comprising death from CHD, nonfatal MI, fatal or nonfatal ischemic stroke, or unstable angina requiring hospital admission) occurred in 903 individuals (9.5%) in the alirocumab group and in 1,052 individuals (11.1%) in the placebo group (HR 0.85 [95% CI 0.78–0.93]; $P < 0.001$). Combination therapy with alirocumab plus statin therapy resulted in a greater absolute reduction in the incidence of the primary end point in people with diabetes (2.3% [95% CI 0.4–4.2]) than in those with prediabetes (1.2% [0.0–2.4]) or normoglycemia (1.2% [−0.3 to 2.7]) (118).

In addition to monoclonal antibodies targeting PCSK9, the siRNA inclisiran has been developed and has recently become available in the U.S. In the Inclisiran for Participants With Atherosclerotic Cardiovascular Disease and Elevated Low-density Lipoprotein Cholesterol (ORION-10) and

Inclisiran for Subjects With ASCVD or ASCVD-Risk Equivalents and Elevated Low-density Lipoprotein Cholesterol (ORION-11) trials (123), individuals with established cardiovascular disease or ASCVD risk equivalent were randomized to receive inclisiran or placebo. Inclisiran allows less frequent administration compared with monoclonal antibodies and was administered on day 1, on day 90, and every 6 months in these trials. In the ORION-10 trial, 47.5% of individuals in the inclisiran group and 42.4% in the placebo group had diabetes; in the ORION-11 trial, 36.5% of individuals in the inclisiran group and 33.7% in the placebo group had diabetes. The coprimary end point of placebo-corrected percent change in LDL cholesterol level from baseline to day 510 was 52.3% in the ORION-10 trial and 49.9% in the ORION-11 trial. In an exploratory analysis, the prespecified cardiovascular end point, defined as a cardiovascular basket of nonadjudicated terms, including those classified within cardiac death, and any signs or symptoms of cardiac arrest, nonfatal MI, or stroke, occurred in 7.4% of the inclisiran group and 10.2% of the placebo group in the ORION-10 trial and in 7.8% of the inclisiran group and 10.3% of the placebo group in the ORION-11 trial. A cardiovascular outcome trial using inclisiran in people with established cardiovascular disease is currently ongoing (124).

Intolerance to Statin Therapy

Statin therapy is a hallmark approach to cardiovascular prevention and treatment; however, a subset of individuals experience partial (inability to tolerate sufficient dosage necessary to achieve therapeutic objectives due to adverse effects) or complete (inability to tolerate any dose) intolerance to statin therapy (125). Although the definition of statin intolerance differs between organizations and within clinical study methods, these individuals will require an alternative treatment approach. Initial steps in people intolerant to statins may include switching to a different high-intensity statin if a high-intensity statin is indicated, switching to moderate-intensity or low-intensity statin, lowering the statin dose, or using nondaily dosing of statins. While considering these alternative treatment plans, the addition of nonstatin treatment plans to maximum tolerated statin therapy should be considered, as these are frequently associated with

improved adherence and target LDL cholesterol goal achievement (125).

PCSK9 Inhibition

The PCSK9 monoclonal antibodies alirocumab and evolocumab have both been studied within populations considered statin intolerant. The Study of Alirocumab in Patients With Primary Hypercholesterolemia and Moderate, High, or Very High Cardiovascular Risk, Who Are Intolerant to Statins (ODYSSEY ALTERNATIVE) trial evaluated the LDL cholesterol–lowering efficacy of alirocumab compared with ezetimibe in addition to the safety of each of the prior two treatments compared with a statin rechallenge arm with 20 mg atorvastatin in 314 individuals with primary hypercholesterolemia and statin intolerance. The proportion of the study population with type 2 diabetes was ~24%. After the 24 weeks, alirocumab lowered LDL cholesterol levels by 54.8% compared with 20.1% with ezetimibe. Although there were similar rates of any adverse event for all treatments, there were fewer events that led to treatment discontinuation for alirocumab (18.3% vs. 25.0% for ezetimibe and 25.4% for atorvastatin) as well as fewer skeletal muscle–related adverse events (32.5% vs. 41.1% with ezetimibe and 46% with atorvastatin) (126). Individuals in all treatment arms were offered the opportunity of an open-label extension phase, in which all received alirocumab for ~3 years. LDL cholesterol reductions of more than 50% were either achieved or maintained for the 281 individuals who either continued with or switched to alirocumab for the extension phase, and these reductions were sustained throughout the treatment period (127).

Evolocumab was evaluated for its safety and efficacy in people with statin intolerance in the Goal Achievement After Utilizing an Anti-PCSK9 Antibody in Statin Intolerant Subjects 1, 2, and 3 (GAUSS 1, 2, and 3) trials as well as the OSLER open-label extension of the GAUSS 1 and 2 trials. There were 160 and 307 individuals in the GAUSS 1 and 2 trials, respectively, who were randomized to various doses of evolocumab plus ezetimibe 10 mg or ezetimibe 10 mg plus placebo injection for 12 weeks. Reductions in LDL cholesterol ranged from 41% to 63% (depending on the dose) for evolocumab/ezetimibe compared with 15% to 18% for

ezetimibe/placebo. Similar to what was found in the alirocumab studies, musculoskeletal adverse effects occurred in fewer of those treated with evolocumab/ezetimibe than with ezetimibe/placebo, although rates of discontinuation were similar (5% and 6%, respectively) due to these effects. Use of low-dose statins was allowed in these studies and was associated with an increase in the incidence of musculoskeletal adverse effects (128,129). One hundred twenty-eight individuals from the GAUSS 1 trial were rerandomized to evolocumab (420 mg monthly) plus standard of care compared with the standard of care for 1 year, and then all participants were treated with 420 mg of evolocumab monthly plus standard of care. Two hundred fifty-four individuals were rerandomized into two dosing options of evolocumab (140 mg biweekly or 420 mg monthly) compared with standard of care for 1 year, and then all were continued on 420 mg monthly for an additional year. After 1 year, the LDL cholesterol was reduced from baseline (beginning of the GAUSS 1 or 2 trial) by a mean of 57% in those treated with evolocumab compared with 13% with the standard of care, with a 59% reduction (from baseline) by the end of year 2. Fourteen percent of participants in the extension trials experienced musculoskeletal adverse effects; however, these effects did not lead to any participant discontinuing the trials (130). Similar LDL cholesterol reductions were demonstrated in the GAUSS 3 trial after 24 weeks (~54.5% with evolocumab compared with ~16.7% with ezetimibe), with slightly higher rates of musculoskeletal adverse events (20.7% with evolocumab and 28.8% with ezetimibe). The higher rates of these adverse events may be due in part to the first phase of this trial, which randomized individuals to a statin rechallenge with either atorvastatin or placebo (131).

Inclisiran has also been proposed as an option for individuals with statin intolerance. Although most individuals (90–95%) in the later ORION-10 and ORION-11 trials were on statin therapy (123), the Trial to Evaluate the Effect of ALN-PCSSC Treatment on Low-density Lipoprotein Cholesterol (ORION-1) included individuals with documented statin intolerance. The percentages of individuals not taking statins were 26% of the 253 who received either a single injection of inclisiran or placebo and 28% of the 248 who

received two injections of inclisiran or placebo. Both groups were followed for 1 year to assess the durability of the initial (30–45%, dose-dependent) lowering of LDL cholesterol. Almost all individuals treated with inclisiran maintained their LDL cholesterol levels at 180 days; however, the levels returned to within 20% change from the baseline for 17–52% of individuals, and response depended on both the number and strength of the inclisiran dose(s) received (132). A proportion of these individuals continued into the open-label Extension Trial of Inclisiran in Participants With Cardiovascular Disease and High Cholesterol (ORION-3), in which they returned to inclisiran 300 mg given every 6 months and were compared with individuals given placebo in ORION-1 and those who were given evolocumab 140 mg every 2 weeks for 1 year and then transitioned to inclisiran. The change in LDL cholesterol levels was compared with the baseline, which was defined as the baseline for the ORION-1 trial for those initially treated with inclisiran (as they were inclisarin naïve at that point) or the start of the ORION-3 trial for those previously treated with placebo. It is important to note that of the ORION-3 participants, only 23% had diabetes and 33% were not taking statin therapy. Both arms maintained an LDL cholesterol reduction of ~45% through the end of year 4 (133). The significant response was seen across the groups during the ORION-1 trial and in the ORION-3 extension, and it may be expected that those with statin intolerance experienced a response similar to the response of those on statin therapy; however, evaluation of response based on background lipid-lowering therapy was not described.

Bempedoic Acid

Bempedoic acid is a novel LDL cholesterol–lowering agent that is indicated as an adjunct to diet and maximum tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or established ASCVD who require additional lowering of LDL cholesterol. A pooled analysis suggests that bempedoic acid therapy lowers LDL cholesterol levels by about 23% compared with placebo (134). This agent should be considered for individuals who cannot use or tolerate other evidence-based LDL cholesterol–lowering approaches or for whom those other

therapies are inadequately effective (135). The Evaluation of Major Cardiovascular Events in Patients With, or at High Risk for, Cardiovascular Disease Who Are Statin Intolerant Treated With Bempedoic Acid or Placebo (CLEAR Outcomes) trial evaluated the impact of bempedoic acid on cardiovascular events for individuals with established ASCVD (70% of population) or at high risk for ASCVD (30% of population) and considered to be intolerant to statin therapy. It is important to note that ~19% of individuals were on very-low-dose statin therapy at baseline. Bempedoic acid was found to reduce the composite outcome of four-point major adverse cardiovascular events by 13% compared with placebo (136). The HR for the primary outcome was more reduced in the primary prevention group (HR 0.68 [95% CI 0.53–0.87]) compared with the secondary prevention group of individuals with established cardiovascular disease (HR 0.91 [95% CI 0.81–1.01]). In addition, in a preplanned subanalysis of the primary prevention population, the use of bempedoic acid resulted in a 30% reduction in primary composite outcome compared with placebo (137).

Treatment of Other Lipoprotein Fractions or Targets

Recommendations

10.29 For individuals with fasting triglyceride levels ≥ 500 mg/dL (≥ 5.7 mmol/L), evaluate for secondary causes of hypertriglyceridemia and consider medical therapy to reduce the risk of pancreatitis. **C**

10.30 In adults with moderate hypertriglyceridemia (fasting or nonfasting triglycerides 175–499 mg/dL [2.0–5.6 mmol/L]), clinicians should address and treat lifestyle factors (obesity and metabolic syndrome), secondary factors (diabetes, chronic liver or kidney disease and/or nephrotic syndrome, and hypothyroidism), and medications that raise triglycerides. **C**

10.31 In individuals with ASCVD or other cardiovascular risk factors on a statin with controlled LDL cholesterol but elevated triglycerides (135–499 mg/dL [1.5–5.6 mmol/L]), the addition of icosapent ethyl can be considered to reduce cardiovascular risk. **A**

Hypertriglyceridemia should be addressed with dietary and lifestyle changes including

weight loss and abstinence from alcohol (138). Severe hypertriglyceridemia (fasting triglycerides ≥ 500 mg/dL and especially $> 1,000$ mg/dL) may warrant pharmacologic therapy (fibrin acid derivatives and/or fish oil) and reduction in dietary fat to reduce the risk of acute pancreatitis. Moderate- or high-intensity statin therapy should also be used as indicated to reduce risk of cardiovascular events (see STATIN TREATMENT). In people with moderate hypertriglyceridemia, lifestyle interventions, treatment of secondary factors, and avoidance of medications that might raise triglycerides are recommended.

The Reduction of Cardiovascular Events with Icosapent Ethyl-Intervention Trial (REDUCE-IT) enrolled 8,179 adults receiving statin therapy with moderately elevated triglycerides (135–499 mg/dL, median baseline of 216 mg/dL) who had either established cardiovascular disease (secondary prevention cohort) or diabetes plus at least one other cardiovascular risk factor (primary prevention cohort) (139). Individuals were randomized to icosapent ethyl 4 g/day (2 g twice daily with food) versus placebo. The trial met its primary end point, demonstrating a 25% relative risk reduction ($P < 0.001$) for the primary end point composite of cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, or unstable angina. This risk reduction was seen in people with or without diabetes at baseline. The composite of cardiovascular death, nonfatal MI, or nonfatal stroke was reduced by 26% ($P < 0.001$). Additional ischemic end points were significantly lower in the icosapent ethyl group than in the placebo group, including cardiovascular death, which was reduced by 20% ($P = 0.03$). The proportions of individuals experiencing adverse events and serious adverse events were similar between the active and placebo treatment groups. It should be noted that data are lacking for other n-3 fatty acids, and results of the REDUCE-IT trial should not be extrapolated to other products (139). As an example, the addition of 4 g per day of a carboxylic acid formulation of the n-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) (n-3 carboxylic acid) to statin therapy in individuals with atherosgenic dyslipidemia and high cardiovascular risk, 70% of whom had diabetes, did not reduce the risk of major adverse

cardiovascular events compared with the inert comparator of corn oil (140).

Low levels of HDL cholesterol, often associated with elevated triglyceride levels, are the most prevalent pattern of dyslipidemia in people with type 2 diabetes. However, the evidence for the use of drugs that target these lipid fractions is substantially less robust than that for statin therapy (141). In a large trial in people with diabetes, fenofibrate failed to reduce overall cardiovascular outcomes (142).

Other Combination Therapy

Recommendations

10.32 Statin plus fibrate combination therapy has not been shown to improve ASCVD outcomes and is generally not recommended. **A**

10.33 Statin plus niacin combination therapy has not been shown to provide additional cardiovascular benefit above statin therapy alone, may increase the risk of stroke with additional side effects, and is generally not recommended. **A**

Statin and Fibrate Combination Therapy

Combination therapy (statin and fibrate) is associated with an increased risk for abnormal transaminase levels, myositis, and rhabdomyolysis. The risk of rhabdomyolysis is more common with higher doses of statins and renal insufficiency and appears to be higher when statins are combined with gemfibrozil (compared with fenofibrate) (143).

In the ACCORD study, in people with type 2 diabetes who were at high risk for ASCVD, the combination of fenofibrate and simvastatin did not reduce the rate of fatal cardiovascular events, nonfatal MI, or nonfatal stroke compared with simvastatin alone. Prespecified subgroup analyses suggested heterogeneity in treatment effects with possible benefit for men with both a triglyceride level ≥ 204 mg/dL (≥ 2.3 mmol/L) and an HDL cholesterol level ≤ 34 mg/dL (≤ 0.9 mmol/L) (144).

Statin and Niacin Combination Therapy

The Atherothrombosis Intervention in Metabolic Syndrome With Low HDL/High Triglycerides: Impact on Global Health Outcomes (AIM-HIGH) trial randomized over 3,000 people (about one-third with diabetes) with established

ASCVD, LDL cholesterol levels <180 mg/dL (<4.7 mmol/L), low HDL cholesterol levels (men <40 mg/dL [<1.0 mmol/L] and women <50 mg/dL [<1.3 mmol/L]), and triglyceride levels of 150–400 mg/dL (1.7–4.5 mmol/L) to statin therapy plus extended-release niacin or placebo. The trial was halted early due to lack of efficacy on the primary ASCVD outcome (first event of the composite of death from CHD, nonfatal MI, ischemic stroke, hospitalization for an acute coronary syndrome, or symptom-driven coronary or cerebral revascularization) and a possible increase in ischemic stroke in those on combination therapy (145).

The much larger Heart Protection Study 2—Treatment of HDL to Reduce the Incidence of Vascular Events (HPS2-THRIVE) trial also failed to show a benefit of adding niacin to background statin therapy (146). A total of 25,673 individuals with prior vascular disease were randomized to receive 2 g of extended-release niacin and 40 mg of laropiprant (an antagonist of the prostaglandin D2 receptor DP1 that has been shown to improve participation in niacin therapy) versus a matching placebo daily and followed for a median follow-up period of 3.9 years. There was no significant difference in the rate of coronary death, MI, stroke, or coronary revascularization with the addition of niacin–laropiprant versus placebo (13.2% vs. 13.7%; rate ratio 0.96; $P = 0.29$). Niacin–laropiprant was associated with an increased incidence of new-onset diabetes (absolute excess, 1.3 percentage points; $P < 0.001$) and disturbances in diabetes management among those with diabetes. In addition, there was an increase in serious adverse events associated with the gastrointestinal system, musculoskeletal system, skin, and, unexpectedly, infection and bleeding.

Therefore, combination therapy with a statin and niacin is not recommended, given the lack of efficacy on major ASCVD outcomes and increased side effects.

Diabetes Risk With Statin Use

Several studies have reported a modestly increased risk of incident type 2 diabetes with statin use (147,148), which may be limited to those with diabetes risk factors. An analysis of one of the initial studies suggested that although statin use was associated with diabetes risk, the cardiovascular event rate reduction with statins far outweighed

the risk of incident diabetes, even for individuals at highest risk for diabetes (149). The absolute risk increase was small (over 5 years of follow-up, 1.2% of participants on placebo developed diabetes and 1.5% on rosuvastatin developed diabetes) (149). A meta-analysis of 13 randomized statin trials with 91,140 participants showed an odds ratio of 1.09 for a new diagnosis of diabetes, so that (on average) treatment of 255 individuals with statins for 4 years resulted in one additional case of diabetes while simultaneously preventing 5.4 vascular events among those 255 individuals (148).

Lipid-Lowering Agents and Cognitive Function

Although concerns regarding a potential adverse impact of lipid-lowering agents on cognitive function have been raised, several lines of evidence point against this association, as detailed in a 2018 European Atherosclerosis Society Consensus Panel statement (150). First, there are three large randomized trials of statin versus placebo where specific cognitive tests were performed, and no differences were seen between statin and placebo (151–154). In addition, no change in cognitive function has been reported in studies with the addition of ezetimibe (112) or PCSK9 inhibitors (115,155) to statin therapy, including among individuals treated to very low LDL cholesterol levels. In addition, the most recent systematic review of the U.S. Food and Drug Administration's (FDA's) postmarketing surveillance databases, randomized controlled trials, and cohort, case-control, and cross-sectional studies evaluating cognition in individuals receiving statins found that published data do not reveal an adverse effect of statins on cognition (156). Therefore, a concern that statins or other lipid-lowering agents might cause cognitive dysfunction or dementia is not currently supported by evidence and should not deter their use in individuals with diabetes at high risk for ASCVD (156).

ANTIPLATELET AGENTS

Recommendations

10.34 Use aspirin therapy (75–162 mg/day) as a secondary prevention strategy in those with diabetes and a history of ASCVD. **A**

10.35a For individuals with ASCVD and documented aspirin allergy, clopidogrel (75 mg/day) should be used. **B**

10.35b The length of treatment with dual antiplatelet therapy using low-dose aspirin and a P2Y12 inhibitor in individuals with diabetes after an acute coronary syndrome or acute ischemic stroke/transient ischemic attack should be determined by an interprofessional team approach that includes a cardiovascular or neurological specialist, respectively. **E**

10.36 Combination therapy with aspirin plus low-dose rivaroxaban should be considered for individuals with stable coronary and/or peripheral artery disease (PAD) and low bleeding risk to prevent major adverse limb and cardiovascular events. **A**

10.37 Aspirin therapy (75–162 mg/day) may be considered as a primary prevention strategy in those with diabetes who are at increased cardiovascular risk, after a comprehensive discussion with the individual on the benefits versus the comparable increased risk of bleeding. **A**

Risk Reduction

Aspirin has been shown to be effective in reducing cardiovascular morbidity and mortality in high-risk individuals with previous MI or stroke (secondary prevention) and is strongly recommended. In primary prevention, however, among individuals with no previous cardiovascular events, its net benefit is more controversial (147,157).

Previous randomized controlled trials of aspirin, specifically in people with diabetes, failed to consistently show a significant reduction in overall ASCVD end points, raising questions about the efficacy of aspirin for primary prevention in people with diabetes, although some sex differences were suggested (158–160).

The Antithrombotic Trialists' Collaboration published an individual patient-level meta-analysis (161) of the six large trials of aspirin for primary prevention in the general population. These trials collectively enrolled over 95,000 participants, including almost 4,000 with diabetes. Overall, they found that aspirin reduced the risk of serious vascular events by 12% (relative risk 0.88 [95% CI 0.82–0.94]). The largest reduction

was for nonfatal MI, with little effect on CHD death (relative risk 0.95 [95% CI 0.78–1.15]) or total stroke.

Most recently, the ASCEND (A Study of Cardiovascular Events iN Diabetes) trial randomized 15,480 people with diabetes but no evident cardiovascular disease to aspirin 100 mg daily or placebo (162). The primary efficacy end point was vascular death, MI, stroke, or transient ischemic attack. The primary safety outcome was major bleeding (i.e., intracranial hemorrhage, sight-threatening bleeding in the eye, gastrointestinal bleeding, or other serious bleeding). During a mean follow-up of 7.4 years, there was a significant 12% reduction in the primary efficacy end point (8.5% vs. 9.6%; $P = 0.01$). In contrast, major bleeding was significantly increased from 3.2 to 4.1% in the aspirin group (rate ratio 1.29; $P = 0.003$), with most of the excess being gastrointestinal bleeding and other extracranial bleeding. There were no significant differences by sex, weight, or duration of diabetes or other baseline factors, including ASCVD risk score.

Two other large, randomized trials of aspirin for primary prevention, in people without diabetes (ARRIVE [Aspirin to Reduce Risk of Initial Vascular Events]) (163) and in the elderly (ASPREE [Aspirin in Reducing Events in the Elderly]) (164), which included 11% with diabetes, found no benefit of aspirin on the primary efficacy end point and an increased risk of bleeding. In ARRIVE, with 12,546 individuals over a period of 60 months of follow-up, the primary end point occurred in 4.29% vs. 4.48% of individuals in the aspirin versus placebo groups (HR 0.96 [95% CI 0.81–1.13]; $P = 0.60$). Gastrointestinal bleeding events (characterized as mild) occurred in 0.97% of individuals in the aspirin group vs. 0.46% in the placebo group (HR 2.11 [95% CI 1.36–3.28]; $P = 0.0007$). In ASPREE, including 19,114 individuals, for cardiovascular disease (fatal CHD, MI, stroke, or hospitalization for heart failure) after a median of 4.7 years of follow-up, the rates per 1,000 person-years were 10.7 vs. 11.3 events in aspirin vs. placebo groups (HR 0.95 [95% CI 0.83–1.08]). The rate of major hemorrhage per 1,000 person-years was 8.6 events vs. 6.2 events, respectively (HR 1.38 [95% CI 1.18–1.62]; $P < 0.001$).

Thus, aspirin appears to have a modest effect on ischemic vascular events,

with the absolute decrease in events depending on the underlying ASCVD risk. The main adverse effect is an increased risk of gastrointestinal bleeding. The excess risk may be as high as 5 per 1,000 per year in real-world settings. However, for adults with ASCVD risk $>1\%$ per year, the number of ASCVD events prevented will be similar to the number of episodes of bleeding induced, although these complications do not have equal effects on long-term health (165).

Recommendations for using aspirin as primary prevention include both men and women aged ≥ 50 years with diabetes and at least one additional major risk factor (family history of premature ASCVD, hypertension, dyslipidemia, smoking, or CKD/albuminuria) who are not at increased risk of bleeding (e.g., older age, anemia, or renal disease) (166–169). Non-invasive imaging techniques such as coronary calcium scoring may help further tailor aspirin therapy, particularly in those at low risk (170,171). For people >70 years of age (with or without diabetes), the balance appears to have greater risk than benefit (162,164). Thus, for primary prevention, the use of aspirin needs to be carefully considered and may generally not be recommended. Aspirin may be considered in the context of high cardiovascular risk with low bleeding risk but generally not in older adults. Aspirin therapy for primary prevention may be considered in the context of shared decision-making, which carefully weighs the cardiovascular benefits with the fairly comparable increase in risk of bleeding.

For people with documented ASCVD, use of aspirin for secondary prevention has far greater benefit than risk; for this indication, aspirin is still recommended (157).

Aspirin Use in People <50 Years of Age

Aspirin is not recommended for those at low risk of ASCVD (such as men and women aged <50 years with diabetes with no other major ASCVD risk factors), as the low benefit is likely to be outweighed by the risk of bleeding. Clinical judgment should be used for those at intermediate risk (younger individuals with one or more risk factors or older individuals with no risk factors) until further research is available. Individuals' willingness to undergo long-term aspirin therapy should also be considered in shared decision-making (172).

Aspirin use in individuals aged <21 years is generally contraindicated due to the associated risk of Reye syndrome.

Aspirin Dosing

Average daily dosages used in most clinical trials involving people with diabetes ranged from 50 to 650 mg but were mostly in the range of 100–325 mg/day. There is little evidence to support any specific dose, but using the lowest possible dose may help to reduce side effects (173). In the ADAPTABLE (Aspirin Dosing: A Patient-Centric Trial Assessing Benefits and Long-term Effectiveness) trial of individuals with established cardiovascular disease, 38% of whom had diabetes, there were no significant differences in cardiovascular events or major bleeding between individuals assigned to 81 mg and those assigned to 325 mg of aspirin daily (174). In the U.S., the most common low-dose tablet is 81 mg. Although platelets from people with diabetes have altered function, it is unclear what, if any, effect that finding has on the required dose of aspirin for cardioprotective effects in people with diabetes. Many alternate pathways for platelet activation exist that are independent of thromboxane A₂ and thus are not sensitive to the effects of aspirin (175). “Aspirin resistance” has been described in people with diabetes when measured by a variety of ex vivo and in vitro methods (platelet aggregometry and measurement of thromboxane B₂) (176), but other studies suggest no impairment in aspirin response among people with diabetes (177). A trial suggested that more frequent dosing of aspirin may reduce platelet reactivity in individuals with diabetes (178); however, these observations alone are insufficient to empirically recommend that higher doses of aspirin be used in this group at this time. Another meta-analysis raised the hypothesis that low-dose aspirin efficacy is reduced in those weighing >70 kg (179); however, the ASCEND trial found benefit of low-dose aspirin in those in this weight range, which would thus not validate this suggested hypothesis (162). It appears that 75–162 mg/day is optimal.

Indications for P2Y12 Receptor Antagonist Use

Combination dual antiplatelet therapy with aspirin and a P2Y12 receptor antagonist is indicated after acute coronary

syndromes and coronary revascularization with stenting (180). In addition, current guidelines recommend short-term dual antiplatelet therapy after high-risk transient ischemic attack and minor stroke (181). The indications for dual antiplatelet therapy and length of treatment are rapidly evolving and should be determined by an interprofessional team approach that includes a cardiovascular or neurological specialist, respectively. Evidence supports use of either ticagrelor or clopidogrel if no percutaneous coronary intervention was performed and clopidogrel, ticagrelor, or prasugrel if a percutaneous coronary intervention was performed (182). In people with diabetes and prior MI (1–3 years before), adding ticagrelor to aspirin significantly reduces the risk of recurrent ischemic events, including cardiovascular and CHD death (183). Similarly, the addition of ticagrelor to aspirin reduced the risk of ischemic cardiovascular events compared with aspirin alone in people with diabetes and stable coronary artery disease (184,185). However, a higher incidence of major bleeding, including intracranial hemorrhage, was noted with dual antiplatelet therapy. The net clinical benefit (ischemic benefit vs. bleeding risk) was improved with ticagrelor therapy in the large prespecified subgroup of individuals with history of percutaneous coronary intervention, while no net benefit was seen in individuals without prior percutaneous coronary intervention (185). However, early aspirin discontinuation compared with continued dual antiplatelet therapy after coronary stenting may reduce the risk of bleeding without a corresponding increase in the risks of mortality and ischemic events, as shown in a prespecified analysis of people with diabetes enrolled in the TWILIGHT (Ticagrelor With Aspirin or Alone in High-Risk Patients After Coronary Intervention) trial and a recent meta-analysis (186,187).

Combination Antiplatelet and Anticoagulation Therapy

Combination therapy with aspirin plus low-dose rivaroxaban may be considered for people with stable coronary and/or PAD to prevent major adverse limb and cardiovascular complications. In the COMPASS (Cardiovascular Outcomes for People Using Anticoagulation Strategies) trial of 27,395 individuals with established coronary artery disease and/or PAD, aspirin plus rivaroxaban

2.5 mg twice daily was superior to aspirin plus placebo in the reduction of cardiovascular ischemic events, including major adverse limb events. The absolute benefits of combination therapy appeared larger in people with diabetes, who comprised 10,341 of the trial participants (188,189). A similar treatment strategy was evaluated in the Vascular Outcomes Study of ASA (acetylsalicylic acid) Along with Rivaroxaban in Endovascular or Surgical Limb Revascularization for Peripheral Artery Disease (VOYAGER PAD) trial (190), in which 6,564 individuals with PAD who had undergone revascularization were randomly assigned to receive rivaroxaban 2.5 mg twice daily plus aspirin or placebo plus aspirin. Rivaroxaban treatment in this group of individuals was also associated with a significantly lower incidence of ischemic cardiovascular events, including major adverse limb events. However, an increased risk of major bleeding was noted with rivaroxaban added to aspirin treatment in both COMPASS and VOYAGER PAD.

The risks and benefits of dual antiplatelet or antiplatelet plus anticoagulant treatment strategies should be thoroughly discussed with eligible individuals, and shared decision-making should be used to determine an individually appropriate treatment approach. This field of cardiovascular risk reduction is evolving rapidly, as are the definitions of optimal care for individuals with differing types and circumstances of cardiovascular complications.

CARDIOVASCULAR DISEASE

Screening

Recommendations

10.38a In asymptomatic individuals, routine screening for coronary artery disease is not recommended, as it does not improve outcomes as long as ASCVD risk factors are treated. **A**

10.38b Consider investigations for coronary artery disease in the presence of any of the following: atypical cardiac symptoms; signs or symptoms of associated vascular disease, including carotid bruits, transient ischemic attack, stroke, claudication, or PAD; or electrocardiogram abnormalities (e.g., Q waves). **E**

10.39a Adults with diabetes are at increased risk for the development

of asymptomatic cardiac structural or functional abnormalities (stage B heart failure) or symptomatic (stage C) heart failure. Consider screening adults with diabetes by measuring a natriuretic peptide (B-type natriuretic peptide [BNP] or N-terminal pro-BNP [NT-proBNP]) to facilitate prevention of stage C heart failure. **B**

10.39b In asymptomatic individuals with diabetes and abnormal natriuretic peptide levels, echocardiography is recommended to identify stage B heart failure. **A**

10.40 In asymptomatic individuals with diabetes and age ≥ 50 years, microvascular disease in any location, or foot complications or any end-organ damage from diabetes, screening for PAD with ankle-brachial index testing is recommended to guide treatment for cardiovascular disease prevention and limb preservation. **A** In individuals with diabetes duration ≥ 10 years, screening for PAD should be considered. **B**

Treatment

Recommendations

10.41 Among people with type 2 diabetes who have established ASCVD or established kidney disease, a sodium–glucose cotransporter 2 (SGLT2) inhibitor or glucagon-like peptide 1 (GLP-1) receptor agonist with demonstrated cardiovascular disease benefit (**Table 10.3B** and **Table 10.3C**) is recommended as part of the comprehensive cardiovascular risk reduction and/or glucose-lowering treatment plans. **A**

10.41a In people with type 2 diabetes and established ASCVD, multiple ASCVD risk factors, or diabetic kidney disease, an SGLT2 inhibitor with demonstrated cardiovascular benefit is recommended to reduce the risk of major adverse cardiovascular events and/or heart failure hospitalization. **A**

10.41b In people with type 2 diabetes and established ASCVD or multiple risk factors for ASCVD, a GLP-1 receptor agonist with demonstrated cardiovascular benefit is recommended to reduce the risk of major adverse cardiovascular events. **A**

10.41c In people with type 2 diabetes and established ASCVD or multiple risk factors for ASCVD, combined therapy with an SGLT2 inhibitor with demonstrated cardiovascular benefit and a GLP-1 receptor agonist with demonstrated cardiovascular benefit may be considered for additive reduction of the risk of adverse cardiovascular and kidney events. **A**

10.42a In people with type 2 diabetes and established heart failure with either preserved or reduced ejection fraction, an SGLT2 inhibitor (including SGLT1/2 inhibitor) with proven benefit in this patient population is recommended to reduce the risk of worsening heart failure and cardiovascular death. **A**

10.42b In people with type 2 diabetes and established heart failure with either preserved or reduced ejection fraction, an SGLT2 inhibitor with proven benefit in this patient population is recommended to improve symptoms, physical limitations, and quality of life. **A**

10.43 For individuals with type 2 diabetes and chronic kidney disease with albuminuria treated with maximum tolerated doses of ACE inhibitor or ARB, addition of finerenone is recommended to improve cardiovascular outcomes and reduce the risk of chronic kidney disease progression. **A**

10.44 In individuals with diabetes with established ASCVD or aged ≥ 55 years with additional cardiovascular risk factors, ACE inhibitor or ARB therapy is recommended to reduce the risk of cardiovascular events and mortality. **A**

10.45a In individuals with diabetes and asymptomatic stage B heart failure, an interprofessional approach to optimize guideline-directed medical therapy, which should include a cardiovascular disease specialist, is recommended to reduce the risk for progression to symptomatic (stage C) heart failure. **A**

10.45b In individuals with diabetes and asymptomatic stage B heart failure, ACE inhibitors/ARBs and β -blockers are recommended to reduce the risk for progression to symptomatic (stage C) heart failure. **A**

10.45c In individuals with type 2 diabetes and asymptomatic stage B heart failure or with high risk of or

established cardiovascular disease, treatment with an SGLT inhibitor (including SGLT2 or SGLT1/2 inhibitors) is recommended to reduce the risk of hospitalization for heart failure. **A**

10.45d In individuals with type 2 diabetes and diabetic kidney disease, finerenone is recommended to reduce the risk of hospitalization for heart failure. **A**

10.45e In individuals with diabetes, guideline-directed medical therapy for myocardial infarction and symptomatic stage C heart failure is recommended with ACE inhibitors/ARBs, MRAs, angiotensin receptor/neprilysin inhibitor, β -blockers, and SGLT2 inhibitors, similar to guideline-directed medical therapy for people without diabetes. **A**

10.46 In people with type 2 diabetes with stable heart failure, metformin may be continued for glucose lowering if estimated glomerular filtration rate remains >30 mL/min/ 1.73 m^2 but should be avoided in unstable or hospitalized individuals with heart failure. **B**

10.47 Individuals with type 1 diabetes and those with type 2 diabetes who are ketosis prone and/or those consuming ketogenic diets who are treated with SGLT inhibition should be educated on the risks and signs of ketoacidosis and methods of risk management and provided with appropriate tools for accurate ketone measurement (i.e., serum β -hydroxybutyrate). **E**

Cardiac Testing

Candidates for advanced or invasive cardiac testing include those with 1) typical or atypical cardiac symptoms and 2) an abnormal resting electrocardiogram (ECG). Exercise ECG testing without or with echocardiography may be used as the initial test. In adults with diabetes ≥ 40 years of age, measurement of coronary artery calcium is also reasonable for cardiovascular risk assessment. Pharmacologic stress echocardiography or nuclear imaging should be considered in individuals with diabetes in whom resting ECG abnormalities preclude exercise stress testing (e.g., left bundle branch block or ST-T abnormalities). In addition, individuals who require stress testing and are unable to exercise should

Table 10.3A—Cardiovascular and cardiorenal outcomes trials of available antihyperglycemic medications completed after the issuance of the FDA 2008 guidelines: DPP-4 inhibitors

	SAVOR-TIMI 53 (289) (n = 16,492)	EXAMINE (311) (n = 5,380)	TECOS (291) (n = 14,671)	CARMELINA (292,312) (n = 6,979)	CAROLINA (246) (n = 6,042)
Intervention	Saxagliptin/placebo	Alogliptin/placebo	Sitagliptin/placebo	Linagliptin/placebo	Linagliptin/glimepiride
Main inclusion criteria	Type 2 diabetes and history of or multiple risk factors for CVD	Type 2 diabetes and ACS within 15–90 days before randomization	Type 2 diabetes and preexisting CVD	Type 2 diabetes and high CV and renal risk	Type 2 diabetes and high CV risk
A1C inclusion criteria (%)	≥6.5	6.5–11.0	6.5–8.0	6.5–10.0	6.5–8.5
Age (years)*	65.1	61.0	65.4	65.8	64.0
Race (% White)	75.2	72.7	67.9	80.2	73.0
Sex (% male)	66.9	67.9	70.7	62.9	60.0
Diabetes duration (years)*	10.3	7.1	11.6	14.7	6.2
Median follow-up (years)	2.1	1.5	3.0	2.2	6.3
Statin use (%)	78	91	80	71.8	64.1
Metformin use (%)	70	66	82	54.8	82.5
Prior CVD/CHF (%)	78/13	100/28	74/18	57/26.8	34.5/4.5
Mean baseline A1C (%)	8.0	8.0	7.2	7.9	7.2
Mean difference in A1C between groups at end of treatment (%)	−0.3†	−0.3†	−0.3†	−0.36†	0
Year started/reported	2010/2013	2009/2013	2008/2015	2013/2018	2010/2019
Primary outcome‡	3-point MACE 1.00 (0.89–1.12)	3-point MACE 0.96 (95% UL ≤1.16)	4-point MACE 0.98 (0.89–1.08)	3-point MACE 1.02 (0.89–1.17)	3-point MACE 0.98 (0.84–1.14)
Key secondary outcome‡	Expanded MACE 1.02 (0.94–1.11)	4-point MACE 0.95 (95% UL ≤1.14)	3-point MACE 0.99 (0.89–1.10)	Kidney composite (ESRD, sustained ≥40% decrease in eGFR, or renal death) 1.04 (0.89–1.22)	4-point MACE 0.99 (0.86–1.14)
Cardiovascular death‡	1.03 (0.87–1.22)	0.85 (0.66–1.10)	1.03 (0.89–1.19)	0.96 (0.81–1.14)	1.00 (0.81–1.24)
MI‡	0.95 (0.80–1.12)	1.08 (0.88–1.33)	0.95 (0.81–1.11)	1.12 (0.90–1.40)	1.03 (0.82–1.29)
Stroke‡	1.11 (0.88–1.39)	0.91 (0.55–1.50)	0.97 (0.79–1.19)	0.91 (0.67–1.23)	0.86 (0.66–1.12)
HF hospitalization‡	1.27 (1.07–1.51)	1.19 (0.90–1.58)	1.00 (0.83–1.20)	0.90 (0.74–1.08)	1.21 (0.92–1.59)
Unstable angina hospitalization‡	1.19 (0.89–1.60)	0.90 (0.60–1.37)	0.90 (0.70–1.16)	0.87 (0.57–1.31)	1.07 (0.74–1.54)
All-cause mortality‡	1.11 (0.96–1.27)	0.88 (0.71–1.09)	1.01 (0.90–1.14)	0.98 (0.84–1.13)	0.91 (0.78–1.06)
Worsening nephropathy‡§	1.08 (0.88–1.32)	—	—	Kidney composite (see above)	—

—, not assessed/reported; ACS, acute coronary syndrome; CHF, congestive heart failure; CV, cardiovascular; CVD, cardiovascular disease; DPP-4, dipeptidyl peptidase 4; eGFR, estimated glomerular filtration rate; ESRD, end-stage renal disease; GLP-1, glucagon-like peptide 1; HF, heart failure; MACE, major adverse cardiovascular event; MI, myocardial infarction; UL, upper limit. Data in this table were adapted from Cefalu et al. (313). *Age was reported as means in all trials except EXAMINE, which reported medians; diabetes duration was reported as means in all trials except SAVOR-TIMI 53 and EXAMINE, which reported medians. †Significant difference in A1C between groups ($P < 0.05$). ‡Outcomes reported as hazard ratio (95% CI). §Worsening nephropathy is defined as a doubling of creatinine level, initiation of dialysis, renal transplantation, or creatinine >6.0 mg/dL (>530 mmol/L) in SAVOR-TIMI 53. Worsening nephropathy was a prespecified exploratory adjudicated outcome in SAVOR-TIMI 53.

undergo pharmacologic stress echocardiography or nuclear imaging.

Screening Asymptomatic Individuals for Atherosclerotic Cardiovascular Disease

The screening of asymptomatic individuals with high ASCVD risk is not recommended (191), in part because these high-risk people should already be receiving intensive medical therapy—an approach that provides benefits similar to those of invasive revascularization (192,193). There is also some evidence that silent ischemia may reverse over time, adding to the controversy concerning aggressive screening strategies (194). In prospective studies, coronary artery calcium has been established as an independent predictor of future ASCVD events in people with diabetes and is consistently superior to both the UK Prospective Diabetes Study (UKPDS) risk engine and the Framingham Risk Score in predicting risk in this population (195–197). However, a randomized observational trial demonstrated no clinical benefit to routine screening of asymptomatic people with type 2 diabetes and normal ECGs (198). Despite abnormal myocardial perfusion imaging in more than one in five individuals, cardiac outcomes were essentially equal (and very low) in screened versus unscreened individuals. Accordingly, indiscriminate screening is not considered cost-effective. Studies have found that a risk factor-based approach to the initial diagnostic evaluation and subsequent follow-up for coronary artery disease fails to identify which people with type 2 diabetes will have silent ischemia on screening tests (199,200).

Any benefit of newer noninvasive coronary artery disease screening methods, such as computed tomography calcium scoring and computed tomography angiography, to identify patient subgroups for different treatment strategies remains unproven in asymptomatic people with diabetes, though research is ongoing. Since asymptomatic people with diabetes with higher coronary disease burden have more future cardiac events (195,201,202), these additional imaging tests may provide reasoning for treatment intensification and/or guide informed individual decision-making and willingness for medication initiation and participation.

While coronary artery screening methods, such as calcium scoring, may improve

cardiovascular risk assessment in people with type 2 diabetes (203), their routine use leads to radiation exposure and may result in unnecessary invasive testing, such as coronary angiography and revascularization procedures. The ultimate balance of benefit, cost, and risk of such an approach in asymptomatic individuals remains controversial, particularly in the modern setting of aggressive ASCVD risk factor control.

Screening for Asymptomatic Heart Failure in People With Diabetes

People with diabetes are at an increased risk for developing heart failure, as shown in multiple longitudinal, observational studies (9,204–206). This association is not only observed in people with type 2 diabetes but also evident in people with type 1 diabetes (9,207,208). In a large multinational cohort of 750,000 people with diabetes without established cardiovascular disease, heart failure and CKD were the most frequent first cardiovascular disease manifestations (209). For a detailed review of screening, diagnosis, and treatment recommendations of heart failure in people with diabetes, the reader is further referred to the ADA consensus report “Heart Failure: An Underappreciated Complication of Diabetes. A Consensus Report of the American Diabetes Association” (7).

The increased risk for heart failure in people with diabetes is classified as the presence of stage A heart failure, i.e., an increased risk for heart failure but without symptoms, structural heart disease, or biomarker evidence of myocardial strain (210). Similar to those with stage A heart failure, people with stage B heart failure are asymptomatic but have evidence of structural heart disease or functional cardiac abnormalities, including elevated biomarkers of myocardial strain or increased filling pressures. During these asymptomatic stages of heart failure, people with diabetes are at particularly high risk for progression to symptomatic stage C and D heart failure (211,212).

Identification, risk stratification, and early treatment of risk factors in people with diabetes and asymptomatic stages of heart failure reduce the risk for progression to symptomatic heart failure (213,214). In people with type 2 diabetes, measurement of natriuretic peptides, including B-type natriuretic peptide (BNP) or N-terminal pro-BNP (NT-proBNP), identifies

people at risk for heart failure development, progression of symptoms, and heart failure-related mortality. For example, in the Canagliflozin Cardiovascular Assessment Study (CANVAS), a baseline NT-proBNP level ≥ 125 pg/mL predicted heart failure hospitalization and all-cause mortality (215). In the Examination of Cardiovascular Outcomes with Alogliptin versus Standard of Care (EXAMINE) trial, increased baseline NT-proBNP levels or an increase after a repeated measurement at 6 months was associated with an increased risk for symptomatic heart failure (216). In a combined analysis of three cohorts without disease (Atherosclerosis Risk In Communities [ARIC], Dallas Heart Study, and Multi-Ethnic Study of Atherosclerosis [MESA]), including 33% of participants with diabetes and 66% with prediabetes, biomarker screening stratifies people at high risk for incident heart failure (217). A similar association and prognostic values of increased NT-proBNP with increased cardiovascular and all-cause mortality has been reported in people with type 1 diabetes (218).

Results from several randomized controlled trials revealed that more intensive treatment of risk factors in people with increased levels of natriuretic peptides reduces the risk for symptomatic heart failure, heart failure hospitalization, and newly diagnosed left ventricular dysfunction. The NT-proBNP Guided Primary Prevention of CV Events in Diabetic Patients (PONTIAC) trial enrolled 300 people with type 2 diabetes and elevated NT-proBNP levels to usual diabetes care versus usual diabetes care plus intensified risk factor treatment in cardiac outpatient clinics (214). After 12 months, there was significant increased use of renin-angiotensin system antagonists and β -blockers in the intensive treatment group but no difference in blood pressure. The primary outcome of hospitalization or death due to cardiac disease was significantly reduced in the intensive treatment group (HR 0.35 [95% CI 0.13–0.98]; $P = 0.044$). The St Vincent’s Screening to Prevent Heart Failure Study (STOP-HF) was a randomized controlled trial that enrolled 1,374 participants with cardiovascular risk factors to receive either usual primary care or screening with BNP testing. Participants with an elevated BNP level of 50 pg/mL or higher underwent echocardiography followed by collaborative care between primary care

Table 10.3B—Cardiovascular and cardiorenal outcomes trials of available antihyperglycemic medications completed after the issuance of the FDA 2008 guidelines: GLP-1 receptor agonists

Intervention	ELIXA (260) (n = 6,068)	LEADER (255) (n = 9,340)	SUSTAIN-6 (256)* (n = 3,297)	EXSCEL (261) (n = 14,752)	REWIND (259) (n = 9,901)	Dulaglutide/placebo	PIONEER-6 (257) (n = 3,183)	Semaglutide oral/placebo
Main inclusion criteria	Type 2 diabetes and history of ACS (<180 days)	Type 2 diabetes and preexisting CVD, CKD, or HF at ≥50 years of age or CV risk at ≥60 years of age	Type 2 diabetes and preexisting CVD, HF, or CKD at ≥50 years of age or CV risk at ≥60 years of age	Type 2 diabetes with or without preexisting CVD	Type 2 diabetes and prior ASCVD event or risk factors for ASCVD	Type 2 diabetes and high CV risk (age of ≥50 years with established CVD or CKD, or age of ≥60 years with CV risk factors only)	Type 2 diabetes and high CV risk (age of ≥50 years with established CVD or CKD, or age of ≥60 years with CV risk factors only)	Type 2 diabetes and high CV risk (age of ≥50 years with established CVD or CKD, or age of ≥60 years with CV risk factors only)
A1C inclusion criteria (%)	5.5–11.0	≥7.0	≥7.0	6.5–10.0	≤9.5	None	None	None
Age (years)†	60.3	64.3	64.6	62	66.2	66	66	66
Race (% White)	75.2	77.5	83.0	75.8	75.7	72.3	72.3	72.3
Sex (% male)	69.3	64.3	60.7	62	53.7	68.4	68.4	68.4
Diabetes duration (years)†	9.3	12.8	13.9	12	10.5	14.9	14.9	14.9
Median follow-up (years)	2.1	3.8	2.1	3.2	5.4	1.3	1.3	1.3
Statin use (%)	93	72	73	74	66	85.2 (all lipid-lowering)	84.7/12.2	84.7/12.2
Metformin use (%)	66	76	73	77	81	77.4	77.4	77.4
Prior CVD/CHF (%)	100/22	81/18	60/24	73.1/16.2	32/9			
Mean baseline A1C (%)	7.7	8.7	8.7	8.0	7.4	8.2	8.2	8.2
Mean difference in A1C between groups at end of treatment (%)	−0.3‡	−0.4‡	−0.7 or −1.0§	−0.53‡	−0.61‡	−0.7	−0.7	−0.7
Year started/reported	2010/2015	2010/2016	2013/2016	2010/2017	2011/2019	2017/2019	2017/2019	2017/2019
Primary outcome§	4-point MACE 1.02 (0.89–1.17)	3-point MACE 0.87 (0.78–0.97)	3-point MACE 0.74 (0.58–0.95)	3-point MACE 0.91 (0.83–1.00)	3-point MACE 0.88 (0.79–0.99)	3-point MACE 0.79 (0.57–1.11)	3-point MACE 0.79 (0.57–1.11)	3-point MACE 0.79 (0.57–1.11)

Continued on p. S199

Table 10.3B—Continued

	ELIXA (260) (n = 6,068)	LEADER (255) (n = 9,340)	SUSTAIN-6 (256)* (n = 3,297)	EXSEL (261) (n = 14,752)	REWIND (259) (n = 9,901)	PIONEER-6 (257) (n = 3,183)
Key secondary outcome§	Expanded MACE 1.02 (0.90–1.11)	Expanded MACE 0.88 (0.81–0.96)	Expanded MACE 0.74 (0.62–0.89)	Individual components of MACE (see below)	Composite microvascular outcome (eye or renal outcome)	Expanded MACE or HF hospitalization 0.82 (0.61–1.10)
Cardiovascular death§	0.98 (0.78–1.22)	0.78 (0.66–0.93)	0.98 (0.65–1.48)	0.88 (0.76–1.02)	0.91 (0.78–1.06)	0.49 (0.27–0.92)
MI§	1.03 (0.87–1.22)	0.86 (0.73–1.00)	0.74 (0.51–1.08)	0.97 (0.85–1.10)	0.96 (0.79–1.15)	1.18 (0.73–1.90)
Stroke§	1.12 (0.79–1.58)	0.86 (0.71–1.06)	0.61 (0.38–0.99)	0.85 (0.70–1.03)	0.76 (0.61–0.95)	0.74 (0.35–1.57)
HF hospitalization§	0.96 (0.75–1.23)	0.87 (0.73–1.05)	1.11 (0.77–1.61)	0.94 (0.78–1.13)	0.93 (0.77–1.12)	0.86 (0.48–1.55)
Unstable angina hospitalization§	1.11 (0.47–2.62)	0.98 (0.76–1.26)	0.82 (0.47–1.44)	1.05 (0.94–1.18)	1.14 (0.84–1.24)	1.56 (0.60–4.01)
All-cause mortality§	0.94 (0.78–1.13)	0.85 (0.74–0.97)	1.05 (0.74–1.50)	0.86 (0.77–0.97)	0.90 (0.80–1.01)	0.51 (0.31–0.84)
Worsening nephropathy§	—	0.78 (0.67–0.92)	0.64 (0.46–0.88)	—	0.85 (0.77–0.93)	—

—, not assessed/reported; ACS, acute coronary syndrome; ASCVD, atherosclerotic cardiovascular disease; CHF, congestive heart failure; CKD, chronic kidney disease; CV, cardiovascular; CVD, cardiovascular disease; GLP-1, glucagon-like peptide 1; HF, heart failure; MACE, major adverse cardiovascular event; MI, myocardial infarction. Data in this table were adapted from Cefalu et al. (313). *Powered to rule out a hazard ratio of 1.8; superiority hypothesis not prespecified. †Age was reported as means in all trials; diabetes duration was reported as means in all trials except EXSEL, which reported medians. ‡Significant difference in A1C between groups ($P < 0.05$). §A1C change of 0.66% with 0.5 mg and 1.05% with 1-mg dose of semaglutide. ||Outcomes reported as hazard ratio (95% CI). ¶Worsening nephropathy is defined as the new onset of urine albumin-to-creatinine ratio >300 mg/g creatinine or a doubling of the serum creatinine level and an estimated glomerular filtration rate of <45 mL/min/1.73 m², the need for continuous renal replacement therapy, or death from renal disease in LEADER and SUSTAIN-6 and as new macroalbuminuria, a sustained decline in estimated glomerular filtration rate of 30% or more from baseline, or chronic renal replacement therapy in REWIND. Worsening nephropathy was a prespecified exploratory adjudicated outcome in LEADER, SUSTAIN-6, and REWIND.

physicians and cardiovascular specialist services. Participants in the intervention group received more renin-angiotensin/aldosterone system-based therapy. The primary end point of left ventricular dysfunction and heart failure was significantly reduced in the intervention group (odds ratio 0.55 [95% CI 0.37–0.82]; $P = 0.003$). The risk for left ventricular systolic dysfunction, diastolic dysfunction, or heart failure was significantly reduced in the BNP-guided intervention group. Approximately 18% of people in the control and intervention groups had a diagnosis of diabetes. Finally, in the original Intensified Multifactorial Intervention in Patients With Type 2 Diabetes and Microalbuminuria (Steno-2) trial, 160 individuals with type 2 diabetes and microalbuminuria were enrolled, and the trial examined the effect of multifactorial treatment on cardiovascular outcomes (219). When stratified in a later analysis by NT-proBNP levels, plasma NT-proBNP levels above the median were associated with increased risk of cardiovascular disease, and a decrease in plasma NT-proBNP of 10 pg/mL for the first 2 years of intervention was associated with a 1% relative risk reduction in the primary cardiovascular end point (220).

Based on this collective evidence, the committee recommends considering screening asymptomatic adults with diabetes for the development of cardiac structural or functional abnormalities (stage B heart failure) by measurement of natriuretic peptides, including BNP or NT-proBNP levels. The biomarker threshold for abnormal values is a BNP level ≥ 50 pg/mL and NT-proBNP ≥ 125 pg/mL. Abnormal levels of natriuretic peptide will need to be evaluated in the context of each individual, using clinical judgment, and in the absence of any possible competing diagnoses, particularly recognizing conditions that may lead to increased levels of natriuretic peptide, including renal insufficiency, pulmonary disease including pulmonary hypertension and chronic obstructive lung disease, obstructive sleep apnea, ischemic and hemorrhagic stroke, and anemia. Conversely, natriuretic peptide levels may be decreased in the population with obesity, which impairs sensitivity of testing.

Risk stratification for incident heart failure (stage A) and identification of people with asymptomatic cardiac abnormalities (stage B) may prevent progression to

Table 10.3C—Cardiovascular and cardiorenal outcomes trials of available antihyperglycemic medications completed after the issuance of the FDA 2008 guidelines: SGLT2 inhibitors

Intervention	CANVAS Program (12) (n = 10,142)	DECLARE-TIMI 58 (249) (n = 17,160)	CREDENCE (247) (n = 4,401)	DAPA-CKD (250;314) (n = 4,304; 2,906 with diabetes)	VERTIS CV (254;315) (n = 8,246)	DAPA-HF (14) (n = 4,744; 1,983 with diabetes)	EMPEROR-Reduced (253) (n = 7,730; 1,856 with diabetes)	DELIVER (252) (n = 6,263; 2,807 with diabetes)
Main inclusion criteria	Type 2 diabetes and preexisting CVD	Type 2 diabetes and preexisting CVD at ≥30 years of age or multiple risk factors for ASCVD	Type 2 diabetes and established ASCVD	Albuminuric kidney disease, with or without diabetes	Type 2 diabetes and ASCVD	NYHA class II, III, or IV heart failure and an ejection fraction ≤40%, with or without diabetes	NYHA class II, III, or IV heart failure and an ejection fraction >40%, with or without diabetes	NYHA class II, III, or IV heart failure and an ejection fraction >40%, with or without diabetes
A1C inclusion criteria (%)	7.0–10.5	≥6.5	6.5–12	—	7.0–10.5	—	—	—
Age (years)†	63.3	64.0	63	61.8	64.4	66	67.2, 66.5	71.8, 71.9
Race (% White)	72.4	78.3	79.6	66.6	53.2	87.8	70.3	71.1, 69.8
Sex (% male)	71.5	64.2	62.6	66.1	66.9	70	76.6	76.5, 75.6
Diabetes duration (years)†	5.7% >10	13.5	11.0	15.8	—	12.9	—	—
Median follow-up (years)	3.1	3.6	4.2	2.6	2.4	3.5	1.5	1.3
Statin use (%)	77	75	75 (statin or ezetimibe use)	69	64.9	—	—	—
Metformin use (%)	74	77	82	57.8	29	—	51.2% (of people with diabetes)	—
Prior CVD/CHF (%)	99/10	65.6/14.4	40/10	50.4/14.8	37.4/10.9	99.9/23.1	100% with CHF	100% with CHF
Mean baseline A1C (%)	8.1	8.2	8.3	8.3	8.3	7.1% (7.8% in those with diabetes)	—	—
Mean difference in A1C between groups at end of treatment (%)	−0.3 [▲]	−0.58 [#]	−0.43 [‡]	−0.31	—	−0.48 to −0.5	—	—
Year started/reported	2010/2015	2009/2017	2013/2018	2017/2019	2013/2020	2017/2019	2017/2020	2017/2020
							2018/2022	2018/2022

Continued on p. S202

Table 10.3C—Continued

	EMPA-REG OUTCOME (11) (n = 7,020)	CANVAS Program (12) (n = 10,142)	DECLARE-TIMI 58 (249) (n = 17,160)	CREDENCE (247) (n = 4,401)	DAPA-CKD (250,314) (n = 4,304; 2,906 with diabetes)	VERTIS CV (254,315) (n = 8,246)	DAPA-HF (14) (n = 4,744; 1,983 with diabetes)	EMPEROR-Reduced (253) (n = 3,730; 1,856 (n = 5,988; 2,938 with diabetes) with diabetes)	DELIVER (252)
Primary Outcomes§	3-point MACE 0.86 (0.74–0.99)	3-point MACE 0.86 (0.75–0.97)	3-point MACE 0.93 (0.84–1.03)	ESRD, doubling of creatinine, or death from renal or CV cause 0.83 (0.73–0.95)	≥50% decline in eGFR, ESKD, or death from renal or CV cause 0.70 (0.59–0.82)	3-point MACE 0.97 (0.85–1.11)	Worsening heart failure or death from CV causes 0.74 (0.55–0.85)	CV death or HF hospitalization 0.79 (0.69–0.90)	Worsening HF or CV death 0.82 (0.73–0.92)
Key secondary outcomes	4-point MACE 0.89 (0.78–1.01)	All-cause and CV mortality (see below)	Death from any cause 0.93 (0.82–1.04)	CV death or HF hospitalization 0.69 (0.57–0.83)	≥50% decline in eGFR, ESKD, or death from renal cause 0.69 (0.57–0.80)	CV death or HF hospitalization 0.88 (0.75–1.03)	CV death or HF hospitalization 0.75 (0.65–0.85)	All HF hospitalizations (first and recurrent) 0.73 (0.61–0.88)	Total number worsening HF and CV deaths 0.77 (0.67–0.89)
		Renal composite (≥40% decrease in eGFR rate to <60 mL/min/1.73 m ² , new ESRD, or death from renal or CV causes)	3-point MACE 0.80 (0.67–0.95)	CV death or HF hospitalization 0.56 (0.45–0.68)	CV death or HF hospitalization 0.92 (0.77–1.11)	CV death or HF hospitalization 0.92 (0.77–1.11)	Mean slope of change in eGFR 1.73 (1.10–2.37) <i>P</i> < 0.0001	Rate of decline in eGFR –1.25 vs. –2.62 ml/min/1.73 m ² ; Mean change in KCCQ TSS at month 8 1.11 (1.03–1.21)	Change in KCCQ TSS at month 8 1.11 (1.03–1.21)
				Renal death, renal replacement therapy, or doubling of creatinine 0.69 (0.53–0.88)	Death from any cause 0.69 (0.53–0.88)	0.81 (0.63–1.04)		Mean change in KCCQ TSS 2.4 (1.5–3.4); All-cause mortality 0.94 (0.83–1.07)	
Cardiovascular death§	0.62 (0.49–0.77)	0.87 (0.72–1.06)	0.98 (0.82–1.17)	0.78 (0.61–1.00)	0.81 (0.58–1.12)	0.92 (0.77–1.11)	0.82 (0.69–0.98)	0.92 (0.75–1.12)	0.91 (0.76–1.09)
MIs§	0.87 (0.70–1.09)	0.89 (0.73–1.09)	0.89 (0.77–1.01)	—	—	1.04 (0.86–1.26)	—	—	—
Stroke§	1.18 (0.89–1.56)	0.87 (0.69–1.09)	1.01 (0.84–1.21)	—	—	1.06 (0.82–1.37)	—	—	—
HF hospitalizations§	0.65 (0.50–0.85)	0.67 (0.52–0.87)	0.73 (0.61–0.88)	0.61 (0.47–0.80)	—	0.70 (0.54–0.90)	0.70 (0.59–0.83)	0.69 (0.59–0.81)	0.73 (0.61–0.88)
Unstable angina hospitalizations§	0.99 (0.74–1.34)	—	—	—	—	—	—	—	—
All-cause mortality§	0.68 (0.57–0.82)	0.87 (0.74–1.01)	0.93 (0.82–1.04)	0.83 (0.68–1.02)	0.69 (0.53–0.88)	0.93 (0.80–1.08)	0.83 (0.71–0.97)	0.92 (0.77–1.10)	1.00 (0.87–1.15)
Worsening nephropathy	0.61 (0.53–0.70)	0.60 (0.47–0.77)	0.53 (0.43–0.66)	(See primary outcome)	(See primary outcome)	(See secondary outcomes)	0.71 (0.44–1.16)	Composite renal outcome 0.50 (0.32–0.77)	Composite renal outcome* 0.95 (0.73–1.24)

—, not assessed/reported; CHF, congestive heart failure; CV, cardiovascular; CVD, cardiovascular disease; eGFR, estimated glomerular filtration rate; ESRD, end-stage renal disease; HF, heart failure; KCCQ TSS, Kansas City Cardiomyopathy Questionnaire Total Symptom Score; MACE, major adverse cardiovascular event; MI, myocardial infarction; SGLT2, sodium-glucose cotransporter 2; NYHA, New York Heart Association. Data in this table were adapted from Cefalu et al. (31). *Baseline characteristics for EMPEROR-Reduced displayed as empagliflozin, placebo. +Age was reported as means in all trials; diabetes duration was reported as means in all trials except EMPA-REG OUTCOME, which reported as percentage of population with diabetes duration >10 years, and DECLARE-TIMI 58, which reported median. †Significant difference in A1C between groups ($P < 0.05$). ||Definitions of worsening nephropathy differed between trials. **Composite outcome in EMPEROR-Preserved: time to first occurrence of chronic dialysis, renal transplantation; sustained reduction of ≥40% in eGFR, sustained eGFR <15 mL/min/1.73 m² for individuals with baseline eGFR ≥30 mL/min/1.73 m².

the symptomatic stages of heart failure (stages C and D). People with diabetes and an elevated natriuretic peptide level without any symptoms of heart failure should be considered to have stage B heart failure, as there is evidence for increased filling pressure and wall strain. In people with diabetes and an abnormal natriuretic peptide level, echocardiography is recommended as the next step to screen for structural heart disease and echocardiographic Doppler indices for evidence of diastolic dysfunction and increased filling pressures (221). At this stage, an interprofessional approach, which should include a cardiovascular disease specialist, is recommended to implement a guideline-directed medical treatment strategy, which may reduce the risk of progression to symptomatic stages of heart failure (213). The recommendations for screening and treatment of heart failure in people with diabetes discussed in this section are consistent with the ADA consensus report on heart failure (7) and with current American Heart Association/American College of Cardiology/Heart Failure Society of America guidelines for the management of heart failure (210).

Screening for Asymptomatic Peripheral Artery Disease in People With Diabetes

The risk for PAD in people with diabetes is higher than that in people without diabetes (222–224). In the PAD Awareness, Risk, and Treatment: New Resources for Survival (PARTNERS) program, 30% of people aged 50–69 years with a history of cigarette smoking or diabetes, or aged ≥70 years regardless of risk factors, had PAD (225). Similarly, in other screening studies, 26% of people with diabetes have been shown to have PAD (226), and diabetes increased the odds of having PAD by 85% (227). Notably, classical symptoms of claudication are uncommon, and almost half of people with newly diagnosed PAD were asymptomatic (225). Conversely, up to two-thirds of people with asymptomatic PAD have been shown to have comorbid diabetes (228). Risk factors associated with an increased risk for PAD in people with diabetes include age, smoking, hypertension, dyslipidemia, worse glycemic control, longer duration of diabetes, neuropathy, and retinopathy as well as a prior history of cardiovascular disease (229,230). In

addition, the presence of microvascular disease is associated with adverse outcomes in people with PAD, including an increased risk for future limb amputation (231,232).

Screening for asymptomatic PAD may lead to early detection and treatment strategies to reduce the risk for progression of PAD and limb preservation. In addition, secondary prevention of PAD has been shown to reduce adverse cardiovascular and limb outcomes. While a positive screening test for PAD in an asymptomatic population has been associated with increased cardiovascular event rates (233,234), prospective, randomized studies addressing whether screening for PAD in people with diabetes improves long-term limb outcomes and cardiovascular event rates are limited. In the randomized controlled Viborg Vascular (VIVA) trial, 50,156 participants were randomized to combined vascular screening for abdominal aortic aneurysm, PAD, and hypertension or to no screening. Vascular screening was associated with increased pharmacologic therapy (antiplatelet, lipid-lowering, and antihypertensive therapy), reduced in-hospital time for PAD and CAD, and reduced mortality (235). Therefore, the committee recommends screening for asymptomatic PAD using ankle-brachial index in people with diabetes at high risk for PAD, including any of the following: age ≥50 years, diabetes with duration ≥10 years, comorbid microvascular disease, clinical evidence of foot complications, or any end-organ damage from diabetes. Therefore, the committee recommends screening people with diabetes and high risk for PAD, including age ≥50 years, diabetes with duration ≥10 years, microvascular disease, clinical evidence of foot complications, or any end-organ damage from diabetes.

Lifestyle and Pharmacologic Interventions

Intensive lifestyle intervention focusing on weight loss through decreased caloric intake and increased physical activity, as performed in the Action for Health in Diabetes (Look AHEAD) trial, may be considered for improving glucose control, fitness, and some ASCVD risk factors (236). Individuals at increased ASCVD risk should receive statin, ACE inhibitor, or ARB therapy if the individual has hypertension, and possibly aspirin, unless there

are contraindications to a particular drug class.

Clear cardiovascular benefit exists for ACE inhibitor or ARB therapy in people with diabetes. The Heart Outcomes Prevention Evaluation (HOPE) study randomized 9,297 individuals aged ≥55 years with a history of vascular disease or diabetes plus one other cardiovascular risk factor to either ramipril or placebo. Ramipril significantly reduced cardiovascular and all-cause mortality, MI, and stroke (237). ACE inhibitors or ARB therapy also have well-established long-term benefit in people with diabetes and diabetic kidney disease or hypertension, and these agents are recommended for hypertension management in people with known ASCVD (particularly coronary artery disease) (65,66,238). People with type 2 diabetes and CKD should be considered for treatment with finerenone to reduce cardiovascular outcomes and the risk of CKD progression (239–242). β-Blockers should be used in individuals with active angina or HFrEF and for 3 years after MI in those with preserved left ventricular function (243,244).

Glucose-Lowering Therapies and Cardiovascular Outcomes

In 2008, the FDA issued guidance for industry to perform cardiovascular outcomes trials for all new medications for the treatment of type 2 diabetes amid concerns of increased cardiovascular risk (245). Previously approved diabetes medications were not subject to the guidance. Recently published cardiovascular outcomes trials have provided additional data on cardiovascular and renal outcomes in people with type 2 diabetes with cardiovascular disease or at high risk for cardiovascular disease (**Table 10.3A**, **Table 10.3B**, and **Table 10.3C**). An expanded review of the effects of glucose-lowering and other therapies in people with CKD is included in Section 11, “Chronic Kidney Disease and Risk Management.”

Cardiovascular outcomes trials of dipeptidyl peptidase 4 (DPP-4) inhibitors have all, so far, not shown cardiovascular benefits relative to placebo. In addition, the CAROLINA (Cardiovascular Outcome Study of Linagliptin Versus Glimepiride in Type 2 Diabetes) study demonstrated noninferiority between a DPP-4 inhibitor, linagliptin, and a sulfonylurea, glimepiride, on cardiovascular outcomes despite

lower rates of hypoglycemia in the liniagliptin treatment group (246). However, results from other new agents have provided a mix of results.

SGLT2 Inhibitor Trials

The BI 10773 (Empagliflozin) Cardiovascular Outcome Event Trial in Type 2 Diabetes Mellitus Patients (EMPA-REG OUTCOME) was a randomized, double-blind trial that assessed the effect of empagliflozin, an SGLT2 inhibitor, versus placebo on cardiovascular outcomes in 7,020 people with type 2 diabetes and existing cardiovascular disease. Study participants had a mean age of 63 years, 57% had diabetes for over 10 years, and 99% had established cardiovascular disease. EMPA-REG OUTCOME showed that over a median follow-up of 3.1 years, treatment reduced the composite outcome of MI, stroke, and cardiovascular death by 14% (absolute rate 10.5% vs. 12.1% in the placebo group, HR in the empagliflozin group 0.86 [95% CI 0.74–0.99]; $P = 0.04$ for superiority) and cardiovascular death by 38% (absolute rate 3.7% vs. 5.9%, HR 0.62 [95% CI 0.49–0.77]; $P < 0.001$) (11).

Two large outcomes trials of the SGLT2 inhibitor canagliflozin have been conducted that separately assessed 1) the cardiovascular effects of treatment in individuals at high risk for major adverse cardiovascular events (12) and 2) the impact of canagliflozin therapy on cardiorenal outcomes in people with diabetes-related CKD (247). First, the CANVAS Program integrated data from two trials. The CANVAS trial that started in 2009 was partially unblinded prior to completion because of the need to file interim cardiovascular outcomes data for regulatory approval of the drug (248). Thereafter, the postapproval CANVAS-Renal (CANVAS-R) trial was started in 2014. Combining both trials, 10,142 participants with type 2 diabetes were randomized to canagliflozin or placebo and were followed for an average of 3.6 years. The mean age of individuals was 63 years, and 66% had a history of cardiovascular disease. The combined analysis of the two trials found that canagliflozin significantly reduced the composite outcome of cardiovascular death, MI, or stroke versus placebo (occurring in 26.9 vs. 31.5 participants per 1,000 patient-years; HR 0.86 [95% CI 0.75–0.97]). The specific estimates for canagliflozin

versus placebo on the primary composite cardiovascular outcome were HR 0.88 (95% CI 0.75–1.03) for the CANVAS trial and 0.82 (0.66–1.01) for CANVAS-R, with no heterogeneity found between trials. Of note, there was an increased risk of lower-limb amputation with canagliflozin (6.3 vs. 3.4 participants per 1,000 patient-years; HR 1.97 [95% CI 1.41–2.75]) (12). Second, the Canagliflozin and Renal Events in Diabetes with Established Nephropathy Clinical Evaluation (CREDENCE) trial randomized 4,401 people with type 2 diabetes and chronic diabetes-related kidney disease ($\text{UACR} > 300 \text{ mg/g}$ and $\text{eGFR} 30 \text{ to } < 90 \text{ mL/min/1.73 m}^2$) to canagliflozin 100 mg daily or placebo (247). The primary outcome was a composite of end-stage kidney disease, doubling of serum creatinine, or death from renal or cardiovascular causes. The trial was stopped early due to conclusive evidence of efficacy identified during a prespecified interim analysis with no unexpected safety signals. The risk of the primary composite outcome was 30% lower with canagliflozin treatment than with placebo (HR 0.70 [95% CI 0.59–0.82]). Moreover, it reduced the prespecified end point of end-stage kidney disease alone by 32% (HR 0.68 [95% CI 0.54–0.86]). Canagliflozin was additionally found to have a lower risk of the composite of cardiovascular death, MI, or stroke (HR 0.80 [95% CI 0.67–0.95]) as well as lower risk of hospitalizations for heart failure (HR 0.61 [95% CI 0.47–0.80]) and of the composite of cardiovascular death or hospitalization for heart failure (HR 0.69 [95% CI 0.57–0.83]). In terms of safety, no significant increase in lower-limb amputations, fractures, AKI, or hyperkalemia was noted for canagliflozin relative to placebo in CREDENCE. An increased risk for diabetic ketoacidosis was noted, however, with 2.2 and 0.2 events per 1,000 patient-years noted in the canagliflozin and placebo groups, respectively (HR 10.80 [95% CI 1.39–83.65]) (247).

The Dapagliflozin Effect on Cardiovascular Events-Thrombosis in Myocardial Infarction 58 (DECLARE-TIMI 58) trial was another randomized, double-blind trial that assessed the effects of dapagliflozin versus placebo on cardiovascular and renal outcomes in 17,160 people with type 2 diabetes and established ASCVD or multiple risk factors for ASCVD (249). Study participants had a mean age of 64 years, with ~40% of study participants having established

ASCVD at baseline—a characteristic of this trial that differs from other large cardiovascular trials where a majority of participants had established cardiovascular disease. DECLARE-TIMI 58 met the prespecified criteria for noninferiority to placebo with respect to major adverse cardiovascular events but did not show a lower rate of major adverse cardiovascular events when compared with placebo (8.8% in the dapagliflozin group and 9.4% in the placebo group; HR 0.93 [95% CI 0.84–1.03]; $P = 0.17$). A lower rate of cardiovascular death or hospitalization for heart failure was noted (4.9% vs. 5.8%; HR 0.83 [95% CI 0.73–0.95]; $P = 0.005$), which reflected a lower rate of hospitalization for heart failure (HR 0.73 [95% CI 0.61–0.88]). No difference was seen in cardiovascular death between groups.

In the Dapagliflozin and Prevention of Adverse Outcomes in Chronic Kidney Disease (DAPA-CKD) trial (250), 4,304 individuals with CKD ($\text{UACR} 200\text{--}5,000 \text{ mg/g}$ and $\text{eGFR} 25\text{--}75 \text{ mL/min/1.73 m}^2$), with or without diabetes, were randomized to dapagliflozin 10 mg daily or placebo. The primary outcome was a composite of sustained decline in eGFR of at least 50%, end-stage kidney disease, or death from renal or cardiovascular causes. Over a median follow-up period of 2.4 years, a primary outcome event occurred in 9.2% of participants in the dapagliflozin group and 14.5% of those in the placebo group. The risk of the primary composite outcome was significantly lower with dapagliflozin therapy compared with placebo (HR 0.61 [95% CI 0.51–0.72]), as were the risks for a renal composite outcome of sustained decline in eGFR of at least 50%, end-stage kidney disease, or death from renal causes (HR 0.56 [95% CI 0.45–0.68]), and a composite of cardiovascular death or hospitalization for heart failure (HR 0.71 [95% CI 0.55–0.92]). The effects of dapagliflozin therapy were similar in individuals with and without type 2 diabetes.

Results of the Dapagliflozin and Prevention of Adverse Outcomes in Heart Failure (DAPA-HF) trial, the Empagliflozin Outcome Trial in Patients With Chronic Heart Failure and a Reduced Ejection Fraction (EMPEROR-Reduced), Empagliflozin Outcome Trial in Patients With Chronic Heart Failure With Preserved Ejection Fraction (EMPEROR-Preserved), Effects of Dapagliflozin on Biomarkers, Symptoms and Functional Status in

Patients With PRESERVED Ejection Fraction Heart Failure (PRESERVED-HF), and Dapagliflozin Evaluation to Improve the Lives of Patients with Preserved Ejection Fraction Heart Failure (DELIVER), which assessed the effects of dapagliflozin and empagliflozin in individuals with established heart failure (14,242,251–253), are described below in GLUCOSE-LOWERING THERAPIES AND HEART FAILURE.

The Evaluation of Ertugliflozin Efficacy and Safety Cardiovascular Outcomes Trial (VERTIS CV) (254) was a randomized, double-blind trial that established the effects of ertugliflozin versus placebo on cardiovascular outcomes in 8,246 people with type 2 diabetes and established ASCVD. Participants were assigned to the addition of 5 mg or 15 mg of ertugliflozin or to placebo once daily to background standard care. Study participants had a mean age of 64.4 years and a mean duration of diabetes of 13 years at baseline and were followed for a median of 3.0 years. VERTIS CV met the prespecified criteria for noninferiority of ertugliflozin to placebo with respect to the primary outcome of major adverse cardiovascular events (11.9% in the pooled ertugliflozin group and 11.9% in the placebo group; HR 0.97 [95% CI 0.85–1.11]; $P < 0.001$). Ertugliflozin was not superior to placebo for the key secondary outcomes of death from cardiovascular causes or hospitalization for heart failure; death from cardiovascular causes; or the composite of death from renal causes, renal replacement therapy, or doubling of the serum creatinine level. The HR for a secondary outcome of hospitalization for heart failure (ertugliflozin vs. placebo) was 0.70 [95% CI 0.54–0.90], consistent with findings from other SGLT2 inhibitor cardiovascular outcomes trials.

GLP-1 Receptor Agonist Trials

The Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results (LEADER) trial was a randomized, double-blind trial that assessed the effect of liraglutide, a glucagon-like peptide 1 (GLP-1) receptor agonist, versus placebo on cardiovascular outcomes in 9,340 people with type 2 diabetes at high risk for cardiovascular disease or with cardiovascular disease (256). Study participants had a mean age of 64 years and a mean duration of diabetes of nearly 13 years. Over 80% of study participants had established cardiovascular

disease. After a median follow-up of 3.8 years, LEADER showed that the primary composite outcome (MI, stroke, or cardiovascular death) occurred in fewer participants in the treatment group (13.0%) than in the placebo group (14.9%) (HR 0.87 [95% CI 0.78–0.97]; $P < 0.001$ for noninferiority; $P = 0.01$ for superiority). Deaths from cardiovascular causes were significantly reduced in the liraglutide group (4.7%) compared with the placebo group (6.0%) (HR 0.78 [95% CI 0.66–0.93]; $P = 0.007$) (256).

Results from a moderate-sized trial of another GLP-1 receptor agonist, semaglutide, were consistent with the LEADER trial (257). Semaglutide is a once-weekly GLP-1 receptor agonist approved by the FDA for the treatment of type 2 diabetes. The Trial to Evaluate Cardiovascular and Other Long-term Outcomes With Semaglutide in Subjects With Type 2 Diabetes (SUSTAIN-6) was the initial randomized trial powered to test noninferiority of semaglutide for the purpose of regulatory approval (257). In this study, 3,297 people with type 2 diabetes were randomized to receive once-weekly semaglutide (0.5 mg or 1.0 mg) or placebo for 2 years. The primary outcome (the first occurrence of cardiovascular death, nonfatal MI, or nonfatal stroke) occurred in 108 individuals (6.6%) in the semaglutide group vs. 146 individuals (8.9%) in the placebo group (HR 0.74 [95% CI 0.58–0.95]; $P < 0.001$). More individuals discontinued treatment in the semaglutide group because of adverse events, mainly gastrointestinal. The cardiovascular effects of the oral formulation of semaglutide compared with placebo have been assessed in Peptide Innovation for Early Diabetes Treatment (PIONEER) 6, a preapproval trial designed to rule out an unacceptable increase in cardiovascular risk (257). In this trial of 3,183 people with type 2 diabetes and high cardiovascular risk followed for a median of 15.9 months, oral semaglutide was noninferior to placebo for the primary composite outcome of cardiovascular death, nonfatal MI, or nonfatal stroke (HR 0.79 [95% CI 0.57–1.11]; $P < 0.001$ for noninferiority) (257). The cardiovascular effects of this formulation of semaglutide will be further tested in a large, longer-term outcomes trial.

The Harmony Outcomes trial randomized 9,463 people with type 2 diabetes and cardiovascular disease to once-weekly subcutaneous albiglutide or matching placebo, in addition to their standard care

(258). Over a median duration of 1.6 years, the GLP-1 receptor agonist reduced the risk of cardiovascular death, MI, or stroke to an incidence rate of 4.6 events per 100 person-years in the albiglutide group vs. 5.9 events in the placebo group (HR 0.78, $P = 0.0006$ for superiority) (258). This agent is not currently available for clinical use.

The Researching Cardiovascular Events With a Weekly Incretin in Diabetes (REWIND) trial was a randomized, double-blind, placebo-controlled trial that assessed the effect of the once-weekly GLP-1 receptor agonist dulaglutide versus placebo on major adverse cardiovascular events in ~9,990 people with type 2 diabetes at risk for cardiovascular events or with a history of cardiovascular disease (259). Study participants had a mean age of 66 years and a mean duration of diabetes of ~10 years. Approximately 32% of participants had history of atherosclerotic cardiovascular events at baseline. After a median follow-up of 5.4 years, the primary composite outcome of nonfatal MI, nonfatal stroke, or death from cardiovascular causes occurred in 12.0% and 13.4% of participants in the dulaglutide and placebo treatment groups, respectively (HR 0.88 [95% CI 0.79–0.99]; $P = 0.026$). These findings equated to incidence rates of 2.4 and 2.7 events per 100 person-years, respectively. The results were consistent across the subgroups of individuals with and without history of CV events. All-cause mortality did not differ between groups ($P = 0.067$).

The Evaluation of Lixisenatide in Acute Coronary Syndrome (ELIXA) trial studied the once-daily GLP-1 receptor agonist lixisenatide on cardiovascular outcomes in people with type 2 diabetes who had had a recent acute coronary event (261). A total of 6,068 people with type 2 diabetes with a recent hospitalization for MI or unstable angina within the previous 180 days were randomized to receive lixisenatide or placebo in addition to standard care and were followed for a median of ~2.1 years. The primary outcome of cardiovascular death, MI, stroke, or hospitalization for unstable angina occurred in 406 individuals (13.4%) in the lixisenatide group vs. 399 (13.2%) in the placebo group (HR 1.2 [95% CI 0.89–1.17]), which demonstrated the noninferiority of lixisenatide to placebo

($P < 0.001$) but did not show superiority ($P = 0.81$).

The Exenatide Study of Cardiovascular Event Lowering (EXSCEL) trial also reported results with the once-weekly GLP-1 receptor agonist extended-release exenatide and found that major adverse cardiovascular events were numerically lower with use of extended-release exenatide compared with placebo, although this difference was not statistically significant (261). A total of 14,752 people with type 2 diabetes (of whom 10,782 [73.1%] had previous cardiovascular disease) were randomized to receive extended-release exenatide 2 mg or placebo and followed for a median of 3.2 years. The primary end point of cardiovascular death, MI, or stroke occurred in 839 individuals (11.4%; 3.7 events per 100 person-years) in the exenatide group and in 905 individuals (12.2%; 4.0 events per 100 person-years) in the placebo group (HR 0.91 [95% CI 0.83–1.00]; $P < 0.001$ for non-inferiority), but exenatide was not superior to placebo with respect to the primary end point ($P = 0.06$ for superiority). However, all-cause mortality was lower in the exenatide group (HR 0.86 [95% CI 0.77–0.97]). The incidence of acute pancreatitis, pancreatic cancer, medullary thyroid carcinoma, and serious adverse events did not differ significantly between the two groups.

In summary, there are now numerous large randomized controlled trials reporting statistically significant reductions in cardiovascular events for three of the FDA-approved SGLT2 inhibitors (empagliflozin, canagliflozin, and dapagliflozin, with lesser benefits seen with ertugliflozin) and four FDA-approved GLP-1 receptor agonists (liraglutide, albiglutide [although that agent was removed from the market for business reasons], semaglutide [lower risk of cardiovascular events in a moderate-sized clinical trial but one not powered as a cardiovascular outcomes trial], and dulaglutide). Meta-analyses of the trials reported to date suggest that GLP-1 receptor agonists and SGLT2 inhibitors reduce risk of atherosclerotic major adverse cardiovascular events to a comparable degree in people with type 2 diabetes and established ASCVD (262,263). SGLT2 inhibitors also reduce risk of heart failure hospitalization and progression of kidney disease in people with established

ASCVD, multiple risk factors for ASCVD, or albuminuric kidney disease (264,265). In people with type 2 diabetes and established ASCVD, multiple ASCVD risk factors, or diabetic kidney disease, an SGLT2 inhibitor with demonstrated cardiovascular benefit is recommended to reduce the risk of major adverse cardiovascular events and/or heart failure hospitalization. In people with type 2 diabetes and established ASCVD or multiple risk factors for ASCVD, a GLP-1 receptor agonist with demonstrated cardiovascular benefit is recommended to reduce the risk of major adverse cardiovascular events. For many individuals, use of either an SGLT2 inhibitor or a GLP-1 receptor agonist to reduce cardiovascular risk is appropriate. Emerging data suggest that use of both classes of drugs will provide an additive cardiovascular and kidney outcomes benefit; thus, combination therapy with an SGLT2 inhibitor and a GLP-1 receptor agonist may be considered to provide the complementary outcomes benefits associated with these classes of medication. Evidence to support such an approach includes findings from AMPLITUDE-O (Effect of Efpeglenatide on Cardiovascular Outcomes), an outcomes trial of people with type 2 diabetes and either cardiovascular or kidney disease plus at least one other risk factor randomized to the investigational GLP-1 receptor agonist efpeglenatide or placebo (266). Randomization was stratified by current or potential use of SGLT2 inhibitor therapy, a class ultimately used by >15% of the trial participants. Over a median follow-up of 1.8 years, efpeglenatide therapy reduced the risk of incident major adverse cardiovascular events by 27% and of a composite renal outcome event by 32%. Importantly, the effects of efpeglenatide did not vary by use of SGLT2 inhibitors, suggesting that the beneficial effects of the GLP-1 receptor agonist were independent of those provided by SGLT2 inhibitor therapy (267). Efpeglenatide is currently not approved by the FDA for use in the U.S.

Prevention and Treatment of Heart Failure

Prevention of Symptomatic Heart Failure
ACE Inhibitors/ARBs and β -Blockers. Early primary prevention strategies and treatment of associated risk factors reduce incident, symptomatic heart failure and should include lifestyle intervention with diet, physical activity, weight control, and

smoking cessation (268–271). The vast majority of incident heart failure is preceded by hypertension; up to 91% of all new heart failure development in the Framingham cohort occurred in people with a previous diagnosis of hypertension (272). Therefore, management of hypertension constitutes a key goal in people with diabetes and stage A or B heart failure. For example, in the UKPDS trial, intensive blood pressure control in people with type 2 diabetes reduced the risk for heart failure by 56% (273). Similarly, in the SPRINT trial, intensive treatment of hypertension decreased the risk for development of incident heart failure by 36% (274). As discussed in the HYPERTENSION/BLOOD PRESSURE CONTROL section above, use of ACE inhibitors or ARBs is the preferred treatment strategy for management of hypertension in people with diabetes, particularly in the presence of albuminuria or coronary artery disease. People with diabetes and stage B heart failure who remain asymptomatic but have evidence of structural heart disease, including history of MI, acute coronary syndrome, or left ventricular ejection fraction (LVEF) $\leq 40\%$, should be treated with ACE inhibitors/ARBs plus β -blockers according to current treatment guidelines (210). In the landmark Studies of Left Ventricular Dysfunction (SOLVD) study, in which 15% of people had diabetes, treatment with enalapril reduced incident heart failure in people with asymptomatic left ventricular dysfunction by 20% (275). In the Survival and Ventricular Enlargement (SAVE) study, which enrolled asymptomatic people with reduced LVEF after MI, including 23% people with diabetes, treatment with captopril reduced the development of heart failure by 37% (276). Subsequent retrospective analyses from both trials revealed that concomitant use of β -blockers was associated with decreased risk of progression to symptomatic heart failure (277,278). The Carvedilol Post-Infarct Survival Control in Left Ventricular Dysfunction (CAPRICORN) study randomized people with a history of MI and reduced LVEF to treatment with carvedilol (279). Approximately half of the study participants were asymptomatic, and 23% of study participants had a history of diabetes. Treatment with carvedilol reduced mortality by 23%, and there was a 14% risk reduction for heart failure hospitalization. Finally, in the Reversal of Ventricular Remodeling With Toprol-XL

(REVERT) trial, in which 45% of the people enrolled had diabetes, metoprolol improved adverse cardiac remodeling in asymptomatic individuals with an LVEF <40% and mild left ventricular dilatation (280).

SGLT Inhibitors. As reviewed in detail in the following section, SGLT2 inhibitors constitute a key treatment approach to reduce cardiovascular disease and heart failure outcomes in people with diabetes. People with type 2 diabetes and increased cardiovascular risk or established cardiovascular disease should be treated with an SGLT2 inhibitor to prevent the development of incident heart failure. This includes people with type 2 diabetes and asymptomatic stage B heart failure. In the EMPA-REG OUTCOME trial, only 10% of study participants had a prior history of heart failure, and treatment with empagliflozin reduced the relative risk for hospitalization from heart failure by 35% (11). In the CANVAS Program, hospitalization from heart failure was reduced by 33% in people allocated to canagliflozin, and only 14% of individuals enrolled had a prior history of heart failure (12). In the DAPA-HF study, only 10% of study participants had a prior history of heart failure, and dapagliflozin reduced cardiovascular mortality and hospitalization for heart failure by 17%, which was consistent across multiple study subgroups regardless of a prior history of heart failure (249). Finally, in the Effect of Sotagliflozin on Cardiovascular and Renal Events in Participants With Type 2 Diabetes and Moderate Renal Impairment Who Are at Cardiovascular Risk (SCORED) trial, randomization to the SGLT1/2 inhibitor sotagliflozin reduced the primary outcome of death from cardiovascular causes, hospitalizations for heart failure, and urgent visits for heart failure in people with type 2 diabetes, CKD, and risk for cardiovascular disease (281). Therefore, SGLT inhibitor treatment is recommended in asymptomatic people with type 2 diabetes at risk or with established cardiovascular disease to prevent incident heart failure and hospitalization from heart failure.

Finerenone. Finerenone is a nonsteroidal MRA and has recently been studied in people with diabetes and diabetic kidney disease, including the Finerenone in Reducing Kidney Failure and Disease

Progression in Diabetic Kidney Disease (FIDELIO-DKD) and the Efficacy and Safety of Finerenone in Subjects With Type 2 Diabetes Mellitus and the Clinical Diagnosis of Diabetic Kidney Disease (FIGARO-DKD) studies. In FIDELIO-DKD, finerenone was compared with placebo for the primary outcome of kidney failure, a sustained decrease of at least 40% in the eGFR from baseline, or death from renal causes in people with type 2 diabetes and diabetic kidney disease (282). A prespecified secondary outcome was death from cardiovascular causes, nonfatal MI, nonfatal stroke, or hospitalization for heart failure, which was reduced by 13% in the finerenone group. The incidence of heart failure hospitalization occurred less in the finerenone-treated group, and only 7.7% of study participants had a prior history of heart failure. In the FIGARO-DKD trial, finerenone reduced the primary outcome of death from cardiovascular causes, nonfatal MI, nonfatal stroke, or hospitalization for heart failure (HR 0.87 [95% CI 0.76–0.98]; $P = 0.03$) in people with type 2 diabetes and diabetic kidney disease (240). Only 7.8% of all participants had a prior history of heart failure, and the incidence of hospitalization for heart failure was reduced in the finerenone-allocated treatment arm (HR 0.71 [95% CI 0.56–0.90]). Owing to these observations, treatment with finerenone is recommended in people with type 2 diabetes and diabetic kidney disease to reduce the risk of progression from stage A heart failure to symptomatic incident heart failure.

Treatment of Symptomatic Heart Failure

In general, current guideline-directed medical therapy for a history of MI and symptomatic stage C and D heart failure in people with diabetes is similar to that for people without diabetes. At these advanced stages of heart failure, a collaborative approach with a cardiovascular specialist is recommended. The treatment recommendations are detailed in current 2022 American Heart Association/American College of Cardiology/Heart Failure Society of America guidelines for the management of heart failure (210).

Glucose-Lowering Medications and Heart Failure: Discussion of Heart Failure Outcomes

Data on the effects of glucose-lowering agents on heart failure outcomes have demonstrated that thiazolidinediones have a strong and consistent relationship with increased risk of heart failure (283–285). Therefore, thiazolidinedione use should be avoided in people with symptomatic heart failure. Restrictions to use of metformin in people with medically treated heart failure were removed by the FDA in 2006 (286). Observational studies of people with type 2 diabetes and heart failure suggest that metformin users have better outcomes than individuals treated with other anti-hyperglycemic agents (287); however, no randomized trial of metformin therapy has been conducted in people with heart failure. Metformin may be used for the management of hyperglycemia in people with stable heart failure as long as kidney function remains within the recommended range for use (288).

Recent studies examining the relationship between DPP-4 inhibitors and heart failure have had mixed results. The Saxagliptin Assessment of Vascular Outcomes Recorded in Patients with Diabetes Mellitus—Thrombolysis in Myocardial Infarction 53 (SAVOR-TIMI 53) study showed that individuals treated with the DPP-4 inhibitor saxagliptin were more likely to be hospitalized for heart failure than those given placebo (3.5% vs. 2.8%, respectively) (289). However, three other cardiovascular outcomes trials—Examination of Cardiovascular Outcomes with Alogliptin versus Standard of Care (EXAMINE) (290), Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS) (291), and the Cardiovascular and Renal Microvascular Outcome Study With Linagliptin (CARMELINA) (292)—did not find a significant increase in risk of heart failure hospitalization with DPP-4 inhibitor use compared with placebo. No increased risk of heart failure hospitalization has been identified in the cardiovascular outcomes trials of the GLP-1 receptor agonists lixisenatide, liraglutide, semaglutide, exenatide once weekly, albiglutide, or dulaglutide compared with placebo (**Table 10.3B**) (255,256, 259–261).

Reduced incidence of heart failure has been observed with the use of SGLT2 inhibitors (11,247,249). In EMPA-REG

OUTCOME, the addition of empagliflozin to standard care led to a significant 35% reduction in hospitalization for heart failure compared with placebo (11). Although the majority of individuals in the study did not have heart failure at baseline, this benefit was consistent in individuals with and without a history of heart failure (13). Similarly, in CANVAS and DECLARE-TIMI 58, there were 33% and 27% reductions in hospitalization for heart failure, respectively, with SGLT2 inhibitor use versus placebo (12,249). Additional data from the CREDENCE trial with canagliflozin showed a 39% reduction in hospitalization for heart failure, and 31% reduction in the composite of cardiovascular death or hospitalization for heart failure, in a diabetic kidney disease population with albuminuria ($\text{UACR} > 300\text{--}5,000 \text{ mg/g}$) (247). These combined findings from four large outcomes trials of three different SGLT2 inhibitors are highly consistent and clearly indicate robust benefits of SGLT2 inhibitors in the prevention of heart failure hospitalizations. The EMPA-REG OUTCOME, CANVAS, DECLARE-TIMI 58, and CREDENCE trials suggested, but did not prove, that SGLT2 inhibitors would be beneficial in the treatment of people with established heart failure. More recently, the placebo-controlled DAPA-HF trial evaluated the effects of dapagliflozin on the primary outcome of a composite of worsening heart failure or cardiovascular death in individuals with New York Heart Association (NYHA) class II, III, or IV heart failure and an ejection fraction of 40% or less. Of the 4,744 trial participants, 45% had a history of type 2 diabetes. Over a median of 18.2 months, the group assigned to dapagliflozin treatment had a lower risk of the primary outcome (HR 0.74 [95% CI 0.65–0.85]), lower risk of first worsening heart failure event (HR 0.70 [95% CI 0.59–0.83]), and lower risk of cardiovascular death (HR 0.82 [95% CI 0.69–0.98]) compared with placebo. The effect of dapagliflozin on the primary outcome was consistent regardless of the presence or absence of type 2 diabetes (14).

EMPEROR-Reduced assessed the effects of empagliflozin 10 mg once daily versus placebo on a primary composite outcome of cardiovascular death or hospitalization for worsening heart failure in a population of 3,730 individuals with NYHA class II, III, or IV heart failure and an ejection fraction of 40% or less

(253). At baseline, 49.8% of participants had a history of diabetes. Over a median follow-up of 16 months, those in the empagliflozin-treated group had a reduced risk of the primary outcome (HR 0.75 [95% CI 0.65–0.86]; $P < 0.001$) and fewer total hospitalizations for heart failure (HR 0.70 [95% CI 0.58–0.85]; $P < 0.001$). The effect of empagliflozin on the primary outcome was consistent irrespective of diabetes diagnosis at baseline. The risk of a prespecified renal composite outcome (chronic dialysis, renal transplantation, or a sustained reduction in eGFR) was lower in the empagliflozin group than in the placebo group (1.6% in the empagliflozin group vs. 3.1% in the placebo group; HR 0.50 [95% CI 0.32–0.77]).

EMPEROR-Preserved, a randomized double-blinded placebo-controlled trial of 5,988 adults with NYHA functional class I–IV chronic HFpEF (LVEF >40%), evaluated the efficacy of empagliflozin 10 mg daily versus placebo on top of standard of care on the primary outcome of composite cardiovascular death or hospitalization for heart failure (242). Approximately 50% of subjects had type 2 diabetes at baseline. Over a median of 26.2 months, there was a 21% reduction (HR 0.79 [95% CI 0.69–0.90]; $P < 0.001$) of the primary outcome. The effects of empagliflozin were consistent in people with or without diabetes (242).

In the DELIVER trial, 6,263 individuals with heart failure and an ejection fraction >40% were randomized to receive either dapagliflozin or placebo (252). The primary outcome of a composite of worsening heart failure, defined as hospitalization or urgent visit for heart failure, or cardiovascular death was reduced by 18% in individuals treated with dapagliflozin compared with placebo (HR 0.82 [95% CI 0.73–0.92]; $P < 0.001$). Approximately 44% of individuals randomized to either dapagliflozin or placebo had type 2 diabetes, and results were consistent regardless of the presence of type 2 diabetes.

A large recent meta-analysis (293) of data from EMPEROR-Reduced, EMPEROR-Preserved, DAPA-HF, DELIVER, and the Effect of Sotagliflozin on Cardiovascular Events in Patients With Type 2 Diabetes Post Worsening Heart Failure (SOLOIST-WHF) trial included 21,947 individuals and demonstrated reduced risk for the composite of cardiovascular death or hospitalization for heart failure, cardiovascular

death, first hospitalization for heart failure, and all-cause mortality. The findings on the studied end points were consistent in both trials of heart failure with mildly reduced or preserved ejection fraction and in all five trials combined. Collectively, these studies indicate that SGLT2 inhibitors reduce the risk for heart failure hospitalization and cardiovascular death in a wide range of people with heart failure.

In addition to the hospitalization and mortality benefit in people with heart failure, several recent analyses have addressed whether SGLT2 inhibitor treatment improves clinical stability and functional status in individuals with heart failure. In 3,730 individuals with NYHA class II–IV heart failure with an ejection fraction of $\leq 40\%$, treatment with empagliflozin reduced the combined risk of death, hospitalization for heart failure, or an emergent/urgent heart failure visit requiring intravenous treatment and reduced the total number of hospitalizations for heart failure requiring intensive care, a vasopressor or positive inotropic drug, or mechanical or surgical intervention (294). In addition, individuals treated with empagliflozin were more likely to experience an improvement in NYHA functional class (294). In people hospitalized for acute de novo or decompensated chronic heart failure, initiation of empagliflozin treatment during hospitalization reduced the primary outcome of a composite of death from any cause, number of heart failure events and time to first heart failure event, or a 5-point or greater difference in change from baseline in the Kansas City Cardiomyopathy Questionnaire Total Symptom Score (295). Furthermore, PRESERVED-HF, a multicenter study (26 sites in the U.S.), showed that dapagliflozin treatment leads to significant improvement in both symptoms and physical limitation as well as objective measures of exercise function in people with chronic HFpEF, regardless of diabetes status (251). Finally, canagliflozin improved heart failure symptoms assessed using the Kansas City Cardiomyopathy Questionnaire Total Symptom Score, irrespective of LVEF or the presence of diabetes (296). Therefore, in people with type 2 diabetes and established HFpEF or HFrEF, an SGLT2 inhibitor with proven benefit in this patient population is recommended to reduce the risk of worsening heart failure and cardiovascular death. In addition, an SGLT2 inhibitor is recommended in

this patient population to improve symptoms, physical limitations, and quality of life. The benefits seen in this patient population likely represent a class effect, and they appear unrelated to glucose lowering, given comparable outcomes in people with heart failure with and without diabetes.

Sotagliflozin


Sotagliflozin, a dual SGLT1 and SGLT2 inhibitor, was recently approved by the FDA in the U.S. to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure in people with heart failure or type 2 diabetes, CKD, and other cardiovascular risk factors. This drug is distinct from other SGLT inhibitors, as it lowers glucose via delayed glucose absorption in the gut via inhibition of the cotransporter SGLT1 in addition to increasing urinary glucose excretion; however, it is not currently approved by the FDA for glycemic management of type 1 or type 2 diabetes. Sotagliflozin was evaluated in the SCORED trial (281) and SOLOIST-WHF trial (297). A total of 10,584 people with type 2 diabetes, CKD, and additional cardiovascular risk were enrolled in SCORED and randomized to sotagliflozin 200 mg once daily (up titrated to 400 mg once daily if tolerated) or placebo. SCORED ended early due to a lack of funding; thus, changes to the prespecified primary end points were made prior to unblinding to accommodate a lower-than-anticipated number of end point events. The primary end point of the trial was the total number of deaths from cardiovascular causes, hospitalizations for heart failure, and urgent visits for heart failure. After a median of 16 months of follow-up, the rate of primary end point events was reduced with sotagliflozin (5.6 events per 100 patient-years in the sotagliflozin group and 7.5 events per 100 patient-years in the placebo group [HR 0.74; [95% CI 0.63–0.88]; $P < 0.001$]). Sotagliflozin also reduced the risk of the secondary end point of total number of hospitalizations for heart failure and urgent visits for heart failure (3.5% in the sotagliflozin group and 5.1% in the placebo group [HR 0.67 [95% CI 0.55–0.82]; $P < 0.001$]) but not the secondary end point of deaths from cardiovascular causes. No significant between-group differences were found for the outcome of all-cause mortality or for a

composite renal outcome comprising the first occurrence of long-term dialysis, renal transplantation, or a sustained reduction in eGFR. In general, the adverse effects of sotagliflozin were similar to those seen with use of SGLT2 inhibitors, but they also included an increased rate of diarrhea potentially related to the inhibition of SGLT1. In general, the adverse effects of sotagliflozin were similar to those seen with use of SGLT2 inhibitors, but they also included an increased rate of diarrhea potentially related to the inhibition of SGLT1.

In SOLOIST-WHF, 1,222 people with type 2 diabetes who were recently hospitalized for worsening heart failure were randomized to sotagliflozin 200 mg once daily (with up titration to 400 mg once daily if tolerated) or placebo either before or within 3 days after hospital discharge. Individuals were eligible if hospitalized for signs and symptoms of heart failure (including elevated natriuretic peptide levels) requiring treatment with intravenous diuretic therapy. Exclusion criteria included end-stage heart failure, recent acute coronary syndrome or intervention, or an eGFR <30 mL/min/1.73 m². Individuals were required to be clinically stable prior to randomization, which was defined as no use of supplemental oxygen, systolic blood pressure ≥ 100 mmHg, and no need for intravenous inotropic or vasodilator therapy other than nitrates. Similar to SCORED, SOLOIST-WHF ended early due to a lack of funding, resulting in a change to the prespecified primary end point prior to unblinding to accommodate a lower-than-anticipated number of end point events. At a median follow-up of 9 months, the rate of primary end point events (the total number of cardiovascular deaths and hospitalizations and urgent visits for heart failure) was lower in the sotagliflozin group than in the placebo group (51.0 vs. 76.3; HR 0.67 [95% CI 0.52–0.85]; $P < 0.001$). No significant between-group differences were found in the rates of cardiovascular death or all-cause mortality. Both diarrhea (6.1% vs. 3.4%) and severe hypoglycemia (1.5% vs. 0.3%) were more common with sotagliflozin than with placebo. The trial was originally also intended to evaluate the effects of SGLT inhibition in people with HFpEF, and ultimately no evidence of heterogeneity of treatment effect by ejection fraction was noted. However, the relatively small percentage

of such individuals enrolled (only 21% of participants had ejection fraction $>50\%$) and the early termination of the trial limited the ability to determine the effects of sotagliflozin in HFpEF specifically (297).

One concern with expanded use of SGLT inhibition is the infrequent but serious risk of diabetic ketoacidosis, including the atypical presentation of euglycemic ketoacidosis. There are multiple proposed pathways through which SGLT inhibition results in ketosis (increased β -hydroxybutyrate and acetacetate), such as increased production due to reduction in insulin doses, increases in glucagon levels leading to increased lipolysis and ketone production, and decreased renal clearance of ketones (298,299). Thus, the use of SGLT inhibitors (whether for glycemic control or another indication) increases the susceptibility to diabetic ketoacidosis, particularly when other risk factors or situations occur (including, but not limited to, insulin pump malfunctions, significant reduction in insulin doses, and nutritional intake plans with prolonged periods of fasting or carbohydrate restriction). Although there were low rates of ketoacidosis in the cardiovascular and heart failure outcomes trials evaluating SGLT inhibition, these studies excluded individuals with type 1 diabetes and/or recent history of diabetic ketoacidosis (297,300). To decrease the risk of ketoacidosis when using SGLT inhibition in people with type 1 diabetes, it is recommended that clinicians assess the underlying susceptibility; provide education regarding the risks, symptoms, and prevention strategies; and prescribe home monitoring supplies for β -hydroxybutyrate (299,301). Use of these processes may have contributed to the lower rates of ketoacidosis seen in some of the studies of these agents for adjunctive glycemic management in people with type 1 diabetes (302–304) compared with those that did not include preventative strategies (298,305). Reassessment of susceptibility, education, and provision of monitoring supplies should reoccur throughout the duration of SGLT inhibitor treatment, particularly as preventative strategies and monitoring can minimize, but not eliminate, the risk of ketoacidosis in those who are susceptible (306,307).



*ASCVD is defined as a history of an acute coronary syndrome or MI, stable or unstable angina, coronary heart disease with or without revascularization, other arterial revascularization, stroke, or peripheral artery disease assumed to be atherosclerotic in origin.

†DKD is a clinical diagnosis marked by reduced eGFR, the presence of albuminuria, or both.

‡ Consider an SGLT2 inhibitor when your patient has established ASCVD, HF, DKD or is at high risk for ASCVD. Consider a GLP-1RA when your patient has established ASCVD or is at high risk for ASCVD.

§ Patients at high risk for ASCVD include those with end organ damage such as left ventricular hypertrophy or retinopathy or with multiple CV risk factors (e.g., age, hypertension, smoking, dyslipidemia, obesity).

! Most patients enrolled in the relevant trials were on metformin at baseline as glucose-lowering therapy.

ASCVD = atherosclerotic cardiovascular disease; CV = cardiovascular; DKD = diabetic kidney disease; eGFR = estimated glomerular filtration rate; GLP-1RA = glucagon-like peptide-1 receptor agonist; HF = heart failure; MI = myocardial infarction; SGLT2 = sodium-glucose cotransporter-2; T2D = type 2 diabetes

Figure 10.3—Approach to risk reduction with sodium-glucose cotransporter 2 inhibitor or glucagon-like peptide 1 receptor agonist therapy in conjunction with other traditional, guideline-based preventive medical therapies for blood pressure, lipids, and glycemia and antiplatelet therapy. Reprinted with permission from Das et al. (309).

Finerenone in People With Type 2 Diabetes and Chronic Kidney Disease

As discussed in detail in Section 11, “Chronic Kidney Disease and Risk Management,” people with diabetes are at an increased risk for CKD, which increases cardiovascular risk (308). Finerenone, a selective nonsteroidal MRA, has been shown in the FIDELIO-DKD trial to improve CKD outcomes in people with type 2 diabetes with stage 3 or 4 CKD and severe albuminuria (281). In the FIGARO-DKD trial, 7,437 individuals with UACR 30–300 mg/g and eGFR 25–90 mL/

min/1.73 m² or UACR 300–5,000 and eGFR ≥60 mL/min/1.73 m² on maximum dose of renin-angiotensin system blockade were randomized to receive finerenone or placebo (240). The HR of the primary outcome of cardiovascular death, nonfatal MI, nonfatal stroke, or hospitalization from heart failure was reduced by 13% in individuals treated with finerenone. A pre-specified subgroup analysis from FIGARO-DKD further revealed that in individuals without symptomatic HFrEF, finerenone reduces the risk for new-onset heart failure and improves heart failure outcomes

in people with type 2 diabetes and CKD (239). Finally, in the pooled analysis of 13,026 people with type 2 diabetes and CKD from both FIDELIO-DKD and FIGARO-DKD, the HR for the composite of cardiovascular death, nonfatal MI, nonfatal stroke, or hospitalization for heart failure as well as a composite of kidney failure, a sustained ≥57% decrease in eGFR from baseline over ≥4 weeks, or renal death were 0.86 and 0.77, respectively (241). These collective studies indicate that finerenone improves cardiovascular and renal outcomes in people with type 2

diabetes. Therefore, in people with type 2 diabetes and CKD with albuminuria treated with maximum tolerated doses of ACE inhibitor or ARB, addition of finerenone should be considered to improve cardiovascular outcomes and reduce the risk of CKD progression.

Clinical Approach

As has been carefully outlined in **Fig. 9.3** in Section 9, “Pharmacologic Approaches to Glycemic Treatment,” people with type 2 diabetes with or at high risk for ASCVD, heart failure, or CKD should be treated with a cardioprotective SGLT2 inhibitor and/or GLP-1 receptor agonist as part of the comprehensive approach to cardiovascular and kidney risk reduction. Importantly, these agents should be included in the plan of care irrespective of the need for additional glucose lowering and irrespective of metformin use. Such an approach has also been described in the ADA-endorsed American College of Cardiology “2020 Expert Consensus Decision Pathway on Novel Therapies for Cardiovascular Risk Reduction in Patients With Type 2 Diabetes” (309). **Figure 10.3**, reproduced from that decision pathway, outlines the approach to risk reduction with SGLT2 inhibitor or GLP-1 receptor agonist therapy in conjunction with other traditional, guideline-based preventive medical therapies for blood pressure, lipids, and glycemia and antiplatelet therapy.

Adoption of these agents should be reasonably straightforward in people with established cardiovascular or kidney disease who are later diagnosed with diabetes, as the cardioprotective agents can be used from the outset of diabetes management. On the other hand, incorporation of SGLT2 inhibitor or GLP-1 receptor agonist therapy in the care of individuals with more long-standing diabetes may be more challenging, particularly if individuals are using an already complex glucose-lowering plan. In such individuals, SGLT2 inhibitor or GLP-1 receptor agonist therapy may need to replace some or all of their existing medications to minimize risks of hypoglycemia and adverse side effects and potentially to minimize medication costs. Close collaboration between primary and specialty care professionals can help to facilitate these transitions in clinical care and, in

turn, improve outcomes for high-risk people with type 2 diabetes.

References

- Parker ED, Lin J, Mahoney T, et al. Economic costs of diabetes in the U.S. in 2022. *Diabetes Care* 1 November 2023 [Epub ahead of print]. DOI: 10.2337/dc23-0085
- Gæde P, Oelgaard J, Carstensen B, et al. Years of life gained by multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: 21 years follow-up on the Steno-2 randomised trial. *Diabetologia* 2016;59:2298–2307
- Gæde P, Lund-Andersen H, Parving HH, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008;358:580–591
- Khunti K, Kosiborod M, Ray KK. Legacy benefits of blood glucose, blood pressure and lipid control in individuals with diabetes and cardiovascular disease: time to overcome multifactorial therapeutic inertia? *Diabetes Obes Metab* 2018;20:1337–1341
- Ali MK, Bullard KM, Saaddine JB, Cowie CC, Imperatore G, Gregg EW. Achievement of goals in U.S. diabetes care, 1999–2010. *N Engl J Med* 2013;368:1613–1624
- Buse JB, Ginsberg HN, Bakris GL, et al; American Heart Association; American Diabetes Association. Primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Diabetes Care* 2007;30:162–172
- Pop-Busui R, Januzzi JL, Bruemmer D, et al. Heart failure: An underappreciated complication of diabetes. A consensus report of the american diabetes association. *Diabetes Care* 2022;45: 1670–1690
- Cavender MA, Steg PG, Smith SC Jr, et al; REACH Registry Investigators. Impact of diabetes mellitus on hospitalization for heart failure, cardiovascular events, and death: outcomes at 4 years from the Reduction of Atherothrombosis for Continued Health (REACH) Registry. *Circulation* 2015;132:923–931
- McAllister DA, Read SH, Kerssens J, et al. Incidence of hospitalization for heart failure and case-fatality among 3.25 million people with and without diabetes mellitus. *Circulation* 2018;138: 2774–2786
- Lam CSP, Voors AA, de Boer RA, Solomon SD, van Veldhuisen DJ. Heart failure with preserved ejection fraction: from mechanisms to therapies. *Eur Heart J* 2018;39:2780–2792
- Zinman B, Wanner C, Lachin JM, et al; EMPA-REG OUTCOME Investigators. Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015;373: 2117–2128
- Neal B, Perkovic V, Mahaffey KW, et al; CANVAS Program Collaborative Group. Canaglifllozin and cardiovascular and renal events in type 2 diabetes. *N Engl J Med* 2017;377:644–657
- Fitchett D, Butler J, van de Borne P, et al; EMPA-REG OUTCOME trial investigators. Effects of empagliflozin on risk for cardiovascular death and heart failure hospitalization across the spectrum of heart failure risk in the EMPA-REG OUTCOME trial. *Eur Heart J* 2018;39:363–370
- McMurray JJV, Solomon SD, Inzucchi SE, et al; DAPA-HF Trial Committees and Investigators. Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 2019;381:1995–2008
- Arnett C, Li Q, Kang A, et al. Sodium-glucose cotransporter 2 inhibition for the prevention of cardiovascular events in patients with type 2 diabetes mellitus: a systematic review and meta-analysis. *J Am Heart Assoc* 2020;9:e014908
- Lloyd-Jones DM, Braun LT, Ndumele CE, et al. Use of risk assessment tools to guide decision-making in the primary prevention of atherosclerotic cardiovascular disease: a special report from the American Heart Association and American College of Cardiology. *Circulation* 2019;139:e1162–e1177
- Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/APA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2018;71:e127–e248
- de Boer IH, Bangalore S, Benetos A, et al. Diabetes and hypertension: a position statement by the American Diabetes Association. *Diabetes Care* 2017;40:1273–1284
- Unger T, Borghi C, Charchar F, et al. 2020 International Society of Hypertension global hypertension practice guidelines. *Hypertension* 2020;75:1334–1357
- Williams B, Mancia G, Spiering W, et al; ESC Scientific Document Group. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J* 2018;39:3021–3104
- Ishigami J, Charleston J, Miller ER 3rd, Matsushita K, Appel LJ, Brady TM. Effects of cuff size on the accuracy of blood pressure readings: the Cuff(SZ) randomized crossover trial. *JAMA Intern Med* 2023;183:1061–1068
- Bobrie G, Genès N, Vaur L, et al. Is “isolated home” hypertension as opposed to “isolated office” hypertension a sign of greater cardiovascular risk? *Arch Intern Med* 2001;161: 2205–2211
- Sega R, Facchetti R, Bombelli M, et al. Prognostic value of ambulatory and home blood pressures compared with office blood pressure in the general population: follow-up results from the Pressioni Arteriose Monitorate e Loro Associazioni (PAMELA) study. *Circulation* 2005; 111:1777–1783
- Omboni S, Gazzola T, Carabelli G, Parati G. Clinical usefulness and cost effectiveness of home blood pressure telemonitoring: meta-analysis of randomized controlled studies. *J Hypertens* 2013;31:455–467; discussion 467–468
- Emdin CA, Rahimi K, Neal B, Callender T, Perkovic V, Patel A. Blood pressure lowering in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2015;313:603–615
- Arguedas JA, Leiva V, Wright JM. Blood pressure targets for hypertension in people with diabetes mellitus. *Cochrane Database Syst Rev* 2013;10:CD008277
- Ettehad D, Emdin CA, Kiran A, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016;387:957–967

28. Brunström M, Carlberg B. Effect of anti-hypertensive treatment at different blood pressure levels in patients with diabetes mellitus: systematic review and meta-analyses. *BMJ* 2016; 352:i717
29. Bangalore S, Kumar S, Lobach I, Messerli FH. Blood pressure targets in subjects with type 2 diabetes mellitus/impaired fasting glucose: observations from traditional and Bayesian random-effects meta-analyses of randomized trials. *Circulation* 2011;123:2799–2810
30. Thomopoulos C, Parati G, Zanchetti A. Effects of blood-pressure-lowering treatment on outcome incidence in hypertension: 10 - Should blood pressure management differ in hypertensive patients with and without diabetes mellitus? Overview and meta-analyses of randomized trials. *J Hypertens* 2017;35:922–944
31. Xie X, Atkins E, Lv J, et al. Effects of intensive blood pressure lowering on cardiovascular and renal outcomes: updated systematic review and meta-analysis. *Lancet* 2016;387:435–443
32. Wright JT Jr, Williamson JD, Whelton PK, et al.; SPRINT Research Group. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015;373:2103–2116
33. Zhang W, Zhang S, Deng Y, et al.; STEP Study Group. Trial of intensive blood-pressure control in older patients with hypertension. *N Engl J Med* 2021;385:1268–1279
34. Cushman WC, Evans GW, Byington RP, et al.; ACCORD Study Group. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1575–1585
35. Patel A, MacMahon S, Chalmers J, et al.; ADVANCE Collaborative Group. Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): a randomised controlled trial. *Lancet* 2007;370:829–840
36. Hansson L, Zanchetti A, Carruthers SG, et al.; HOT Study Group. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *Lancet* 1998;351:1755–1762
37. Reboldi G, Gentile G, Angeli F, Ambrosio G, Mancia G, Verdecchia P. Effects of intensive blood pressure reduction on myocardial infarction and stroke in diabetes: a meta-analysis in 73,913 patients. *J Hypertens* 2011;29:1253–1269
38. de Boer IH, Bakris G, Cannon CP. Individualizing blood pressure targets for people with diabetes and hypertension: comparing the ADA and the ACC/AHA recommendations. *JAMA* 2018; 319:1319–1320
39. Basu S, Sussman JB, Rigdon J, Steimle L, Denton BT, Hayward RA. Benefit and harm of intensive blood pressure treatment: derivation and validation of risk models using data from the SPRINT and ACCORD trials. *PLoS Med* 2017;14: e1002410
40. Phillips RA, Xu J, Peterson LE, Arnold RM, Diamond JA, Schussheim AE. Impact of cardiovascular risk on the relative benefit and harm of intensive treatment of hypertension. *J Am Coll Cardiol* 2018;71:1601–1610
41. Blood Pressure Lowering Treatment Trialists' Collaboration. Blood pressure-lowering treatment based on cardiovascular risk: a meta-analysis of individual patient data. *Lancet* 2014; 384:591–598
42. Sink KM, Evans GW, Shorr RI, et al. Syncope, hypotension, and falls in the treatment of hypertension: results from the randomized clinical Systolic Blood Pressure Intervention Trial. *J Am Geriatr Soc* 2018;66:679–686
43. Beddhu S, Greene T, Boucher R, et al. Intensive systolic blood pressure control and incident chronic kidney disease in people with and without diabetes mellitus: secondary analyses of two randomised controlled trials. *Lancet Diabetes Endocrinol* 2018;6:555–563
44. Ilkun OL, Greene T, Cheung AK, et al. The influence of baseline diastolic blood pressure on the effects of intensive blood pressure lowering on cardiovascular outcomes and all-cause mortality in type 2 diabetes. *Diabetes Care* 2020;43:1878–1884
45. Abalos E, Duley L, Steyn DW, Giardini C. Antihypertensive drug therapy for mild to moderate hypertension during pregnancy. *Cochrane Database Syst Rev* 2018;10:CD002252
46. Magee LA, von Dadelszen P, Rey E, et al. Less-tight versus tight control of hypertension in pregnancy. *N Engl J Med* 2015;372:407–417
47. Brown MA, Magee LA, Kenny LC, et al.; International Society for the Study of Hypertension in Pregnancy (ISSHP). Hypertensive disorders of pregnancy: ISSHP classification, diagnosis, and management recommendations for international practice. *Hypertension* 2018;72:24–43
48. Tita AT, Szychowski JM, Boggess K, et al.; Chronic Hypertension and Pregnancy (CHAP) Trial Consortium. Treatment for mild chronic hypertension during pregnancy. *N Engl J Med* 2022; 386:1781–1792
49. American College of Obstetricians and Gynecologists, Task Force on Hypertension in Pregnancy. Hypertension in pregnancy: report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstet Gynecol* 2013;122:1122–1131
50. Al-Balas M, Bozzo P, Einarsen A. Use of diuretics during pregnancy. *Can Fam Physician* 2009;55:44–45
51. Irgens HU, Reisaeter L, Irgens LM, Lie RT. Long term mortality of mothers and fathers after pre-eclampsia: population based cohort study. *BMJ* 2001;323:1213–1217
52. Sacks FM, Svetkey LP, Vollmer WM, et al.; DASH-Sodium Collaborative Research Group. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. *N Engl J Med* 2001; 344:3–10
53. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014;311: 507–520
54. Mao Y, Lin W, Wen J, Chen G. Impact and efficacy of mobile health intervention in the management of diabetes and hypertension: a systematic review and meta-analysis. *BMJ Open Diabetes Res Care* 2020;8:e001225
55. Stogios N, Kaur B, Huszti E, Vasanthan J, Nolan RP. Advancing digital health interventions as a clinically applied science for blood pressure reduction: a systematic review and meta-analysis. *Can J Cardiol* 2020;36:764–774
56. Bakris GL; Study of Hypertension and the Efficacy of Lotrel in Diabetes (SHIELD) Investigators. Achieving goal blood pressure in patients with type 2 diabetes: conventional versus fixed-dose combination approaches. *J Clin Hypertens (Greenwich)* 2003;5:202–209
57. Feldman RD, Zou GY, Vandervoort MK, Wong CJ, Nelson SA, Feagan BG. A simplified approach to the treatment of uncomplicated hypertension: a cluster randomized, controlled trial. *Hypertension* 2009;53:646–653
58. Webster R, Salam A, de Silva HA, et al.; TRIUMPH Study Group. Fixed low-dose triple combination antihypertensive medication vs usual care for blood pressure control in patients with mild to moderate hypertension in Sri Lanka: a randomized clinical trial. *JAMA* 2018;320: 566–579
59. Bangalore S, Kamalakkannan G, Parkar S, Messerli FH. Fixed-dose combinations improve medication compliance: a meta-analysis. *Am J Med* 2007;120:713–719
60. Catalá-López F, Macías Saint-Gerons D, González-Bermejo D, et al. Cardiovascular and renal outcomes of renin-angiotensin system blockade in adult patients with diabetes mellitus: a systematic review with network meta-analyses. *PLoS Med* 2016;13:e1001971
61. Palmer SC, Mavridis D, Navarese E, et al. Comparative efficacy and safety of blood pressure-lowering agents in adults with diabetes and kidney disease: a network meta-analysis. *Lancet* 2015;385:2047–2056
62. Barzilay JL, Davis BR, Bettencourt J, et al.; ALLHAT Collaborative Research Group. Cardiovascular outcomes using doxazosin vs. chlorothalidone for the treatment of hypertension in older adults with and without glucose disorders: a report from the ALLHAT study. *J Clin Hypertens (Greenwich)* 2004;6:116–125
63. Weber MA, Bakris GL, Jamerson K, et al.; ACCOMPLISH Investigators. Cardiovascular events during differing hypertension therapies in patients with diabetes. *J Am Coll Cardiol* 2010;56:77–85
64. Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;355: 253–259
65. Arnold SV, Bhatt DL, Barsness GW, et al.; American Heart Association Council on Lifestyle and Cardiometabolic Health and Council on Clinical Cardiology. Clinical management of stable coronary artery disease in patients with type 2 diabetes mellitus: a scientific statement from the American Heart Association. *Circulation* 2020; 141:e779–e806
66. Yusuf S, Teo K, Anderson C, et al.; Telmisartan Randomised AssessmeNt Study in ACE iNTolerant subjects with cardiovascular Disease (TRANSCEND) Investigators. Effects of the angiotensin-receptor blocker telmisartan on cardiovascular events in high-risk patients intolerant to angiotensin-converting enzyme inhibitors: a randomised controlled trial. *Lancet* 2008;372:1174–1183
67. Qiao Y, Shin JL, Chen TK, et al. Association between renin-angiotensin system blockade

- discontinuation and all-cause mortality among persons with low estimated glomerular filtration rate. *JAMA Intern Med* 2020;180:718–726
68. Bangalore S, Fakhri R, Toklu B, Messerli FH. Diabetes mellitus as a compelling indication for use of renin angiotensin system blockers: systematic review and meta-analysis of randomized trials. *BMJ* 2016;352:i438
 69. Carlberg B, Samuelsson O, Lindholm LH. Atenolol in hypertension: is it a wise choice? *Lancet* 2004;364:1684–1689
 70. Murphy SP, Ibrahim NE, Januzzi JL Jr. Heart failure with reduced ejection fraction: a review. *JAMA* 2020;324:488–504
 71. Yusuf S, Teo KK, Pogue J, et al.; ONTARGET Investigators. Telmisartan, ramipril, or both in patients at high risk for vascular events. *N Engl J Med* 2008;358:1547–1559
 72. Fried LF, Emanuele N, Zhang JH, et al.; VA NEPHRON-D Investigators. Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med* 2013;369:1892–1903
 73. Makani H, Bangalore S, Desouza KA, Shah A, Messerli FH. Efficacy and safety of dual blockade of the renin-angiotensin system: meta-analysis of randomised trials. *BMJ* 2013;346:f360
 74. Zhao P, Xu P, Wan C, Wang Z. Evening versus morning dosing regimen drug therapy for hypertension. *Cochrane Database Syst Rev* 2011;2011:CD004184
 75. Hermida RC, Ayala DE, Mojón A, Fernández JR. Influence of time of day of blood pressure-lowering treatment on cardiovascular risk in hypertensive patients with type 2 diabetes. *Diabetes Care* 2011;34:1270–1276
 76. Rahman M, Greene T, Phillips RA, et al. A trial of 2 strategies to reduce nocturnal blood pressure in blacks with chronic kidney disease. *Hypertension* 2013;61:82–88
 77. Nilsson E, Gasparini A, Ärnlöv J, et al. Incidence and determinants of hyperkalemia and hypokalemia in a large healthcare system. *Int J Cardiol* 2017;245:277–284
 78. Bandak G, Sang Y, Gasparini A, et al. Hyperkalemia after initiating renin-angiotensin system blockade: the Stockholm Creatinine Measurements (SCREAM) project. *J Am Heart Assoc* 2017;6:e005428
 79. Hughes-Austin JM, Rifkin DE, Beben T, et al. The relation of serum potassium concentration with cardiovascular events and mortality in community-living individuals. *Clin J Am Soc Nephrol* 2017;12:245–252
 80. James MT, Grams ME, Woodward M, et al.; CKD Prognosis Consortium. A meta-analysis of the association of estimated GFR, albuminuria, diabetes mellitus, and hypertension with acute kidney injury. *Am J Kidney Dis* 2015;66:602–612
 81. Williams B, MacDonald TM, Morant S, et al.; British Hypertension Society's PATHWAY Studies Group. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet* 2015;386:2059–2068
 82. Sato A, Hayashi K, Naruse M, Saruta T. Effectiveness of aldosterone blockade in patients with diabetic nephropathy. *Hypertension* 2003;41:64–68
 83. Mehdi UF, Adams-Huet B, Raskin P, Vega GL, Toto RD. Addition of angiotensin receptor blockade or mineralocorticoid antagonism to maximal angiotensin-converting enzyme inhibition in diabetic nephropathy. *J Am Soc Nephrol* 2009;20:2641–2650
 84. Bakris GL, Agarwal R, Chan JC, et al.; Mineralocorticoid Receptor Antagonist Tolerability Study–Diabetic Nephropathy (ARTS-DN) Study Group. Effect of finerenone on albuminuria in patients with diabetic nephropathy: a randomized clinical trial. *JAMA* 2015;314:884–894
 85. Jensen MD, Ryan DH, Apovian CM, et al.; American College of Cardiology/American Heart Association Task Force on Practice Guidelines; Obesity Society. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol* 2014;63(25 Pt B):2985–3023
 86. Eckel RH, Jakicic JM, Ard JD, et al.; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;129(Suppl. 2):S76–S99
 87. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;140:e596–e646
 88. Jia X, Al Rifai M, Ramsey DJ, et al. Association between lipid testing and statin adherence in the Veterans Affairs health system. *Am J Med* 2019;132:e693–e700
 89. Rana JS, Virani SS, Moffet HH, et al. Association of low-density lipoprotein testing after an atherosclerotic cardiovascular event with subsequent statin adherence and intensification. *Am J Med* 2022;135:603–606
 90. Tran C, Vo V, Taylor P, Koehn DA, Virani SS, Dixon DL. Adherence to lipid monitoring and its impact on treat intensification of LDL-C lowering therapies at an urban academic medical center. *J Clin Lipidol* 2022;16:491–497
 91. Chasman DI, Posada D, Subrahmanyam L, Cook NR, Stanton VP Jr, Ridker PM. Pharmacogenetic study of statin therapy and cholesterol reduction. *JAMA* 2004;291:2821–2827
 92. Meek C, Wierzbicki AS, Jewkes C, et al. Daily and intermittent rosuvastatin 5 mg therapy in statin intolerant patients: an observational study. *Curr Med Res Opin* 2012;28:371–378
 93. Mihaylova B, Emberson J, Blackwell L, et al.; Cholesterol Treatment Trialists' (CTT) Collaborators. The effects of lowering LDL cholesterol with statin therapy in people at low risk of vascular disease: meta-analysis of individual data from 27 randomised trials. *Lancet* 2012;380:581–590
 94. Baigent C, Keech A, Kearney PM, et al.; Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet* 2005;366:1267–1278
 95. Pyörälä K, Pedersen TR, Kjekshus J, Faergeman O, Olsson AG, Thorgeirsson G. Cholesterol lowering with simvastatin improves prognosis of diabetic patients with coronary heart disease. A subgroup analysis of the Scandinavian Simvastatin Survival Study (4S). *Diabetes Care* 1997;20:614–620
 96. Collins R, Armitage J, Parish S, Sleigh P; Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003;361:2005–2016
 97. Goldberg RB, Mellies MJ, Sacks FM, et al.; The CARE Investigators. Cardiovascular events and their reduction with pravastatin in diabetic and glucose-intolerant myocardial infarction survivors with average cholesterol levels: subgroup analyses in the Cholesterol and Recurrent Events (CARE) trial. *Circulation* 1998;98:2513–2519
 98. Shepherd J, Barter P, Carmena R, et al. Effect of lowering LDL cholesterol substantially below currently recommended levels in patients with coronary heart disease and diabetes: the Treating to New Targets (TNT) study. *Diabetes Care* 2006;29:1220–1226
 99. Sever PS, Poulter NR, Dahlöf B, et al. Reduction in cardiovascular events with atorvastatin in 2,532 patients with type 2 diabetes: Anglo-Scandinavian Cardiac Outcomes Trial-Lipid-Lowering Arm (ASCOT-LLA). *Diabetes Care* 2005;28:1151–1157
 100. Knopp RH, d'Emden M, Smilde JG, Pocock SJ. Efficacy and safety of atorvastatin in the prevention of cardiovascular end points in subjects with type 2 diabetes: the atorvastatin study for prevention of coronary heart disease endpoints in non-insulin-dependent diabetes mellitus (ASPEN). *Diabetes Care* 2006;29:1478–1485
 101. Colhoun HM, Betteridge DJ, Durrington PN, et al.; CARDS Investigators. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 2004;364:685–696
 102. Kearney PM, Blackwell L, Collins R, et al.; Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy of cholesterol-lowering therapy in 18,686 people with diabetes in 14 randomised trials of statins: a meta-analysis. *Lancet* 2008;371:117–125
 103. Taylor F, Huffman MD, Macedo AF, et al. Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2013;2013:CD004816
 104. Carter AA, Gomes T, Camacho X, Juurlink DN, Shah BR, Mamdani MM. Risk of incident diabetes among patients treated with statins: population based study. *BMJ* 2013;346:f2610
 105. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. *Endocr Pract* 2017;23(Suppl. 2):1–87
 106. Goldberg RB, Stone NJ, Grundy SM. The 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA guidelines on the management of blood cholesterol in diabetes. *Diabetes Care* 2020;43:1673–1678
 107. Mach F, Baigent C, Catapano AL, et al.; ESC Scientific Document Group. 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020;41:111–188

108. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: executive summary: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol* 2019; 73:3168–3209
109. Sabatine MS, Giugliano RP, Wiviott SD, et al.; Open-Label Study of Long-Term Evaluation against LDL Cholesterol (OSLER) Investigators. Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med* 2015;372:1500–1509
110. Robinson JG, Farnier M, Krempf M, et al.; ODYSSEY LONG TERM Investigators. Efficacy and safety of alirocumab in reducing lipids and cardiovascular events. *N Engl J Med* 2015; 372:1489–1499
111. Khan SU, Yedlapati SH, Lone AN, et al. PCSK9 inhibitors and ezetimibe with or without statin therapy for cardiovascular risk reduction: a systematic review and network meta-analysis. *BMJ* 2022;377:e069116
112. Cannon CP, Blazing MA, Giugliano RP, et al.; IMPROVE-IT Investigators. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;372:2387–2397
113. de Ferranti SD, de Boer IH, Fonseca V, et al. Type 1 diabetes mellitus and cardiovascular disease: a scientific statement from the American Heart Association and American Diabetes Association. *Diabetes Care* 2014;37:2843–2863
114. Cannon CP, Braunwald E, McCabe CH, et al.; Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis in Myocardial Infarction 22 Investigators. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350: 1495–1504
115. Sabatine MS, Giugliano RP, Keech AC, et al.; FOURIER Steering Committee and Investigators. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017;376:1713–1722
116. Giugliano RP, Cannon CP, Blazing MA, et al.; IMPROVE-IT (Improved Reduction of Outcomes: Vytorin Efficacy International Trial) Investigators. Benefit of adding ezetimibe to statin therapy on cardiovascular outcomes and safety in patients with versus without diabetes mellitus: results from IMPROVE-IT (Improved Reduction of Outcomes: Vytorin Efficacy International Trial). *Circulation* 2018;137:1571–1582
117. Schwartz GG, Steg PG, Szarek M, et al.; ODYSSEY OUTCOMES Committees and Investigators. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med* 2018;379:2097–2107
118. Ray KK, Colhoun HM, Szarek M, et al.; ODYSSEY OUTCOMES Committees and Investigators. Effects of alirocumab on cardiovascular and metabolic outcomes after acute coronary syndrome in patients with or without diabetes: a prespecified analysis of the ODYSSEY OUTCOMES randomised controlled trial. *Lancet Diabetes Endocrinol* 2019;7:618–628
119. Sabatine MS, Leiter LA, Wiviott SD, et al. Cardiovascular safety and efficacy of the PCSK9 inhibitor evolocumab in patients with and without diabetes and the effect of evolocumab on glycaemia and risk of new-onset diabetes: a prespecified analysis of the FOURIER randomised controlled trial. *Lancet Diabetes Endocrinol* 2017;5:941–950
120. Moriarty PM, Jacobson TA, Bruckert E, et al. Efficacy and safety of alirocumab, a monoclonal antibody to PCSK9, in statin-intolerant patients: design and rationale of ODYSSEY ALTERNATIVE, a randomized phase 3 trial. *J Clin Lipidol* 2014;8:554–561
121. Zhang XL, Zhu QQ, Zhu L, et al. Safety and efficacy of anti-PCSK9 antibodies: a meta-analysis of 25 randomized, controlled trials. *BMC Med* 2015;13:123
122. Giugliano RP, Pedersen TR, Saver JL, et al.; FOURIER Investigators. Stroke prevention with the PCSK9 (proprotein convertase subtilisin-kexin type 9) inhibitor evolocumab added to statin in high-risk patients with stable atherosclerosis. *Stroke* 2020;51:1546–1554
123. Ray KK, Wright RS, Kallend D, et al.; ORION-10 and ORION-11 Investigators. Two phase 3 trials of inclisiran in patients with elevated LDL cholesterol. *N Engl J Med* 2020;382:1507–1519
124. University of Oxford. A randomized trial assessing the effects of inclisiran on clinical outcomes among people with cardiovascular disease (ORION-4). In: ClinicalTrials.gov. Bethesda, MD, National Library of Medicine. NLM Identifier: NCT03705234. Accessed 10 October 2023. Available from <https://clinicaltrials.gov/ct2/show/NCT03705234>
125. Cheeley MK, Saseen JJ, Agarwala A, et al. NLA scientific statement on statin intolerance: a new definition and key considerations for ASCVD risk reduction in the statin intolerant patient. *J Clin Lipidol* 2022;16:361–375
126. Moriarty PM, Thompson PD, Cannon CP, et al.; ODYSSEY ALTERNATIVE Investigators. Efficacy and safety of alirocumab vs ezetimibe in statin-intolerant patients, with a statin rechallenge arm: the ODYSSEY ALTERNATIVE randomized trial. *J Clin Lipidol* 2015;9:758–769
127. Moriarty PM, Thompson PD, Cannon CP, et al. Efficacy and safety of alirocumab in statin-intolerant patients over 3 years: open-label treatment period of the ODYSSEY ALTERNATIVE trial. *J Clin Lipidol* 2020;14:88–97.e2
128. Sullivan D, Olsson AG, Scott R, et al. Effect of a monoclonal antibody to PCSK9 on low-density lipoprotein cholesterol levels in statin-intolerant patients: the GAUSS randomized trial. *JAMA* 2012;308:2497–2506
129. Stroes E, Colquhoun D, Sullivan D, et al.; GAUSS-2 Investigators. Anti-PCSK9 antibody effectively lowers cholesterol in patients with statin intolerance: the GAUSS-2 randomized, placebo-controlled phase 3 clinical trial of evolocumab. *J Am Coll Cardiol* 2014;63:2541–2548
130. Cho L, Dent R, Stroes ESG, et al. Persistent safety and efficacy of evolocumab in patients with statin intolerance: a subset analysis of the OSLER open-label extension studies. *Cardiovasc Drugs Ther* 2018;32:365–372
131. Nissen SE, Stroes E, Dent-Acosta RE, et al.; GAUSS-3 Investigators. Efficacy and tolerability of evolocumab vs ezetimibe in patients with muscle-related statin intolerance: the GAUSS-3 randomized clinical trial. *JAMA* 2016;315: 1580–1590
132. Ray KK, Stoekenbroek RM, Kallend D, et al. Effect of 1 or 2 doses of inclisiran on low-density lipoprotein cholesterol levels: one-year follow-up of the ORION-1 randomized clinical trial. *JAMA Cardiol* 2019;4:1067–1075
133. Ray KK, Troquay RPT, Visseren FLJ, et al. Long-term efficacy and safety of inclisiran in patients with high cardiovascular risk and elevated LDL cholesterol (ORION-3): results from the 4-year open-label extension of the ORION-1 trial. *Lancet Diabetes Endocrinol* 2023;11: 109–119
134. Dai L, Zuo Y, You Q, Zeng H, Cao S. Efficacy and safety of bempedoic acid in patients with hypercholesterolemia: a systematic review and meta-analysis of randomized controlled trials. *Eur J Prev Cardiol* 2021;28:825–833
135. Di Minno A, Lupoli R, Calcaterra I, et al. Efficacy and safety of bempedoic acid in patients with hypercholesterolemia: systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc* 2020;9:e016262
136. Nissen SE, Lincoff AM, Brennan D, et al.; CLEAR Outcomes Investigators. Bempedoic acid and cardiovascular outcomes in statin-intolerant patients. *N Engl J Med* 2023;388:1353–1364
137. Nissen SE, Menon V, Nicholls SJ, et al. Bempedoic acid for primary prevention of cardiovascular events in statin-intolerant patients. *JAMA* 2023;330:131–140
138. Berglund L, Brunzell JD, Goldberg AC, et al.; Endocrine Society. Evaluation and treatment of hypertriglyceridemia: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2012;97:2969–2989
139. Bhatt DL, Steg PG, Miller M, et al.; REDUCE-IT Investigators. Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med* 2019;380:11–22
140. Nicholls SJ, Lincoff AM, Garcia M, et al. Effect of high-dose omega-3 fatty acids vs corn oil on major adverse cardiovascular events in patients at high cardiovascular risk: the STRENGTH randomized clinical trial. *JAMA* 2020;324: 2268–2280
141. Singh IM, Shishehbor MH, Ansell BJ. High-density lipoprotein as a therapeutic target: a systematic review. *JAMA* 2007;298:786–798
142. Keech A, Simes RJ, Barter P, et al.; FIELD Study Investigators. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005;366:1849–1861
143. Jones PH, Davidson MH. Reporting rate of rhabdomyolysis with fenofibrate + statin versus gemfibrozil + any statin. *Am J Cardiol* 2005;95: 120–122
144. Ginsberg HN, Elam MB, Lovato LC, et al.; ACCORD Study Group. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1563–1574
145. Boden WE, Probstfield JL, Anderson T, et al.; AIM-HIGH Investigators. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med* 2011; 365:2255–2267
146. Landray MJ, Haynes R, Hopewell JC, et al.; HPS2-THRIVE Collaborative Group. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med* 2014;371:203–212
147. Rajpathak SN, Kumbhani DJ, Crandall J, Barzilai N, Alderman M, Ridker PM. Statin

- therapy and risk of developing type 2 diabetes: a meta-analysis. *Diabetes Care* 2009;32:1924–1929
148. Sattar N, Preiss D, Murray HM, et al. Statins and risk of incident diabetes: a collaborative meta-analysis of randomised statin trials. *Lancet* 2010;375:735–742
149. Ridker PM, Pradhan A, MacFadyen JG, Libby P, Glynn RJ. Cardiovascular benefits and diabetes risks of statin therapy in primary prevention: an analysis from the JUPITER trial. *Lancet* 2012;380:565–571
150. Mach F, Ray KK, Wiklund O, et al.; European Atherosclerosis Society Consensus Panel. Adverse effects of statin therapy: perception vs. the evidence—focus on glucose homeostasis, cognitive, renal and hepatic function, haemorrhagic stroke and cataract. *Eur Heart J* 2018;39:2526–2539
151. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002;360:7–22
152. Shepherd J, Blauw GJ, Murphy MB, et al.; PROSPER Study Group. PROspective Study of Pravastatin in the Elderly at Risk. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet* 2002;360:1623–1630
153. Trompet S, van Vliet P, de Craen AJ, et al. Pravastatin and cognitive function in the elderly. Results of the PROSPER study. *J Neurol* 2010;257:85–90
154. Yusuf S, Bosch J, Dagenais G, et al.; HOPE-3 Investigators. Cholesterol lowering in intermediate-risk persons without cardiovascular disease. *N Engl J Med* 2016;374:2021–2031
155. Giugliano RP, Mach F, Zavitz K, et al.; EBBINGHAUS Investigators. Cognitive function in a randomized trial of evolocumab. *N Engl J Med* 2017;377:633–643
156. Richardson K, Schoen M, French B, et al. Statins and cognitive function: a systematic review. *Ann Intern Med* 2013;159:688–697
157. Perk J, De Backer G, Gohlke H, et al.; European Association for Cardiovascular Prevention & Rehabilitation (EACPR); ESC Committee for Practice Guidelines (CPG). European guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). *Eur Heart J* 2012;33:1635–1701
158. Belch J, MacCuish A, Campbell I, et al.; Prevention of Progression of Arterial Disease and Diabetes Study Group; Diabetes Registry Group; Royal College of Physicians Edinburgh. The Prevention of Progression of Arterial Disease and Diabetes (POPADAD) trial: factorial randomised placebo controlled trial of aspirin and antioxidants in patients with diabetes and asymptomatic peripheral arterial disease. *BMJ* 2008;337:a1840
159. Zhang C, Sun A, Zhang P, et al. Aspirin for primary prevention of cardiovascular events in patients with diabetes: a meta-analysis. *Diabetes Res Clin Pract* 2010;87:211–218
160. De Berardis G, Sacco M, Strippoli GF, et al. Aspirin for primary prevention of cardiovascular events in people with diabetes: meta-analysis of randomised controlled trials. *BMJ* 2009;339:b4531
161. Baigent C, Blackwell L, Collins R, et al.; Antithrombotic Trialists' (ATT) Collaboration. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* 2009;373:1849–1860
162. Bowman L, Mafham M, Wallendszus K, et al.; ASCEND Study Collaborative Group. Effects of aspirin for primary prevention in persons with diabetes mellitus. *N Engl J Med* 2018;379:1529–1539
163. Gaziano JM, Brotons C, Coppolecchia R, et al.; ARRIVE Executive Committee. Use of aspirin to reduce risk of initial vascular events in patients at moderate risk of cardiovascular disease (ARRIVE): a randomised, double-blind, placebo-controlled trial. *Lancet* 2018;392:1036–1046
164. McNeil JJ, Wolfe R, Woods RL, et al.; ASPREE Investigator Group. Effect of aspirin on cardiovascular events and bleeding in the healthy elderly. *N Engl J Med* 2018;379:1509–1518
165. Pignone M, Earnshaw S, Tice JA, Pletcher MJ. Aspirin, statins, or both drugs for the primary prevention of coronary heart disease events in men: a cost-utility analysis. *Ann Intern Med* 2006;144:326–336
166. Huxley RR, Peters SA, Mishra GD, Woodward M. Risk of all-cause mortality and vascular events in women versus men with type 1 diabetes: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol* 2015;3:198–206
167. Peters SA, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia* 2014;57:1542–1551
168. Kalyani RR, Lazo M, Ouyang P, et al. Sex differences in diabetes and risk of incident coronary artery disease in healthy young and middle-aged adults. *Diabetes Care* 2014;37:830–838
169. Peters SA, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775,385 individuals and 12,539 strokes. *Lancet* 2014;383:1973–1980
170. Miedema MD, Duprez DA, Misialek JR, et al. Use of coronary artery calcium testing to guide aspirin utilization for primary prevention: estimates from the Multi-Ethnic Study of Atherosclerosis. *Circ Cardiovasc Qual Outcomes* 2014;7:453–460
171. Dimitriu-Leen AC, Scholte AJ, van Rosendaal AR, et al. Value of coronary computed tomography angiography in tailoring aspirin therapy for primary prevention of atherosclerotic events in patients at high risk with diabetes mellitus. *Am J Cardiol* 2016;117:887–893
172. Mora S, Ames JM, Manson JE. Low-dose aspirin in the primary prevention of cardiovascular disease: shared decision making in clinical practice. *JAMA* 2016;316:709–710
173. Campbell CL, Smyth S, Montalecot G, Steinbuhl SR. Aspirin dose for the prevention of cardiovascular disease: a systematic review. *JAMA* 2007;297:2018–2024
174. Jones WS, Mulder H, Wruck LM, et al.; ADAPTABLE Team. Comparative effectiveness of aspirin dosing in cardiovascular disease. *N Engl J Med* 2021;384:1981–1990
175. Davì G, Patrono C. Platelet activation and atherothrombosis. *N Engl J Med* 2007;357:2482–2494
176. Larsen SB, Grove EL, Neergaard-Petersen S, Würtz M, Hvas AM, Kristensen SD. Determinants of reduced antiplatelet effect of aspirin in patients with stable coronary artery disease. *PLoS One* 2015;10:e0126767
177. Zaccardi F, Rizzi A, Petrucci G, et al. In vivo platelet activation and aspirin responsiveness in type 1 diabetes. *Diabetes* 2016;65:503–509
178. Bethel MA, Harrison P, Sourij H, et al. Randomized controlled trial comparing impact on platelet reactivity of twice-daily with once-daily aspirin in people with type 2 diabetes. *Diabet Med* 2016;33:224–230
179. Rothwell PM, Cook NR, Gaziano JM, et al. Effects of aspirin on risks of vascular events and cancer according to bodyweight and dose: analysis of individual patient data from randomised trials. *Lancet* 2018;392:387–399
180. Levine GN, Bates ER, Bittl JA, et al. 2016 ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines: an update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention, 2011 ACCF/AHA guideline for coronary artery bypass graft surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease, 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction, 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes, and 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery. *Circulation* 2016;134:e123–e155
181. Kleindorfer DO, Towfighi A, Chaturvedi S, et al. 2021 guideline for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline from the American Heart Association/American Stroke Association. *Stroke* 2021;52:e364–e467
182. Vandvik PO, Lincoff AM, Gore JM, et al. Primary and secondary prevention of cardiovascular disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2012;141:e637S–e668S
183. Bhatt DL, Bonaca MP, Bansilal S, et al. Reduction in ischemic events with ticagrelor in diabetic patients with prior myocardial infarction in PEGASUS-TIMI 54. *J Am Coll Cardiol* 2016;67:2732–2740
184. Steg PG, Bhatt DL, Simon T, et al.; THEMIS Steering Committee and Investigators. Ticagrelor in patients with stable coronary disease and diabetes. *N Engl J Med* 2019;381:1309–1320
185. Bhatt DL, Steg PG, Mehta SR, et al.; THEMIS Steering Committee and Investigators. Ticagrelor in patients with diabetes and stable coronary artery disease with a history of previous percutaneous coronary intervention (THEMIS-PCI): a phase 3, placebo-controlled, randomised trial. *Lancet* 2019;394:1169–1180

186. Angiolillo DJ, Baber U, Sartori S, et al. Ticagrelor with or without aspirin in high-risk patients with diabetes mellitus undergoing percutaneous coronary intervention. *J Am Coll Cardiol* 2020;75:2403–2413
187. Wiebe J, Ndrepepa G, Kufner S, et al. Early aspirin discontinuation after coronary stenting: a systematic review and meta-analysis. *J Am Heart Assoc* 2021;10:e018304
188. Bhatt DL, Eikelboom JW, Connolly SJ, et al.; COMPASS Steering Committee and Investigators. Role of combination antiplatelet and anti-coagulation therapy in diabetes mellitus and cardiovascular disease: insights from the COMPASS trial. *Circulation* 2020;141:1841–1854
189. Connolly SJ, Eikelboom JW, Bosch J, et al.; COMPASS Investigators. Rivaroxaban with or without aspirin in patients with stable coronary artery disease: an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2018;391:205–218
190. Bonaca MP, Bauersachs RM, Anand SS, et al. Rivaroxaban in peripheral artery disease after revascularization. *N Engl J Med* 2020;382:1994–2004
191. Bax JJ, Young LH, Frye RL, Bonow RO, Steinberg HO; ADA. Screening for coronary artery disease in patients with diabetes. *Diabetes Care* 2007;30:2729–2736
192. Boden WE, O'Rourke RA, Teo KK, et al.; COURAGE Trial Research Group. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med* 2007;356:1503–1516
193. Frye RL, August P, Brooks MM, et al.; BARI 2D Study Group. A randomized trial of therapies for type 2 diabetes and coronary artery disease. *N Engl J Med* 2009;360:2503–2515
194. Wackers FJ, Chyun DA, Young LH, et al.; Detection of Ischemia in Asymptomatic Diabetics (DIAD) Investigators. Resolution of asymptomatic myocardial ischemia in patients with type 2 diabetes in the Detection of Ischemia in Asymptomatic Diabetics (DIAD) study. *Diabetes Care* 2007;30:2892–2898
195. Elkeles RS, Godslan IF, Feher MD, et al.; PREDICT Study Group. Coronary calcium measurement improves prediction of cardiovascular events in asymptomatic patients with type 2 diabetes: the PREDICT study. *Eur Heart J* 2008;29:2244–2251
196. Raggi P, Shaw LJ, Berman DS, Callister TQ. Prognostic value of coronary artery calcium screening in subjects with and without diabetes. *J Am Coll Cardiol* 2004;43:1663–1669
197. Anand DV, Lim E, Hopkins D, et al. Risk stratification in uncomplicated type 2 diabetes: prospective evaluation of the combined use of coronary artery calcium imaging and selective myocardial perfusion scintigraphy. *Eur Heart J* 2006;27:713–721
198. Young LH, Wackers FJ, Chyun DA, et al.; DIAD Investigators. Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *JAMA* 2009;301:1547–1555
199. Wackers FJ, Young LH, Inzucchi SE, et al.; Detection of Ischemia in Asymptomatic Diabetics Investigators. Detection of silent myocardial ischemia in asymptomatic diabetic subjects: the DIAD study. *Diabetes Care* 2004;27:1954–1961
200. Scognamiglio R, Negut C, Ramondo A, Tiengo A, Avogaro A. Detection of coronary artery disease in asymptomatic patients with type 2 diabetes mellitus. *J Am Coll Cardiol* 2006;47:65–71
201. Hadamitzky M, Hein F, Meyer T, et al. Prognostic value of coronary computed tomographic angiography in diabetic patients without known coronary artery disease. *Diabetes Care* 2010;33:1358–1363
202. Choi EK, Chun EJ, Choi SI, et al. Assessment of subclinical coronary atherosclerosis in asymptomatic patients with type 2 diabetes mellitus with single photon emission computed tomography and coronary computed tomography angiography. *Am J Cardiol* 2009;104:890–896
203. Malik S, Zhao Y, Budoff M, et al. Coronary artery calcium score for long-term risk classification in individuals with type 2 diabetes and metabolic syndrome from the Multi-Ethnic Study of Atherosclerosis. *JAMA Cardiol* 2017;2:1332–1340
204. Kannel WB, Hjortland M, Castelli WP. Role of diabetes in congestive heart failure: the Framingham study. *Am J Cardiol* 1974;34:29–34
205. He J, Ogden LG, Bazzano LA, Vuppuluri S, Loria C, Welton PK. Risk factors for congestive heart failure in US men and women: NHANES I epidemiologic follow-up study. *Arch Intern Med* 2001;161:996–1002
206. Thraulsdottir IS, Aspelund T, Thorsteinsson G, et al. The association between glucose abnormalities and heart failure in the population-based Reykjavik study. *Diabetes Care* 2005;28:612–616
207. Rosengren A, Vestberg D, Svensson AM, et al. Long-term excess risk of heart failure in people with type 1 diabetes: a prospective case-control study. *Lancet Diabetes Endocrinol* 2015;3:876–885
208. Ohkuma T, Komorita Y, Peters SAE, Woodward M. Diabetes as a risk factor for heart failure in women and men: a systematic review and meta-analysis of 47 cohorts including 12 million individuals. *Diabetologia* 2019;62:1550–1560
209. Birkeland KI, Bodegard J, Eriksson JW, et al. Heart failure and chronic kidney disease manifestation and mortality risk associations in type 2 diabetes: a large multinational cohort study. *Diabetes Obes Metab* 2020;22:1607–1618
210. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2022;145:e895–e1032
211. Segar MW, Patel KV, Vaduganathan M, et al. Association of long-term change and variability in glycemia with risk of incident heart failure among patients with type 2 diabetes: a secondary analysis of the ACCORD trial. *Diabetes Care* 2020;43:1920–1928
212. Echouffo-Tcheugui JB, Ndumele CE, Zhang S, et al. Diabetes and progression of heart failure: the Atherosclerosis Risk In Communities (ARIC) study. *J Am Coll Cardiol* 2022;79:2285–2293
213. Ledwidge M, Gallagher J, Conlon C, et al. Natriuretic peptide-based screening and collaborative care for heart failure: the STOP-HF randomized trial. *JAMA* 2013;310:66–74
214. Huelsmann M, Neuhold S, Resl M, et al. PONTIAC (NT-proBNP selected prevention of cardiac events in a population of diabetic patients without a history of cardiac disease): a prospective randomized controlled trial. *J Am Coll Cardiol* 2013;62:1365–1372
215. Januzzi JL Jr, Xu J, Li J, et al. Effects of canagliflozin on amino-terminal pro-B-type natriuretic peptide: implications for cardiovascular risk reduction. *J Am Coll Cardiol* 2020;76:2076–2085
216. Jarolim P, White WB, Cannon CP, Gao Q, Morrow DA. Serial measurement of natriuretic peptides and cardiovascular outcomes in patients with type 2 diabetes in the EXAMINE trial. *Diabetes Care* 2018;41:1510–1515
217. Pandey A, Vaduganathan M, Patel KV, et al. Biomarker-based risk prediction of incident heart failure in pre-diabetes and diabetes. *JACC Heart Fail* 2021;9:215–223
218. Rørt R, Jørgensen PG, Andersen HU, et al. Cardiovascular prognostic value of echocardiography and N terminal pro B-type natriuretic peptide in type 1 diabetes: the Thousand & 1 Study. *Eur J Endocrinol* 2020;182:481–488
219. Gaede P, Vedel P, Larsen N, Jensen GV, Parving HH, Pedersen O. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383–393
220. Gaede P, Hildebrandt P, Hess G, Parving HH, Pedersen O. Plasma N-terminal pro-brain natriuretic peptide as a major risk marker for cardiovascular disease in patients with type 2 diabetes and microalbuminuria. *Diabetologia* 2005;48:156–163
221. Redfield MM, Jacobsen SJ, Burnett JC Jr, Mahoney DW, Bailey KR, Rodeheffer RJ. Burden of systolic and diastolic ventricular dysfunction in the community: appreciating the scope of the heart failure epidemic. *JAMA* 2003;289:194–202
222. Selvin E, Erlinger TP. Prevalence of and risk factors for peripheral arterial disease in the United States: results from the National Health and Nutrition Examination Survey, 1999–2000. *Circulation* 2004;110:738–743
223. Leibson CL, Ransom JE, Olson W, Zimmerman BR, O’Fallon WM, Palumbo PJ. Peripheral arterial disease, diabetes, and mortality. *Diabetes Care* 2004;27:2843–2849
224. Murabito JM, D’Agostino RB, Silbershatz H, Wilson WF. Intermittent claudication. A risk profile from the Framingham Heart Study. *Circulation* 1997;96:44–49
225. Hirsch AT, Criqui MH, Treat-Jacobson D, et al. Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA* 2001;286:1317–1324
226. Lange S, Diehm C, Darius H, et al. High prevalence of peripheral arterial disease and low treatment rates in elderly primary care patients with diabetes. *Exp Clin Endocrinol Diabetes* 2004;112:566–573
227. Grøndal N, Søgaard R, Lindholm JS. Baseline prevalence of abdominal aortic aneurysm, peripheral arterial disease and hypertension in men aged 65–74 years from a population screening study (VIVA trial). *Br J Surg* 2015;102:902–906
228. Eason SL, Petersen NJ, Suarez-Almazor M, Davis B, Collins TC. Diabetes mellitus, smoking, and the risk for asymptomatic peripheral arterial

- disease: whom should we screen? *J Am Board Fam Pract* 2005;18:355–361
229. Adler AI, Stevens RJ, Neil A, Stratton IM, Boulton AJ, Holman RR. UKPDS 59: hyperglycemia and other potentially modifiable risk factors for peripheral vascular disease in type 2 diabetes. *Diabetes Care* 2002;25:894–899
230. Al-Delaimy WK, Merchant AT, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Effect of type 2 diabetes and its duration on the risk of peripheral arterial disease among men. *Am J Med* 2004;116:236–240
231. Beckman JA, Duncan MS, Damrauer SM, et al. Microvascular disease, peripheral artery disease, and amputation. *Circulation* 2019;140:449–458
232. Olesen KKW, Anand S, Thim T, Gyldenkerne C, Maeng M. Microvascular disease increases the risk of lower limb amputation—a western Danish cohort study. *Eur J Clin Invest* 2022;52:e13812
233. Smolderen KG, Ameli O, Chaisson CE, Heath K, Mena-Hurtado C. Peripheral artery disease screening in the community and 1-year mortality, cardiovascular events, and adverse limb events. *AJPM Focus* 2022;1:100016
234. Smolderen KG, Heath K, Scherr T, Bauzon SR, Howell AN, Mena-Hurtado C. The Nevada peripheral artery disease screening effort in a Medicare Advantage population and subsequent mortality and major adverse cardiovascular event risk. *J Vasc Surg* 2022;75:2054–2064.e3
235. Lindholt JS, Søgaard R. Population screening and intervention for vascular disease in Danish men (VIVA): a randomised controlled trial. *Lancet* 2017;390:2256–2265
236. Wing RR, Bolin P, Brancati FL, et al.; Look AHEAD Research Group. Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes. *N Engl J Med* 2013;369:145–154
237. Yusuf S, Sleight P, Pogue J, Bosch J, Davies R; Heart Outcomes Prevention Evaluation Study Investigators. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;342:145–153
238. Braunwald E, Domanski MJ, Fowler SE, et al.; PEACE Trial Investigators. Angiotensin-converting-enzyme inhibition in stable coronary artery disease. *N Engl J Med* 2004;351:2058–2068
239. Filippatos G, Anker SD, Agarwal R, et al.; FIGARO-DKD Investigators. Finerenone reduces risk of incident heart failure in patients with chronic kidney disease and type 2 diabetes: analyses from the FIGARO-DKD trial. *Circulation* 2022;145:437–447
240. Pitt B, Filippatos G, Agarwal R, et al.; FIGARO-DKD Investigators. Cardiovascular events with finerenone in kidney disease and type 2 diabetes. *N Engl J Med* 2021;385:2252–2263
241. Agarwal R, Filippatos G, Pitt B, et al.; FIDELIO-DKD and FIGARO-DKD Investigators. Cardiovascular and kidney outcomes with finerenone in patients with type 2 diabetes and chronic kidney disease: the FIDELITY pooled analysis. *Eur Heart J* 2022;43:474–484
242. Anker SD, Butler J, Filippatos G, et al.; EMPEROR-Preserved Trial Investigators. Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;385:1451–1461
243. Kezerashvili A, Marzo K, De Leon J. Beta blocker use after acute myocardial infarction in the patient with normal systolic function: when is it “ok” to discontinue? *Curr Cardiol Rev* 2012;8:77–84
244. Fihn SD, Gardin JM, Abrams J, et al.; American College of Cardiology Foundation; American Heart Association Task Force on Practice Guidelines; American College of Physicians; American Association for Thoracic Surgery; Preventive Cardiovascular Nurses Association; Society for Cardiovascular Angiography and Interventions; Society of Thoracic Surgeons. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol* 2012;60:e44–e164
245. U.S. Food and Drug Administration. Guidance for industry. Diabetes mellitus—evaluating cardiovascular risk in new antidiabetic therapies to treat type 2 diabetes. Silver Spring, MD, 2008. Accessed 10 October 2023. Available from <https://www.federalregister.gov/documents/2008/12/19/E8-30086/guidance-for-industry-on-diabetes-mellitus-evaluating-cardiovascular-risk-in-new-antidiabetic>
246. Marx N, Rosenstock J, Kahn SE, et al. Design and baseline characteristics of the CARdiovascular Outcome Trial of LINagliptin Versus Glimepiride in Type 2 Diabetes (CAROLINA). *Diab Vasc Dis Res* 2015;12:164–174
247. Perkovic V, Jardine MJ, Neal B, et al.; CREDENCE Trial Investigators. Canaglifllozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019;380:2295–2306
248. Neal B, Perkovic V, Matthews DR, et al.; CANVAS-R Trial Collaborative Group. Rationale, design and baseline characteristics of the CANAglifllozin cardioVascular Assessment Study-Renal (CANVAS-R): a randomized, placebo-controlled trial. *Diabetes Obes Metab* 2017;19:387–393
249. Wiviott SD, Raz I, Bonaca MP, et al.; DECLARE-TIMI 58 Investigators. Dapagliflozin and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2019;380:347–357
250. Heerspink HJL, Stefánsson BV, Correa-Rotter R, et al.; DAPA-CKD Trial Committees and Investigators. Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;383:1436–1446
251. Nassif ME, Windsor SL, Borlaug BA, et al. The SGLT2 inhibitor dapagliflozin in heart failure with preserved ejection fraction: a multicenter randomized trial. *Nat Med* 2021;27:1954–1960
252. Solomon SD, McMurray J JV, Claggett B, et al.; DELIVER Trial Committees and Investigators. Dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2022;387:1089–1098
253. Packer M, Anker SD, Butler J, et al.; EMPEROR-Reduced Trial Investigators. Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* 2020;383:1413–1424
254. Cannon CP, Pratley R, Dagogo-Jack S, et al.; VERTIS CV Investigators. Cardiovascular outcomes with empagliflozin in type 2 diabetes. *N Engl J Med* 2021;385:896–907
255. Marso SP, Daniels GH, Brown-Frandsen K, et al.; LEADER Steering Committee; LEADER Trial Investigators. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016;375:311–322
256. Marso SP, Bain SC, Consoli A, et al.; SUSTAIN-6 Investigators. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016;375:1834–1844
257. Husain M, Birkenfeld AL, Donsmark M, et al.; PIONEER 6 Investigators. Oral semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2019;381:841–851
258. Hernandez AF, Green JB, Janmohamed S, et al.; Harmony Outcomes committees and investigators. Albiglutide and cardiovascular outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet* 2018;392:1519–1529
259. Gerstein HC, Colhoun HM, Dagenais GR, et al.; REWIND Investigators. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *Lancet* 2019;394:121–130
260. Pfeffer MA, Claggett B, Diaz R, et al.; ELIXA Investigators. Lixisenatide in patients with type 2 diabetes and acute coronary syndrome. *N Engl J Med* 2015;373:2247–2257
261. Holman RR, Bethel MA, Mentz RJ, et al.; EXSEL Study Group. Effects of once-weekly exenatide on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2017;377:1228–1239
262. Zelniker TA, Wiviott SD, Raz I, et al. Comparison of the effects of glucagon-like peptide receptor agonists and sodium-glucose cotransporter 2 inhibitors for prevention of major adverse cardiovascular and renal outcomes in type 2 diabetes mellitus. *Circulation* 2019;139:2022–2031
263. Palmer SC, Tendal B, Mustafa RA, et al. Sodium-glucose cotransporter protein-2 (SGLT-2) inhibitors and glucagon-like peptide-1 (GLP-1) receptor agonists for type 2 diabetes: systematic review and network meta-analysis of randomised controlled trials. *BMJ* 2021;372:m4573
264. Zelniker TA, Wiviott SD, Raz I, et al. SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet* 2019;393:31–39
265. McGuire DK, Shih WJ, Cosentino F, et al. Association of SGLT2 inhibitors with cardiovascular and kidney outcomes in patients with type 2 diabetes: a meta-analysis. *JAMA Cardiol* 2021;6:148–158
266. Gerstein HC, Sattar N, Rosenstock J, et al.; AMPLITUDE-O Trial Investigators. Cardiovascular and renal outcomes with efpeglenatide in type 2 diabetes. *N Engl J Med* 2021;385:896–907
267. Lam CSP, Ramasundarahettige C, Branch KR, et al. Efpeglenatide and clinical outcomes with and without concomitant sodium-glucose cotransporter-2 inhibition use in type 2 diabetes: exploratory analysis of the AMPLITUDE-O trial. *Circulation* 2022;145:565–574
268. Del Gobbo LC, Kalantarian S, Imamura F, et al. Contribution of major lifestyle risk factors for incident heart failure in older adults: the

- Cardiovascular Health Study. *JACC Heart Fail* 2015;3:520–528
269. Young DR, Reynolds K, Sidell M, et al. Effects of physical activity and sedentary time on the risk of heart failure. *Circ Heart Fail* 2014;7:21–27
270. Tektonidis TG, Åkesson A, Gigante B, Wolk A, Larsson SC. Adherence to a Mediterranean diet is associated with reduced risk of heart failure in men. *Eur J Heart Fail* 2016;18:253–259
271. Levitan EB, Wolk A, Mittleman MA. Consistency with the DASH diet and incidence of heart failure. *Arch Intern Med* 2009;169:851–857
272. Levy D, Larson MG, Vasan RS, Kannel WB, Ho KK. The progression from hypertension to congestive heart failure. *JAMA* 1996;275:1557–1562
273. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998;317:703–713
274. Upadhyia B, Rocco M, Lewis CE, et al.; SPRINT Research Group. Effect of intensive blood pressure treatment on heart failure events in the systolic blood pressure reduction intervention trial. *Circ Heart Fail* 2017;10:e003613
275. Yusuf S, Pitt B, Davis CE, Hood WB Jr; SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med* 1992;327:685–691
276. Pfeffer MA, Braunwald E, Moyé LA, et al.; The SAVE Investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. Results of the Survival and Ventricular Enlargement trial. *N Engl J Med* 1992;327:669–677
277. Exner DV, Dries DL, Wacławiw MA, Shelton B, Domanski MJ. Beta-adrenergic blocking agent use and mortality in patients with asymptomatic and symptomatic left ventricular systolic dysfunction: a post hoc analysis of the Studies of Left Ventricular Dysfunction. *J Am Coll Cardiol* 1999;33:916–923
278. Vantrimpont P, Rouleau JL, Wun CC, et al.; SAVE Investigators. Additive beneficial effects of beta-blockers to angiotensin-converting enzyme inhibitors in the Survival and Ventricular Enlargement (SAVE) study. *J Am Coll Cardiol* 1997;29:229–236
279. Dargie HJ. Effect of carvedilol on outcome after myocardial infarction in patients with left-ventricular dysfunction: the CAPRICORN randomised trial. *Lancet* 2001;357:1385–1390
280. Colucci WS, Kolias TJ, Adams KF, et al.; REVERT Study Group. Metoprolol reverses left ventricular remodeling in patients with asymptomatic systolic dysfunction: the REversal of VEntricular Remodeling with Toprol-XL (REVERT) trial. *Circulation* 2007;116:49–56
281. Bhatt DL, Szarek M, Pitt B, et al.; SCORED Investigators. Sotagliflozin in patients with diabetes and chronic kidney disease. *N Engl J Med* 2021;384:129–139
282. Bakris GL, Agarwal R, Anker SD, et al.; FIDELIO-DKD Investigators. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *N Engl J Med* 2020;383:2219–2229
283. Dormandy JA, Charbonnel B, Eckland DJ, et al.; PROactive Investigators. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAzone Clinical Trial In macroVascular Events): a randomised controlled trial. *Lancet* 2005;366:1279–1289
284. Singh S, Loke YK, Furberg CD. Long-term risk of cardiovascular events with rosiglitazone: a meta-analysis. *JAMA* 2007;298:1189–1195
285. Lincoff AM, Wolski K, Nicholls SJ, Nissen SE. Pioglitazone and risk of cardiovascular events in patients with type 2 diabetes mellitus: a meta-analysis of randomized trials. *JAMA* 2007;298:1180–1188
286. Inzuchi SE, Masoudi FA, McGuire DK. Metformin in heart failure. *Diabetes Care* 2007;30:e129
287. Eurich DT, Majumdar SR, McAlister FA, Tsuyuki RT, Johnson JA. Improved clinical outcomes associated with metformin in patients with diabetes and heart failure. *Diabetes Care* 2005;28:2345–2351
288. U.S. Food and Drug Administration. FDA drug safety communication: FDA revises warnings regarding use of the diabetes medicine metformin in certain patients with reduced kidney function, 2016. Accessed 10 October 2023. Available from <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-revises-warnings-regarding-use-diabetes-medicine-metformin-certain>
289. Scirica BM, Bhatt DL, Braunwald E, et al.; SAVOR-TIMI 53 Steering Committee and Investigators. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013;369:1317–1326
290. Zannad F, Cannon CP, Cushman WC, et al.; EXAMINE Investigators. Heart failure and mortality outcomes in patients with type 2 diabetes taking alogliptin versus placebo in EXAMINE: a multicentre, randomised, double-blind trial. *Lancet* 2015;385:2067–2076
291. Green JB, Bethel MA, Armstrong PW, et al.; TECOS Study Group. Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015;373:232–242
292. Rosenstock J, Perkovic V, Johansen OE, et al.; CARMELINA Investigators. Effect of linagliptin vs placebo on major cardiovascular events in adults with type 2 diabetes and high cardiovascular and renal risk: the CARMELINA randomized clinical trial. *JAMA* 2019;321:69–79
293. Vaduganathan M, Docherty KF, Claggett BL, et al. SGLT-2 inhibitors in patients with heart failure: a comprehensive meta-analysis of five randomized controlled trials. *Lancet* 2022;400:757–767
294. Packer M, Anker SD, Butler J, et al. Effect of empagliflozin on the clinical stability of patients with heart failure and a reduced ejection fraction: the EMPEROR-Reduced trial. *Circulation* 2021;143:326–336
295. Voors AA, Angermann CE, Teerlink JR, et al. The SGLT2 inhibitor empagliflozin in patients hospitalized for acute heart failure: a multinational randomized trial. *Nat Med* 2022;28:568–574
296. Spertus JA, Birmingham MC, Nassif M, et al. The SGLT2 inhibitor canagliflozin in heart failure: the CHIEF-HF remote, patient-centered randomized trial. *Nat Med* 2022;28:809–813
297. Bhatt DL, Szarek M, Steg PG, et al.; SOLOIST-WHF Trial Investigators. Sotagliflozin in patients with diabetes and recent worsening heart failure. *N Engl J Med* 2021;384:117–128
298. Peters AL, Henry RR, Thakkar P, Tong C, Alba M. Diabetic ketoacidosis with canagliflozin, a sodium–glucose cotransporter 2 inhibitor, in patients with type 1 diabetes. *Diabetes Care* 2016;39:532–538
299. Danne T, Garg S, Peters AL, et al. International consensus on risk management of diabetic ketoacidosis in patients with type 1 diabetes treated with sodium–glucose cotransporter (SGLT) inhibitors. *Diabetes Care* 2019;42:1147–1154
300. Musso G, Sircana A, Saba F, Cassader M, Gambino R. Assessing the risk of ketoacidosis due to sodium–glucose cotransporter (SGLT)-2 inhibitors in patients with type 1 diabetes: a meta-analysis and meta-regression. *PLoS Med* 2020;17:e1003461
301. Holt RIG, DeVries JH, Hess-Fischl A, et al. The management of type 1 diabetes in adults. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2021;44:2589–2625
302. Rosenstock J, Marquardt J, Laffel LM, et al. Empagliflozin as adjunctive to insulin therapy in type 1 diabetes: the EASE trials. *Diabetes Care* 2018;41:2560–2569
303. Mathieu C, Dandona P, Gillard P, et al.; DEPICT-2 Investigators. Efficacy and safety of dapagliflozin in patients with inadequately controlled type 1 diabetes (the DEPICT-2 study): 24-week results from a randomized controlled trial. *Diabetes Care* 2018;41:1938–1946
304. Garg SK, Henry RR, Banks P, et al. Effects of sotagliflozin added to insulin in patients with type 1 diabetes. *N Engl J Med* 2017;377:2337–2348
305. Rodbard HW, Peters AL, Slee A, Cao A, Traina SB, Alba M. The effect of canagliflozin, a sodium–glucose cotransporter 2 inhibitor, on glycemic end points assessed by continuous glucose monitoring and patient-reported outcomes among people with type 1 diabetes. *Diabetes Care* 2017;40:171–180
306. Palanca A, van Nes F, Pardo F, Ampudia Blasco FJ, Mathieu C. Real-world evidence of efficacy and safety of SGLT2 inhibitors as adjunctive therapy in adults with type 1 diabetes: a European two-center experience. *Diabetes Care* 2022;45:650–658
307. U.S. Food and Drug Administration. Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee. 2019. Accessed 10 August 2023. Available from <https://wayback.archive-it.org/7993/20190207212714/https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/EndocrinologicAndMetabolicDrugsAdvisoryCommittee/UCM629782.pdf>
308. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013;382:339–352
309. Das SR, Everett BM, Birtcher KK, et al. 2020 Expert consensus decision pathway on novel therapies for cardiovascular risk reduction in patients with type 2 diabetes: a report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol* 2020;76:1117–1145
310. Zoungas S, Chalmers J, Neal B, et al.; ADVANCE-ON Collaborative Group. Follow-up of blood-pressure lowering and glucose control in type 2 diabetes. *N Engl J Med* 2014;371:1392–1406

311. White WB, Cannon CP, Heller SR, et al.; EXAMINE Investigators. Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med* 2013;369:1327–1335
312. Rosenstock J, Perkovic V, Alexander JH, et al.; CARMELINA Investigators. Rationale, design, and baseline characteristics of the CArdiovascular safety and Renal Microvascular outcomeE study with LINagliptin (CARMELINA): a randomized, double-blind, placebo-controlled clinical trial in patients with type 2 diabetes and high cardio-renal risk. *Cardiovasc Diabetol* 2018;17:39
313. Cefalu WT, Kaul S, Gerstein HC, et al. Cardiovascular outcomes trials in type 2 diabetes: where do we go from here? Reflections from a *Diabetes Care* editors' expert forum. *Diabetes Care* 2018; 41:14–31
314. Wheeler DC, Stefansson BV, Batiushin M, et al. The Dapagliflozin and Prevention of Adverse Outcomes in Chronic Kidney Disease (DAPA-CKD) trial: baseline characteristics. *Nephrol Dial Transplant* 2020;35:1700–1711
315. Cannon CP, McGuire DK, Pratley R, et al.; VERTIS-CV Investigators. Design and baseline characteristics of the eValuation of ERiTugliflozin efflacy and Safety CardioVascular outcomes trial (VERTIS-CV). *Am Heart J* 2018; 206:11–23
316. Anker SD, Butler J, Filippatos G, et al.; EMPEROR-Preserved Trial Committees and Investigators. Baseline characteristics of patients with heart failure with preserved ejection fraction in the EMPEROR-Preserved trial. *Eur J Heart Fail* 2020;22:2383–2392