

Quantum Field Theory

Lecture Notes

Quantization, Fock space, propagators, scattering, and renormalization.



Contents

Preface: What These Notes Are For	1
1 From Classical Normal Modes to Quantum Oscillators	3
1.1 Starting Point: A Classical Field Decomposed into Modes	3
1.2 Each Mode as a Harmonic Oscillator	4
1.3 Quantizing a Single Oscillator	4
1.4 Creation and Annihilation Operators	5
1.5 Number Operator and Energy Spectrum	5
1.6 The Vacuum State of One Oscillator	5
1.7 From One Oscillator to Infinitely Many Oscillators	6
1.8 Field Quanta as Mode Excitations	6
1.9 What Is New Compared with Classical Field Theory?	7
2 Canonical Quantization of the Real Scalar Field	8
2.1 Planned Sections	8
3 The Complex Scalar Field and Antiparticles	9
3.1 Planned Sections	9
4 Quantization of the Dirac Field	10
4.1 Planned Sections	10
5 Quantization of the Electromagnetic Field	11
5.1 Planned Sections	11
6 Propagators as Quantum Correlation Functions	12
6.1 Planned Sections	12
7 The Interaction Picture	13
7.1 Planned Sections	13
8 Wick Theorem and Diagrammatics	14
8.1 Planned Sections	14
9 Scattering Theory and the S-Matrix	15
9.1 Planned Sections	15
10 Path Integral for Quantum Mechanics	16
10.1 Planned Sections	16
11 Path Integral for Scalar Fields	17
11.1 Planned Sections	17
12 Fermionic Path Integrals	18
12.1 Planned Sections	18
13 Gauge Fields in the Path Integral	19
13.1 Planned Sections	19

14 ϕ^4 Theory as the First Interacting QFT	20
14.1 Planned Role	20
15 Yukawa Theory	21
15.1 Planned Role	21
16 Quantum Electrodynamics	22
16.1 Planned Role	22
17 Loop Effects in QED	23
17.1 Planned Role	23
18 Regularization	24
18.1 Planned Role	24
19 Renormalization	25
19.1 Planned Role	25
20 Renormalization Group	26
20.1 Planned Role	26
21 Effective Field Theory	27
21.1 Planned Role	27
22 Non-Abelian Gauge Theory as a Quantum Theory	28
22.1 Planned Role	28
23 BRST Symmetry	29
23.1 Planned Role	29
24 Quantum Chromodynamics	30
24.1 Planned Role	30
25 Symmetries in Quantum Field Theory	31
25.1 Planned Role	31
26 Anomalies	32
26.1 Planned Role	32
27 Vacuum Structure and Nonperturbative Effects	33
27.1 Planned Role	33
28 The Quantum Electroweak Theory	34
28.1 Planned Role	34
29 The Standard Model Lagrangian as a Quantum Theory	35
29.1 Planned Role	35
30 Selected Standard Model Processes	36
30.1 Planned Role	36
31 Euclidean Quantum Field Theory	37
31.1 Planned Role	37

32 Lattice Field Theory	38
32.1 Planned Role	38
33 Large-N and Semiclassical Methods	39
33.1 Planned Role	39
34 Conceptual Synthesis I	40
34.1 Planned Role	40
35 Conceptual Synthesis II	41
35.1 Planned Role	41
36 Summary: The Architecture of Quantum Field Theory	42
36.1 Planned Role	42
A Quantization Dictionary	43
A.1 Classical Mode to Quantum Oscillator	43
A.2 Poisson Bracket to Commutator	43
A.3 Classical Field to Field Operator	43
A.4 Classical Green's Function to Propagator	43
A.5 Classical Symmetry to Quantum Ward Identity	44
B Reference B	45
C Reference C	46
D Reference D	47
E Reference E	48

Preface: What These Notes Are For

Remark

These notes are designed as a continuation of the Classical Field Theory I–II sequence. They do not try to re-teach variational calculus, Lorentz covariance, Noether currents, gauge redundancy, or the classical equations of fundamental fields from the beginning. Instead, they start where the classical notes stop: with fields as dynamical systems ready to be quantized.

Quantum Field Theory After Classical Field Theory

Classical field theory teaches us to describe nature by local fields, actions, symmetries, conservation laws, and equations of motion. Quantum field theory keeps this architecture, but changes the meaning of the basic objects. A classical field configuration is no longer the final description of a physical state. It becomes an ingredient in a quantum theory whose observable content is encoded in states, operators, correlation functions, transition amplitudes, and probabilities.

The guiding idea of the first part of these notes is simple: a free classical field decomposes into normal modes, and each normal mode behaves like a harmonic oscillator. Quantizing the field therefore means quantizing infinitely many oscillators at once. The excitations of these oscillators are interpreted as particles.

What Will Not Be Repeated

The notes assume familiarity with the following ideas from classical field theory:

- the action principle and Euler–Lagrange equations for fields,
- canonical momenta and Hamiltonian density,
- Lorentz covariance and relativistic notation,
- real and complex scalar fields,
- electromagnetic and Yang–Mills gauge fields at the classical level,
- Noether’s theorem and conserved currents,
- Green’s functions as solutions of linear differential equations.

Whenever one of these topics is used in a quantum calculation, the notes recall just enough of the classical structure to make the calculation readable. The aim is not repetition, but conversion: classical structures are translated into quantum structures.

What Is New in QFT

The genuinely new concepts are the vacuum, Fock space, field operators, propagators, Wick contractions, Feynman diagrams, scattering amplitudes, loop corrections, regularization, renormalization, and scale dependence. These ideas are not decorative additions to classical field theory. They are the mechanisms that make quantum field theory predictive.

A central conceptual warning will appear repeatedly: a Feynman diagram is not a literal picture of tiny particles following tiny trajectories. It is a compact bookkeeping device for terms in a perturbative expansion of correlation functions or scattering amplitudes.

How to Read These Notes

A useful rhythm for reading a QFT calculation is:

classical field $\longrightarrow$ mode expansion $\longrightarrow$ operators $\longrightarrow$ states $\longrightarrow$ correlators $\longrightarrow$ amplitudes.

At first this may look like a change of notation. It is not. Each arrow changes what is meant by a physical prediction. The purpose of the notes is to make those changes explicit and technically usable.

Chapter 1

From Classical Normal Modes to Quantum Oscillators

Remark

This chapter follows the structure of the course plan in `ideas_Quantum_Field_Theory.md`. Its purpose is to turn the first entry of the table of contents into a readable lecture: a free classical field is decomposed into normal modes, each mode is recognized as a harmonic oscillator, and the oscillator excitations become the first particle states of quantum field theory.

1.1 Starting Point: A Classical Field Decomposed into Modes

Classical field theory describes a field by a function such as $\phi(t, \mathbf{x})$. For a free field, this function can be decomposed into normal modes. This is the same idea that appears in mechanics when a system of coupled oscillators is rewritten in terms of independent normal coordinates.

Begin with a finite chain of N identical masses connected by springs. If $q_j(t)$ is the displacement of the j -th mass, a typical quadratic Lagrangian is

$$L = \sum_{j=1}^N \frac{m}{2} \dot{q}_j^2 - \sum_{j=1}^N \frac{\kappa}{2} (q_{j+1} - q_j)^2.$$

The variables q_j are coupled. After introducing normal coordinates $Q_r(t)$, one obtains the diagonal form

$$L = \sum_r \left(\frac{1}{2} \dot{Q}_r^2 - \frac{1}{2} \omega_r^2 Q_r^2 \right).$$

The coupled chain is therefore equivalent to a collection of independent oscillators.

A field is the continuum version of this situation. In a one-dimensional chain with spacing a , one may view

$$q_j(t) \approx \phi(t, x_j), \quad x_j = ja.$$

In the continuum limit, sums over sites become integrals over space, and the normal-mode decomposition becomes a Fourier decomposition.

For a real scalar field in a box of volume V , with periodic boundary conditions, write schematically

$$\phi(t, \mathbf{x}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} q_{\mathbf{k}}(t) e^{i\mathbf{k} \cdot \mathbf{x}},$$

with the appropriate reality condition for a real field. The allowed momenta are discrete because the field is placed in a finite box. This is a temporary technical convenience; the continuum limit is obtained later by replacing sums with integrals.

Remark

The field is infinite-dimensional, but the free field is not mysterious at this stage. The normal-mode decomposition rewrites it as a collection of independent oscillating degrees

of freedom.

1.2 Each Mode as a Harmonic Oscillator

For the free real scalar field, the classical Lagrangian is

$$L = \int d^3x \left(\frac{1}{2} \dot{\phi}^2 - \frac{1}{2} (\nabla \phi)^2 - \frac{1}{2} m^2 \phi^2 \right).$$

Its equation of motion is the Klein–Gordon equation,

$$(\partial_t^2 - \nabla^2 + m^2)\phi(t, \mathbf{x}) = 0.$$

After inserting the normal-mode expansion into the free Lagrangian and choosing standard normalizations, the field decomposes into independent oscillator modes,

$$L = \sum_{\mathbf{k}} \left(\frac{1}{2} \dot{q}_{\mathbf{k}}^2 - \frac{1}{2} \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2 \right),$$

where

$$\omega_{\mathbf{k}} = \sqrt{\mathbf{k}^2 + m^2}$$

in units with $c = \hbar = 1$.

The canonical momentum of the mode $q_{\mathbf{k}}$ is

$$p_{\mathbf{k}} = \dot{q}_{\mathbf{k}},$$

and the Hamiltonian becomes

$$H = \sum_{\mathbf{k}} H_{\mathbf{k}}, \quad H_{\mathbf{k}} = \frac{1}{2} p_{\mathbf{k}}^2 + \frac{1}{2} \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2.$$

Thus each mode $\mathbf{k}$ is a harmonic oscillator of frequency $\omega_{\mathbf{k}}$.

Example

The classical field is not quantized all at once. First, the free classical field is diagonalized into modes. Then each mode is quantized using the same oscillator algebra known from elementary quantum mechanics.

1.3 Quantizing a Single Oscillator

For one harmonic oscillator, the classical variables q and p become operators $\hat{q}$ and $\hat{p}$ satisfying

$$[\hat{q}, \hat{p}] = i.$$

The Hamiltonian is

$$\hat{H} = \frac{1}{2} \hat{p}^2 + \frac{1}{2} \omega^2 \hat{q}^2.$$

Introduce the annihilation and creation operators

$$\hat{a} = \sqrt{\frac{\omega}{2}} \hat{q} + \frac{i}{\sqrt{2\omega}} \hat{p}, \quad \hat{a}^\dagger = \sqrt{\frac{\omega}{2}} \hat{q} - \frac{i}{\sqrt{2\omega}} \hat{p}.$$

Equivalently,

$$\hat{q} = \frac{1}{\sqrt{2\omega}} (\hat{a} + \hat{a}^\dagger), \quad \hat{p} = -i\sqrt{\frac{\omega}{2}} (\hat{a} - \hat{a}^\dagger).$$

The canonical commutator implies

$$[\hat{a}, \hat{a}^\dagger] = 1.$$

The Hamiltonian becomes

$$\hat{H} = \omega \left(\hat{a}^\dagger \hat{a} + \frac{1}{2} \right).$$

This is the algebraic form that will be copied for every field mode.

1.4 Creation and Annihilation Operators

The operator $\hat{a}^\dagger$ raises the excitation number of the oscillator, while $\hat{a}$ lowers it. If $|0\rangle$ is the oscillator ground state, then

$$\hat{a}|0\rangle = 0.$$

Repeated action of $\hat{a}^\dagger$ produces excited states,

$$|n\rangle = \frac{(\hat{a}^\dagger)^n}{\sqrt{n!}}|0\rangle, \quad n = 0, 1, 2, \dots$$

In a single oscillator problem, this is a convenient way to organize the energy spectrum. In a field theory, the same algebra acquires a new interpretation: creation operators create field quanta.

1.5 Number Operator and Energy Spectrum

The number operator is

$$\hat{N} = \hat{a}^\dagger \hat{a}.$$

It satisfies

$$\hat{N}|n\rangle = n|n\rangle.$$

The Hamiltonian eigenvalue equation is

$$\hat{H}|n\rangle = \omega \left(n + \frac{1}{2} \right) |n\rangle.$$

The integer n counts oscillator excitations. The term $\omega/2$ is the zero-point energy.

For a field, there will be one number operator for each mode,

$$\hat{N}_{\mathbf{k}} = \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}.$$

The eigenvalue of $\hat{N}_{\mathbf{k}}$ counts how many quanta occupy the mode $\mathbf{k}$.

1.6 The Vacuum State of One Oscillator

The oscillator vacuum is defined algebraically by

$$\hat{a}|0\rangle = 0.$$

It is not the state of zero energy; its energy is

$$E_0 = \frac{1}{2}\omega.$$

This small fact becomes conceptually important in field theory. A free field has one oscillator for every allowed momentum mode, so the formal vacuum energy is

$$E_{\text{vac}} = \frac{1}{2} \sum_{\mathbf{k}} \omega_{\mathbf{k}}.$$

In the continuum limit this expression diverges. Later chapters will explain normal ordering, regularization, renormalization, and the physical meaning of vacuum energy. At this stage the important lesson is simpler: the quantum vacuum is a state with no excitations, not a state with no structure.

1.7 From One Oscillator to Infinitely Many Oscillators

For the scalar field, each normal mode receives its own creation and annihilation operators,

$$\hat{a}_{\mathbf{k}}, \quad \hat{a}_{\mathbf{k}}^\dagger.$$

They satisfy

$$[\hat{a}_{\mathbf{k}}, \hat{a}_{\mathbf{k}'}^\dagger] = \delta_{\mathbf{k}, \mathbf{k}'}, \quad [\hat{a}_{\mathbf{k}}, \hat{a}_{\mathbf{k}'}] = 0, \quad [\hat{a}_{\mathbf{k}}^\dagger, \hat{a}_{\mathbf{k}'}^\dagger] = 0.$$

The free-field vacuum is the state annihilated by every annihilation operator,

$$\hat{a}_{\mathbf{k}}|0\rangle = 0 \quad \text{for every } \mathbf{k}.$$

A one-particle state of momentum $\mathbf{k}$ is

$$|\mathbf{k}\rangle = \hat{a}_{\mathbf{k}}^\dagger|0\rangle.$$

A two-particle state may be written as

$$|\mathbf{k}_1, \mathbf{k}_2\rangle = \hat{a}_{\mathbf{k}_1}^\dagger \hat{a}_{\mathbf{k}_2}^\dagger|0\rangle,$$

up to normalization conventions.

Definition

The *Fock space* of a free quantum field is the Hilbert space generated from the vacuum by applying creation operators for all allowed modes. It is the natural state space for a theory in which the number of particles is not fixed.

1.8 Field Quanta as Mode Excitations

In the free theory, the Hamiltonian can be written as

$$\hat{H} = \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left(\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \frac{1}{2} \right).$$

The first term counts excitations of each mode. The state $\hat{a}_{\mathbf{k}}^\dagger|0\rangle$ has energy $\omega_{\mathbf{k}}$ above the free vacuum and momentum $\mathbf{k}$. It is interpreted as a one-particle state.

The field operator itself is reconstructed from the same mode operators. In a standard continuum normalization,

$$\hat{\phi}(t, \mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{k}}}} \left(\hat{a}_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}t + i\mathbf{k}\cdot\mathbf{x}} + \hat{a}_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}t - i\mathbf{k}\cdot\mathbf{x}} \right).$$

The coefficients of the classical mode expansion have become operators. The field operator can annihilate and create quanta because it contains both $\hat{a}_{\mathbf{k}}$ and $\hat{a}_{\mathbf{k}}^\dagger$.

Remark

A field quantum is not a tiny localized lump of classical field. In the free theory, it is a mode excitation created by a creation operator. Localization, wave packets, interactions, and particle detection require additional steps.

1.9 What Is New Compared with Classical Field Theory?

Classical field theory assigns numbers to field configurations. Quantum field theory assigns operators to fields and vectors to states. The basic object is no longer a definite field profile $\phi(x)$, but an operator $\hat{\phi}(x)$ acting on a Hilbert space.

The free quantum field still satisfies the free equation of motion,

$$(\square + m^2)\hat{\phi}(x) = 0,$$

but the meaning has changed. This is now an equation for an operator-valued distribution, not for a classical function.

The conceptual dictionary of the chapter is:

classical normal mode	→	quantum oscillator,
oscillator excitation	→	field quantum,
many oscillator excitations	→	Fock-space state,
classical field	→	field operator.

Summary

A free classical field can be decomposed into independent normal modes. Each mode is a harmonic oscillator. Quantizing all of these oscillators produces creation and annihilation operators, a vacuum state, number operators, and Fock space. In the free theory, particles are mode excitations of quantum fields. This is the starting point for propagators, perturbation theory, and scattering.

Chapter 2

Canonical Quantization of the Real Scalar Field

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

2.1 Planned Sections

- Equal-time commutation relations.
- Field operator and momentum operator.
- Mode expansion of the field operator.
- Creation and annihilation operators for momentum modes.
- Fock space.
- Vacuum, one-particle states, and many-particle states.
- Hamiltonian and momentum operators.
- Normal ordering.
- Zero-point energy and its interpretation.
- Field operators as operator-valued distributions.
- Causality and spacelike commutators.

Chapter 3

The Complex Scalar Field and Antiparticles

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

3.1 Planned Sections

- Quantizing the complex scalar field.
- Two independent sets of ladder operators.
- Particles and antiparticles.
- Charge operator.
- Positive and negative frequency modes revisited.
- Why antiparticles appear naturally.
- Complex scalar propagator.
- Charged quanta and global symmetry.
- Physical interpretation of the Fock space.

Chapter 4

Quantization of the Dirac Field

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

4.1 Planned Sections

- Why fermionic quantization is different.
- Mode expansion of the Dirac field.
- Anticommutation relations.
- Spinor creation and annihilation operators.
- Electron and positron states.
- Charge, momentum, and Hamiltonian operators.
- Positivity of energy and the role of anticommutators.
- Dirac propagator.
- Spin sums and completeness relations.
- Chirality, helicity, and massless fermions in QFT.
- Majorana quantization: conceptual overview.

Chapter 5

Quantization of the Electromagnetic Field

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

5.1 Planned Sections

- Why gauge fields are harder to quantize.
- Physical and redundant degrees of freedom.
- Coulomb gauge quantization.
- Photon creation and annihilation operators.
- Polarization vectors.
- Photon states and helicity.
- Covariant quantization.
- Gauge fixing and the photon propagator.
- Unphysical polarizations and their removal.
- Gauge-invariant observables.
- What is a photon?

Chapter 6

Propagators as Quantum Correlation Functions

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

6.1 Planned Sections

- From classical Green's functions to quantum two-point functions.
- Wightman functions.
- Time-ordered products.
- Feynman propagator.
- Retarded and advanced propagators in QFT.
- The $i\epsilon$ prescription.
- Momentum-space propagators.
- Pole structure and particle interpretation.
- Propagation vs correlation.
- What a propagator actually means.

Chapter 7

The Interaction Picture

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

7.1 Planned Sections

- Schrödinger, Heisenberg, and interaction pictures.
- Free fields as the starting point.
- Interaction Hamiltonian.
- Time evolution operator.
- Dyson series.
- Time-ordered exponential.
- Perturbation theory as an expansion in couplings.
- Vacuum persistence amplitude.
- Adiabatic switching and the meaning of in/out states.
- Why exact interacting QFT is difficult.

Chapter 8

Wick Theorem and Diagrammatics

Remark

This chapter follows the planned QFT table of contents. It is a placeholder to be developed in small increments.

8.1 Planned Sections

- Normal ordering revisited.
- Contractions.
- Wick theorem for scalar fields.
- Wick theorem for fermions.
- Fermionic sign factors.
- From algebra to diagrams.
- External lines, internal lines, and vertices.
- Connected and disconnected diagrams.
- Vacuum bubbles.
- Symmetry factors.
- Interpretation limits of Feynman diagrams.

Chapter 9

Scattering Theory and the S -Matrix

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

9.1 Planned Sections

- In-states and out-states.
- The S -matrix.
- Transition amplitudes.
- Matrix elements and momentum conservation.
- Cross sections.
- Decay rates.
- Lorentz-invariant phase space.
- The LSZ reduction formula.
- From correlation functions to scattering amplitudes.
- Tree-level $2 \rightarrow 2$ scattering.
- Decay of an unstable particle.
- From amplitudes to measurable predictions.

Chapter 10

Path Integral for Quantum Mechanics

Remark

This chapter is reserved by the course plan in `ideas_Quantum_Field_Theory.md`. It will be developed in small increments.

10.1 Planned Sections

- Transition amplitudes as sums over histories.
- Time slicing.
- Classical limit and stationary phase.
- Gaussian integrals.
- Generating functionals in ordinary quantum mechanics.
- Sources and correlation functions.
- Why the path integral is useful for QFT.
- Relation to canonical quantization.

Chapter 11

Path Integral for Scalar Fields

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

11.1 Planned Sections

- Functional integration.
- Free scalar generating functional.
- Sources and derivatives.
- Correlation functions.
- Gaussian functional integrals.
- Perturbation theory.
- Diagrammatic expansion.
- Connected functional.
- Effective action.
- Classical field as expectation value.
- One-particle-irreducible functions.

Chapter 12

Fermionic Path Integrals

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

12.1 Planned Sections

- Grassmann variables.
- Berezin integration.
- Gaussian integrals for Grassmann variables.
- Fermionic generating functional.
- Dirac propagator from the path integral.
- Fermion loops and minus signs.
- Determinants from integrating out fermions.
- Majorana fermions in the path integral.
- Physical interpretation and common pitfalls.

Chapter 13

Gauge Fields in the Path Integral

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

13.1 Planned Sections

- Overcounting gauge-equivalent configurations.
- Gauge fixing in the functional integral.
- Faddeev–Popov method.
- Ghost fields.
- Why ghosts are not physical particles.
- Gauge-fixed photon path integral.
- Gauge-fixed Yang–Mills path integral.
- First introduction to BRST symmetry.
- Gauge independence of physical observables.
- Meaning of gauge fixing in quantum theory.

Chapter 14

ϕ^4 Theory as the First Interacting QFT

Remark

Placeholder chapter from the QFT course plan in `ideas_Quantum_Field_Theory.md`.
Detailed lecture text will be added later.

14.1 Planned Role

This chapter will introduce the first interacting scalar quantum field theory and the basic structure of perturbative Feynman rules, loop corrections, counterterms, and renormalized parameters.

Chapter 15

Yukawa Theory

Remark

Placeholder chapter from the QFT course plan in `ideas_Quantum_Field_Theory.md`.
Detailed lecture text will be added later.

15.1 Planned Role

This chapter will study a scalar field coupled to fermions, Yukawa vertices, simple scattering and decay processes, and the low-energy interpretation of scalar exchange.

Chapter 16

Quantum Electrodynamics

Remark

Placeholder chapter from the QFT course plan in `ideas_Quantum_Field_Theory.md`.
Detailed lecture text will be added later.

16.1 Planned Role

This chapter will introduce QED as the quantum theory of charged matter and light, including Feynman rules, spinor chains, Ward identities, and standard tree-level processes.

Chapter 17

Loop Effects in QED

Remark

Placeholder chapter from the QFT course plan in `ideas_Quantum_Field_Theory.md`.
Detailed lecture text will be added later.

17.1 Planned Role

This chapter will cover radiative corrections in QED, including self-energies, vacuum polarization, vertex corrections, charge renormalization, infrared divergences, and inclusive observables.

Chapter 18

Regularization

Remark

Placeholder chapter from the QFT course plan in `ideas_Quantum_Field_Theory.md`.
Detailed lecture text will be added later.

18.1 Planned Role

This chapter will explain why divergent loop integrals appear and how cutoff, Pauli–Villars, dimensional regularization, and subtraction schemes organize intermediate infinities.

Chapter 19

Renormalization

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

19.1 Planned Role

This chapter will develop the standard QFT treatment of renormalized masses, fields, couplings, counterterms, and physical predictions.

Chapter 20

Renormalization Group

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

20.1 Planned Role

This chapter will organize the scale dependence of quantum field theories and introduce running couplings, beta functions, fixed points, and effective descriptions.

Chapter 21

Effective Field Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

21.1 Planned Role

This chapter will introduce effective field theory, separation of scales, matching, power counting, symmetries, operator bases, decoupling, and naturalness.

Chapter 22

Non-Abelian Gauge Theory as a Quantum Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

22.1 Planned Role

This chapter will develop quantum non-Abelian gauge fields, self-interactions, gauge-fixed Feynman rules, ghosts, Ward-type identities, and unitarity checks.

Chapter 23

BRST Symmetry

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

23.1 Planned Role

This chapter will introduce BRST transformations, ghost number, nilpotency, physical states as cohomology, gauge independence, and Slavnov–Taylor identities.

Chapter 24

Quantum Chromodynamics

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

24.1 Planned Role

This chapter will introduce QCD field content, color charge, QCD Feynman rules, gluon self-interactions, asymptotic freedom, confinement, jets, and perturbative versus nonperturbative regimes.

Chapter 25

Symmetries in Quantum Field Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

25.1 Planned Role

This chapter will translate classical symmetries into quantum operators, conserved charges, Ward identities, current correlation functions, selection rules, and constraints on observables.

Chapter 26

Anomalies

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

26.1 Planned Role

This chapter will study symmetries that fail after quantization, chiral currents, triangle diagrams, path-integral measure effects, gauge anomalies, anomaly cancellation, and physical consequences.

Chapter 27

Vacuum Structure and Nonperturbative Effects

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

27.1 Planned Role

This chapter will discuss the quantum vacuum, degenerate vacua, tunneling, instantons, theta vacua, false vacuum decay, semiclassical approximation, and nonperturbative contributions.

Chapter 28

The Quantum Electroweak Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

28.1 Planned Role

This chapter will introduce the quantum electroweak sector, gauge fixing after symmetry breaking, massive vector boson propagators, currents, weak processes, and the low-energy Fermi theory.

Chapter 29

The Standard Model Lagrangian as a Quantum Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

29.1 Planned Role

This chapter will organize the Standard Model field content, representations, gauge interactions, Yukawa sector, Higgs sector, flavor mixing, neutrino masses, anomaly cancellation, and EFT viewpoint.

Chapter 30

Selected Standard Model Processes

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

30.1 Planned Role

This chapter will connect QFT calculations with representative Standard Model predictions and collider observables.

Chapter 31

Euclidean Quantum Field Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

31.1 Planned Role

This chapter will introduce Wick rotation, Euclidean correlation functions, reflection positivity, statistical field theory connections, finite temperature, boundary conditions, and Matsubara frequencies.

Chapter 32

Lattice Field Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

32.1 Planned Role

This chapter will introduce lattice regularization, scalar fields on a lattice, gauge fields on links, Wilson action, Wilson loops, fermion difficulties, Monte Carlo sampling, and the continuum limit.

Chapter 33

Large- N and Semiclassical Methods

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

33.1 Planned Role

This chapter will introduce large- N counting, planar diagrams, saddle-point methods, semiclassical expansion, collective fields, solitons, instanton calculus, and reliability of approximations.

Chapter 34

Conceptual Synthesis I

34.1 Planned Role

Placeholder chapter from the QFT course plan.

Chapter 35

Conceptual Synthesis II

35.1 Planned Role

Placeholder chapter from the QFT course plan.

Chapter 36

Summary: The Architecture of Quantum Field Theory

Remark

Placeholder chapter from the QFT course plan. Detailed lecture text will be added later.

36.1 Planned Role

This chapter will summarize quantization, Fock space, propagators, interactions, renormalization, gauge theory, anomalies, nonperturbative effects, the Standard Model, and open directions.

Appendix A

Quantization Dictionary

Remark

This appendix records the translation rules that are used repeatedly in the main text. They should not be treated as mechanical replacements valid in every context. They are starting points that must be checked against constraints, symmetries, domains of operators, and gauge redundancies.

A.1 Classical Mode to Quantum Oscillator

A classical normal mode with amplitude $q_{\mathbf{k}}(t)$ and frequency $\omega_{\mathbf{k}}$ is quantized like a harmonic oscillator. Schematically,

$$q_{\mathbf{k}}, p_{\mathbf{k}} \longrightarrow \hat{q}_{\mathbf{k}}, \hat{p}_{\mathbf{k}}, \quad [\hat{q}_{\mathbf{k}}, \hat{p}_{\mathbf{k}'}] = i\delta_{\mathbf{k}\mathbf{k}'}.$$

The oscillator is then rewritten in terms of ladder operators $\hat{a}_{\mathbf{k}}$ and $\hat{a}_{\mathbf{k}}^\dagger$.

A.2 Poisson Bracket to Commutator

The canonical rule is

$$\{A, B\}_{\text{PB}} \longrightarrow \frac{1}{i}[\hat{A}, \hat{B}].$$

For fields, the equal-time canonical relation is

$$[\hat{\phi}(t, \mathbf{x}), \hat{\pi}(t, \mathbf{y})] = i\delta^{(3)}(\mathbf{x} - \mathbf{y}).$$

A.3 Classical Field to Field Operator

A classical field $\phi(x)$ becomes an operator-valued distribution $\hat{\phi}(x)$. The word “distribution” is important: products of field operators at the same spacetime point are generally singular and require a careful definition.

A.4 Classical Green’s Function to Propagator

A classical Green’s function solves a differential equation with a delta-function source. A quantum propagator is a vacuum expectation value of a time-ordered product,

$$\Delta_F(x - y) = \langle 0|T\{\hat{\phi}(x)\hat{\phi}(y)\}|0\rangle.$$

It is related to Green’s functions, but its physical meaning is quantum correlation, not merely classical signal propagation.

A.5 Classical Symmetry to Quantum Ward Identity

A classical continuous symmetry gives a conserved current. In the quantum theory, the same symmetry constrains correlation functions and scattering amplitudes through Ward identities. If the symmetry fails after quantization, the failure is called an anomaly.

Appendix B

Reference B

Placeholder appendix.

Appendix C

Reference C

Placeholder appendix.

Appendix D

Reference D

Placeholder appendix.

Appendix E

Reference E

Placeholder appendix.

List of Figures