

# **Set Theory**

**CSE-103: Discrete Mathematics**

# Introduction to Set Theory

- A *set* is a structure, representing an unordered collection (group, plurality) of zero or more distinct (different) objects.
- Set theory deals with operations between, relations among, and statements about sets.
- The *objects* in a set are called the elements or members of that set.

# Basic notations for sets

- For sets, we'll use variables  $S$ ,  $T$ ,  $U$ , ...
- We can denote a set  $S$  in writing by listing all of its elements in curly braces:
  - $\{a, b, c\}$  is the set of whatever 3 objects are denoted by  $a$ ,  $b$ ,  $c$ .
- *Set builder notation*: For any proposition  $P(x)$  over any universe of discourse,  $\{x|P(x)\}$  is *the set of all  $x$  such that  $P(x)$* .
  - e.g.,  $\{x \mid x \text{ is an integer where } x>0 \text{ and } x<5 \}$

# Basic properties of sets

- Sets are inherently unordered:
  - No matter what objects  $a$ ,  $b$ , and  $c$  denote,  
 $\{a, b, c\} = \{a, c, b\} = \{b, a, c\} =$   
 $\{b, c, a\} = \{c, a, b\} = \{c, b, a\}.$
- All elements are distinct (unequal); multiple listings make no difference!
  - $\{a, b, c\} = \{a, a, b, a, b, c, c, c, c\}.$
  - This set contains at most 3 elements!

# Definition of Set Equality


- Two sets are declared to be equal *if and only if* they contain exactly the same elements.
  - Two sets A and B are equal if and only if  $\forall x(x \in A \leftrightarrow x \in B)$   
i.e.  $A=B$ .
- In particular, it does not matter *how the set is defined or denoted*.
- For example: The set  $\{1, 2, 3, 4\} =$   
 $\{x \mid x \text{ is an integer where } x > 0 \text{ and } x < 5\} =$   
 $\{x \mid x \text{ is a positive integer whose square}$   
 $\text{is } > 0 \text{ and } < 25\}$

# Infinite Sets

- Conceptually, sets may be *infinite* (i.e., not *finite*, without end, unending).
- Symbols for some special infinite sets:
  - $\mathbf{N} = \{0, 1, 2, 3, \dots\}$ , The natural numbers
  - $\mathbf{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$ , The integers
  - $\mathbf{Z}^+ = \{1, 2, 3, \dots\}$ , The positive integers
  - $\mathbf{Q} = \{p/q \mid p \in \mathbf{Z}, q \in \mathbf{Z}, \text{ and } q \neq 0\}$ , The rational numbers
  - $\mathbf{R}$ , the set of real numbers such as:  
374.1828471929498181917281943125...
  - $\mathbf{R}^+$ , The positive real numbers
  - $\mathbf{C}$ , The of complex numbers.

# Venn Diagram

- Used to represent sets graphically.
  - Universal set consist of all elements of interest. Then the set is drawn as a circle or other geometrical shape with its all members within the rectangle.
  - Sometimes points are used to represent elements of the set.




# Venn Diagrams

Primes  $< 10$

Even integers from 2 to 9

Odd integers from 1 to 9



Positive integers less than 10

\* Integers from -1 to 9



# Basic Set Relations: Member of

- $x \in S$  (“ $x$  is in  $S$ ”) is the proposition that object  $x$  is an *element* or *member* of set  $S$ .
  - e.g.  $3 \in \mathbf{N}$ , “ $a$ ”  $\in \{x \mid x \text{ is a letter of the alphabet}\}$
- Can define set equality in terms of  $\in$  relation:  
 $\forall S, T: S = T \leftrightarrow (\forall x: x \in S \leftrightarrow x \in T)$   
“Two sets are equal **iff** they have all the same members.”
- $x \notin S \equiv \neg(x \in S)$       “ $x$  is not in  $S$ ”

# The Empty Set

- $\emptyset$  (“null”, “the empty set”) is the unique set that contains no elements whatsoever.
- $\emptyset = \{\} = \{x | \mathbf{False}\}$
- No matter the domain of discourse, we have the axiom


$$\neg \exists x: x \in \emptyset.$$

# Subset and Superset Relations

- $S \subseteq T$  (“ $S$  is a subset of  $T$ ”) means that every element of  $S$  is also an element of  $T$ .
- $S \subseteq T \Leftrightarrow \forall x (x \in S \rightarrow x \in T)$
- $\emptyset \subseteq S$ ,  $S \subseteq S$ .
- $S \supseteq T$  (“ $S$  is a superset of  $T$ ”) means  $T \subseteq S$ .
- Note  $S = T \Leftrightarrow S \subseteq T \wedge S \supseteq T$ .
- $S \not\subseteq T$  means  $\neg(S \subseteq T)$ , i.e.  $\exists x (x \in S \wedge x \notin T)$

# Proper (Strict) Subsets & Supersets

- $S \subset T$  (“ $S$  is a proper subset of  $T$ ”) means that  $S \subseteq T$  but  $T \not\subseteq S$ . Similar for  $S \supset T$ .



Example:

$$\{1,2\} \subset \{1,2,3\}$$

Venn Diagram equivalent of  $S \subset T$

# Sets Are Objects, Too!

- The objects that are elements of a set may *themselves* be sets.
- *E.g.* let  $S = \{x \mid x \subseteq \{1, 2, 3\}\}$   
then  $S = \{\emptyset,$   
           $\{1\}, \{2\}, \{3\},$   
           $\{1, 2\}, \{1, 3\}, \{2, 3\},$   
           $\{1, 2, 3\}\}$
- Note that  $1 \neq \{1\} \neq \{\{1\}\} !!!!$



# Cardinality and Finiteness

- $|S|$  (read “the *cardinality* of  $S$ ”) is a measure of how many different elements  $S$  has.
- *E.g.*,  $|\emptyset|=0$ ,  $|\{1,2,3\}| = 3$ ,  $|\{a,b\}| = 2$ ,  
 $|\{\{1,2,3\},\{4,5\}\}| = \underline{\underline{2}}$
- We say  $S$  is *infinite* if it is not *finite*.
- What are some infinite sets we’ve seen?

**N Z R**

# The *Power Set* Operation

- The *power set*  $P(S)$  of a set  $S$  is the set of all subsets of  $S$ .  $P(S) = \{x \mid x \subseteq S\}$ .
- *E.g.*  $P(\{a,b\}) = \{\emptyset, \{a\}, \{b\}, \{a,b\}\}$ .
- Sometimes  $P(S)$  is written  $2^S$ .  
Note that for finite  $S$ ,  $|P(S)| = 2^{|S|}$ .
- It turns out that  $|P(\mathbf{N})| > |\mathbf{N}|$ .  
*There are different sizes of infinite sets!*

# Ordered $n$ -tuples

- For  $n \in \mathbf{N}$ , an *ordered  $n$ -tuple* or a sequence of length  $n$  is written  $(a_1, a_2, \dots, a_n)$ . The *first* element is  $a_1$ , *etc.*
- These are like sets, except that duplicates matter, and the order makes a difference.
- Note  $(1, 2) \neq (2, 1) \neq (2, 1, 1)$ .
- Empty sequence, singlets, pairs, triples, quadruples, quintuples, ...,  $n$ -tuples.



# Cartesian Products of Sets


- For sets  $A, B$ , their *Cartesian product*  $A \times B \equiv \{(a, b) \mid a \in A \wedge b \in B\}$ .
- *E.g.*  $\{a, b\} \times \{1, 2\} = \{(a, 1), (a, 2), (b, 1), (b, 2)\}$
- Note that for finite  $A, B$ ,  $|A \times B| = |A||B|$ .
- Note that the Cartesian product is **not** commutative:  $\neg \forall A, B: A \times B = B \times A$ .
- Extends to  $A_1 \times A_2 \times \dots \times A_n \dots$

# The Union Operator

- For sets  $A$ ,  $B$ , their *union*  $A \cup B$  is the set containing all elements that are either in  $A$ , **or** (“ $\vee$ ”) in  $B$  (or, of course, in both).
- Formally,  $\forall A, B: A \cup B = \{x \mid x \in A \vee x \in B\}$ .
- Note that  $A \cup B$  contains all the elements of  $A$  **and** it contains all the elements of  $B$ :  
 $\forall A, B: (A \cup B \supseteq A) \wedge (A \cup B \supseteq B)$

# Union Examples

- $\{a,b,c\} \cup \{2,3\} = \{a,b,c,2,3\}$  **Required Form**
- $\{2,3,5\} \cup \{3,5,7\} = \{2,3,5,3,5,7\} = \{2,3,5,7\}$




# The Intersection Operator

- For sets  $A$ ,  $B$ , their *intersection*  $A \cap B$  is the set containing all elements that are simultaneously in  $A$  **and** (“ $\wedge$ ”) in  $B$ .
- Formally,  $\forall A, B: A \cap B \equiv \{x \mid x \in A \wedge x \in B\}$ .
- Note that  $A \cap B$  is a subset of  $A$  **and** it is a subset of  $B$ :  
$$\forall A, B: (A \cap B \subseteq A) \wedge (A \cap B \subseteq B)$$


# Intersection Examples

- $\{a,b,c\} \cap \{2,3\} = \underline{\quad \emptyset \quad}$
- $\{2,4,6\} \cap \{3,4,5\} = \underline{\quad \{4\} \quad}$



# Disjointedness

- Two sets  $A$ ,  $B$  are called *disjoint* (i.e., unjoined) iff their intersection is empty. ( $A \cap B = \emptyset$ )
- Example: the set of even integers is disjoint with the set of odd integers.



# Inclusion-Exclusion Principle


- How many elements are in  $A \cup B$ ?

$$|A \cup B| = |A| + |B| - |A \cap B|$$

- Example:

$$\{2,3,5\} \cup \{3,5,7\} = \{2,3,5,3,5,7\} = \{2,3,5,7\}$$

$$\{2,3,5\} \cup \{3,5,7\} = \{3, 5\}$$



Subtract out items in intersection, to compensate for double-counting them!

# Set Difference

- For sets  $A$ ,  $B$ , the *difference of  $A$  and  $B$* , written  $A - B$ , is the set of all elements that are in  $A$  but not  $B$ .
- $A - B \equiv \{x \mid x \in A \wedge x \notin B\}$   
 $= \{x \mid \neg( x \in A \rightarrow x \in B ) \}$
- Also called:  
The *complement of  $B$  with respect to  $A$* .




# Set Difference Examples

- $\{1, 2, 3, 4, 5, 6\} - \{2, 3, 5, 7, 9, 11\} =$   
 $\{1, 4, 6\}$
- $\mathbf{Z} - \mathbf{N} = \{\dots, -1, 0, 1, 2, \dots\} - \{0, 1, \dots\}$   
 $= \{x \mid x \text{ is an integer but not a nat. \#}\}$   
 $= \{x \mid x \text{ is a negative integer}\}$   
 $= \{\dots, -3, -2, -1\}$

# Set Difference - Venn Diagram

- $A-B$  is what's left after  $B$   
“takes a bite out of  $A$ ”



# Set Complements


- The *universe of discourse* can itself be considered a set, call it  $U$ .
- The *complement* of  $A$ , written  $\overline{A}$ , is the complement of  $A$  w.r.t.  $U$ , i.e., it is  $U-A$ .
- E.g., If  $U=\mathbf{N}$ ,

$$\overline{\{3,5\}} = \{0,1,2,4,6,7,\dots\}$$

# More on Set Complements

- An equivalent definition, when  $U$  is clear:

$$\overline{A} = \{x \mid x \notin A\}$$



# Set Identities

- Identity:  $A \cup \emptyset = A$   $A \cap U = A$
- Domination:  $A \cup U = U$   $A \cap \emptyset = \emptyset$
- Idempotent:  $A \cup A = A = A \cap A$
- Double complement:  $\overline{(\overline{A})} = A$
- Commutative:  $A \cup B = B \cup A$   $A \cap B = B \cap A$
- Associative:  $A \cup (B \cup C) = (A \cup B) \cup C$   
 $A \cap (B \cap C) = (A \cap B) \cap C$

# Set Identities

- Distributed:  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$   
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- Absorption:  $A \cup (A \cap B) = A$      $A \cap (A \cup B) = A$
- Complement:  $A \cap A' = \emptyset$      $A \cup A' = U$

# DeMorgan's Law for Sets

- Exactly analogous to (and derivable from) DeMorgan's Law for propositions.

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

# Proving Set Identities

To prove statements about sets, of the form

$E_1 = E_2$  (where  $E$ s are set expressions), here are three useful techniques:

- Prove  $E_1 \subseteq E_2$  and  $E_2 \subseteq E_1$  separately.
- Use logical equivalences.
- Use a *membership table*.



# Method 1: Mutual subsets

Example: Show  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

- Show  $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$ .
  - Assume  $x \in A \cap (B \cup C)$ , & show  $x \in (A \cap B) \cup (A \cap C)$ .
  - We know that  $x \in A$ , and either  $x \in B$  or  $x \in C$ .
    - Case 1:  $x \in B$ . Then  $x \in A \cap B$ , so  $x \in (A \cap B) \cup (A \cap C)$ .
    - Case 2:  $x \in C$ . Then  $x \in A \cap C$ , so  $x \in (A \cap B) \cup (A \cap C)$ .
  - Therefore,  $x \in (A \cap B) \cup (A \cap C)$ .
  - Therefore,  $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$ .
- Show  $(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C)$ . ...

# Method 2: logical Equivalence

- Prove that  $\overline{A \cup (B \cap C)} = (\bar{C} \cup \bar{B}) \cap \bar{A}$ 
  - $\overline{A \cup (B \cap C)} = \bar{A} \cap \overline{(B \cap C)}$  De Morgan's Law
  - $\quad \quad \quad = \bar{A} \cap (\bar{B} \cup \bar{C})$  De Morgan's Law
  - $\quad \quad \quad = (\bar{B} \cup \bar{C}) \cap \bar{A}$  Commutative Law
  - $\quad \quad \quad = (\bar{C} \cup \bar{B}) \cap \bar{A}$  Commutative Law

# Method 3: Membership Tables

- Just like truth tables for propositional logic.
- Columns for different set expressions.
- Rows for all combinations of memberships in constituent sets.
- Use “1” to indicate membership in the derived set, “0” for non-membership.
- Prove equivalence with identical columns.

# Membership Table Example

Prove  $(A \cup B) - B = A - B$ .

$A$	$B$	$A \cup B$	$(A \cup B) - B$	$A - B$
0	0	0	0	0
0	1	1	0	0
1	0	1	1	1
1	1	1	0	0

# Generalized Union

- Binary union operator:  $A \cup B$
- $n$ -ary union:  
 $A \cup A_2 \cup \dots \cup A_n \equiv (((A_1 \cup A_2) \cup \dots) \cup A_n)$   
(grouping & order is irrelevant)
- “Big U” notation:  $\bigcup_{i=1}^n A_i$
- Or for infinite sets of sets:  $\bigcup_{A \in X} A$

# Generalized Intersection

- Binary intersection operator:  $A \cap B$
- $n$ -ary intersection:  
 $A \cap A_2 \cap \dots \cap A_n \equiv (((A_1 \cap A_2) \cap \dots) \cap A_n)$   
(grouping & order is irrelevant)
- “Big Arch” notation:  $\bigcap_{i=1}^n A_i$
- Or for infinite sets of sets:  $\bigcap_{A \in X} A$

# Thank You

- Study all the solved problem from your text book.
- Try to solve related problems from exercise.