



HEAVY PERIODS

Key points

- Signs can include soaking pads or tampons quickly, bleeding that lasts a long time, passing large clots, or needing to change products overnight very often.
- Heavy bleeding can be linked with ovulation problems, fibroids, endometriosis, or PCOS.
- Ongoing heavy bleeding can contribute to anemia.

When to seek care

- Bleeding so heavy you feel dizzy or faint
- Shortness of breath, weakness, or paleness
- Very heavy bleeding after months without a period

Suggested Actions

- Track flow days and product use
- Ask about anemia if tired or dizzy
- Seek urgent care for severe bleeding symptoms

Key Takeaway

Heavy periods should be evaluated, especially if symptoms affect daily life.

Bloom provides educational guidance only. Always consult with a qualified healthcare professional for personalized medical advice.

Educational content informed by American College of Obstetricians and Gynecologists. (n.d.). Heavy menstrual bleeding. ACOG. <https://www.acog.org/womens-health/faqs/heavy-menstrual-bleeding>American College of Obstetricians and Gynecologists. (n.d.). Abnormal uterine bleeding. ACOG. <https://www.acog.org/womens-health/faqs/abnormal-uterine-bleeding>Eunice Kennedy Shriver National Institute of Child Health and Human Development. (n.d.). Menstruation and menstrual problems. <https://www.nichd.nih.gov/health/topics/menstruation>

