





SPRING 2019

Read This First!

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

1. Consider a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that has the following effect on the unit square.



- (a) [2 points] Find the matrix representation of T in the standard basis.
(There are two possible answers based on how you interpret the picture. You need only give one.)

- (b) [2 points] Determine $R(T)$ and $N(T)$ (no explanation is necessary for this part).

(continued on reverse)

- (c) [3 points] Find the matrix representation, in the standard basis, of the inverse transformation T^{-1} .

- (d) [3 points] Determine a point $\vec{x} \in \mathbb{R}^2$ that is sent to $\begin{pmatrix} 5 \\ 10 \end{pmatrix}$ by this transformation (i.e. $T(\vec{x}) = \begin{pmatrix} 5 \\ 10 \end{pmatrix}$).

2. Let A be the following 3×5 matrix.

$$A = \begin{pmatrix} 2 & -6 & -9 & -11 & -8 \\ 2 & -6 & -6 & -8 & -4 \\ 1 & -3 & -3 & -4 & 0 \end{pmatrix}$$

The reduced row echelon form of A is as follows (you do not need to verify this yourself).

$$\begin{pmatrix} 1 & -3 & 0 & -1 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

- (a) [3 points] Give a basis for $N(A)$.

- (b) [3 points] Give a basis for $R(A)$.

(continued on reverse)

- (c) [3 points] Let A be the same matrix above, and let \vec{b} be any vector in \mathbb{R}^3 . Explain why the matrix equation $A\vec{x} = \vec{b}$ is consistent, regardless of the choice of \vec{b} . How many free variables occur in the general solution of $A\vec{x} = \vec{b}$?

3. [9 points] Let

$$A = \begin{pmatrix} 2 & -2 & -3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{pmatrix}.$$

Determine the inverse matrix A^{-1} .

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

4. Suppose that $T : V \rightarrow W$ is a linear transformation of vector spaces, and $S = \{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is a set of three vectors in V .
- (a) [5 points] Suppose $\{T(\vec{v}_1), T(\vec{v}_2), T(\vec{v}_3)\}$ is a linearly independent set in W . Prove that S is a linearly independent set in V .

(continued on reverse)

- (b) [5 points] Prove that if S is a linearly independent set in V and T is one-to-one, then $\{T(\vec{v}_1), T(\vec{v}_2), T(\vec{v}_3)\}$ is a linearly independent set in W .

5. [9 points] Denote by $\vec{u}, \vec{v}, \vec{b}$ the following three vectors in \mathbb{R}^4 .

$$\vec{u} = \begin{pmatrix} 0 \\ -1 \\ 2 \\ 0 \end{pmatrix} \quad \vec{v} = \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} \quad \vec{b} = \begin{pmatrix} 0 \\ -11 \\ -11 \\ 11 \end{pmatrix}$$

Determine the linear combination \vec{w} of $\{\vec{u}, \vec{v}\}$ that is closest to \vec{b} (that is, the linear combination that minimizes $\|\vec{w} - \vec{b}\|$).

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

6. Let V be an inner product space. Suppose that W is a subspace of V with basis $B = \{\vec{w}_1, \vec{w}_2, \vec{w}_3\}$.

(a) [2 points] What is the dimension of W ?

- (b) [4 points] Suppose that \vec{b} is a vector in V , and \vec{u} is a vector in W such that

$$\vec{w}_1 \perp (\vec{b} - \vec{u}), \quad \vec{w}_2 \perp (\vec{b} - \vec{u}), \quad \text{and} \quad \vec{w}_3 \perp (\vec{b} - \vec{u}).$$

Prove that $\vec{b} - \vec{u}$ is orthogonal to every vector in W .

(continued on reverse)

(c) [4 points] Let \vec{b}, \vec{u} be as in part (b). Prove that if \vec{v} is any other vector in W , then

$$\|\vec{b} - \vec{v}\|^2 = \|\vec{b} - \vec{u}\|^2 + \|\vec{u} - \vec{v}\|^2.$$

7. Define a map $T : \mathcal{P}_2 \rightarrow \mathbb{R}^4$ by the formula

$$T(p(x)) = \begin{pmatrix} p(0) \\ p(1) \\ p(2) \\ p(3) \end{pmatrix}.$$

T is a linear transformation (you do not need to prove this).

- (a) [4 points] Let $B = \{(x-1)(x-2), x(x-2), x(x-1)\}$. This is a basis of \mathcal{P}_2 (you do not need to prove this). Let S denote the standard basis of \mathbb{R}^4 . Determine the matrix representation $[T]_B^S$ of T with respect to the basis B of \mathcal{P}_2 and the standard basis S of \mathbb{R}^4 .

- (b) [3 points] Let A be the matrix $[T]_B^S$ obtained in the previous part. Prove that $\vec{b} \in R(T)$ if and only if the matrix equation $A\vec{x} = \vec{b}$ is consistent.

- (c) [3 points] Suppose that we are given four constants a, b, c, d , and wish to find a polynomial $p(x) \in \mathcal{P}_2$ whose graph passes through the four points $(0, a), (1, b), (2, c), (3, d)$. For which values of a, b, c, d is this possible? Express your answer as a linear equation in a, b, c , and d .

Hint: interpret this as asking when a linear system is consistent, using the previous part.

8. [9 points] Let A be an $n \times n$ matrix, and λ a scalar. Define as usual the following set (called the *eigenspace* in class).

$$V_\lambda = \{\vec{v} \in \mathbb{R}^n : A\vec{v} = \lambda\vec{v}\}$$

Prove that V_λ is a *subspace* of \mathbb{R}^n .

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

9. In a certain town, 8000 customers subscribe to internet service from company A . A competitor, company B , enters the market and begins to draw customers from company A .

Suppose that every year, 10% of company A 's customers change their service to company B , and 30% of company B 's customers change their service to company A .

For example:

- In the first year, 800 customers (10% of 8000) change service from A and B , after which company A has 7200 customers and company B has 800 customers.
- In the second year, 720 customers (10% of 7200) switch from A to B , and 240 customers (30% of 800) switch from B to A . So after two years, company A has $7200 - 720 + 240 = 6720$ customers and company B has $800 + 720 - 240 = 1280$ customers.

- (a) [2 points] Find a 2×2 matrix M encoding the change in number of customers of each company from one year to the next. More precisely: if a, b denote the number of customers of companies A and B (respectively) in a given year, and a', b' denote the number of customers in the following year, the matrix M should satisfy

$$\begin{pmatrix} a' \\ b' \end{pmatrix} = M \begin{pmatrix} a \\ b \end{pmatrix}.$$

You can check your answer by verifying that

$$\begin{pmatrix} 7200 \\ 800 \end{pmatrix} = M \begin{pmatrix} 8000 \\ 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 6720 \\ 1280 \end{pmatrix} = M^2 \begin{pmatrix} 8000 \\ 0 \end{pmatrix}.$$

(continued on reverse)

- (b) [6 points] Find the eigenvalues of M , and an eigenvector for each eigenvalue.

- (c) [3 points] Express $\begin{pmatrix} 8000 \\ 0 \end{pmatrix}$ as a linear combination of the eigenvectors you found in part (b).

- (d) [3 points] Find a formula for $M^n \begin{pmatrix} 8000 \\ 0 \end{pmatrix}$, and use your formula to evaluate $\lim_{n \rightarrow \infty} M^n \begin{pmatrix} 8000 \\ 0 \end{pmatrix}$.

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).

This page intentionally left blank. You may use it for scratchwork or to continue answers to any question (note clearly on the original page if you do so).