
## Region 3

WWF

"ArcNet" scenario 33 achievement for region 3. Use Accenter for advanced mode.



Bellugs of the Bristol Bay spring-summer distribution   106.81%   90.90%   94.88   90.90%   94.88   90.90%   94.88   90.90%   94.88   90.90%   94.88   90.90%   94.88   94.8	CF	Name	Target Achieve- ment for Region	Proportion of Target Achievement in Region	Amount Proportion in Region
10.11   10.1	5010	Beluga of the Bristol Bay spring-summer distribution	106.8%	100.0%	96.6%
Belluga of the Easten Bering Sea summer core distribution 107.9% 99.5% 518.58 90.99 Belluga of the Bristol Bay autum-winter distribution 107.9% 99.5% 518.58 15.88 15.89 13.13. Birstol Bay autum-winter distribution 69.9% 69.9% 69.9% 69.98 13.13. Birstol Bay and Kusskokwin Bay inner sheff 61.5% 97.4% 52.38 10.00	3105	polynya Bering SE	785.8%	99.9%	94.4%
5009 Baluga of the Bristol Bay autumn-winter distribution         107.9%         99.5%         61.6%           5079 B.13.13 Bristol Bay and Kistokowin Bay inner shelf         401.5%         97.4%           6007 Common eider (Somatleria mollissima v.ingrum) wintering grounds         56.5%         46.6%         38.8%           5019 Beluga of the Eastern Bering Sea coast of southern Alaska and         206.5%         54.9%         55.8%           722 Kelp breats allow the Bering Sea coast of southern Alaska and         206.5%         54.9%         55.8%           723 Kelp forests of north-eastern coast of Bering Sea         200.3%         76.6%         30.0%           6077 Common murre (Uria asilge inomata) breeding colonies         58.3%         36.4%         38.8%           8077 Murgral Ide Zone distribution in April in the Eastern Bering Sea LME         227.6%         73.5%         28.8%           2008 Sported seal wheeliping areas as predicted by MIZ distribution         144.7%         80.7%         38.0%           2035 Sported seal wheeliping areas as predicted by MIZ distribution         144.7%         80.7%         38.0%           807 Long-lailed duck (Clangula hyemalis) East Asian wintering grounds         33.3%         32.2%         26.8%           807 Spowling graves of the Peadlic Capelini (Mallotus villosus catervarius)         105.4%         36.5%         28.8%	8018	Chukchi Sea estuaries	164.1%	86.2%	82.1%
13.1.3. Bristol Bay and Kuskokwin Bay inner shelf   40.15%   97.4%   65.3%   50.6%   50.8%	5018	Beluga of the Eastern Bering Sea summer core distribution	100.1%	99.4%	81.5%
6007   Common eider (Somateria mollissima v-injourn) withering grounds   60.8%   60.	5009	Beluga of the Bristol Bay autumn-winter distribution	107.9%	99.5%	61.6%
Beluga of the Eastern Bering Sea winter distribution   69,8%   54,9%   55,5%   55,1%   20,5%	7089	I.3.1.3. Bristol Bay and Kuskokwin Bay inner shelf	401.5%	97.4%	52.3%
Rel	6067	Common eider (Somateria mollissima v-nigrum) wintering grounds	56.5%	46.6%	39.8%
Aleutian Islands	5019	Beluga of the Eastern Bering Sea winter distribution	69.8%	69.6%	35.6%
15   15   15   15   15   15   15   15	7221		206.5%	54.9%	35.1%
3017   Marginal Ice Zone distribution in April in the Eastern Bering Sea LME   227.8%   73.5%   28.4%   2060   Ribbon seal whelping areas as predicted by MIZ distribution   144.7%   80.7%   27.0%	7233	Kelp forests of north-eastern coast of Bering Sea	200.3%	76.6%	34.0%
2050 Ribbon seal whelping areas as predicted by MIZ distribution         223.1%         76.4%         27.6%           2063 Spotted seal whelping areas as predicted by MIZ distribution         144.7%         80.7%         27.0%           2058 Bearded seal whelping areas in the Bering Sea as predicted by MIZ distribution         107.0%         38.0%         26.8%           6071 Long-failed duck (Clangula hyemalis) East Asian wintering grounds         33.3%         32.2%         26.0%           7088 I.3.1.2. Notno Sound inner shelf         105.1%         86.8%         25.4%           2054 Spotted seal franging areas         199.9%         39.4%         25.3%           4005 In Alaska (F1)         22.2%         32.2%         23.4%           4005 In Alaska (F1)         32.2%         23.4%         25.3%           4007 Gisucous guil (Larus hyperboreus barrovianus) breeding grounds         176.3%         54.5%         22.9%           4026 Feeding/migration area of the Coho Salmon (Oncorhynchus kisutch) (F2.6%         52.5%         22.8%         22.8%           50408 Black-legged kittiwake (Rissa tridactyla Indactyla) breeding colonies         34.9%         25.5%         22.8%           4026 Feeding/migration area of the Chinok Salmon (Oncorhynchus stutch) (F2.6%         311.1%         33.9%         12.5%         22.6%           504 Read (Warth Salman Salman Salman	6077	Common murre (Uria aalge inornata) breeding colonies	58.3%	36.4%	31.5%
2038         Spotted seal whelping areas as predicted by MIZ distribution         144.7%         80.7%         27.0%           2058         Bearded seal whelping areas in the Bering Sea as predicted by MIZ distribution         107.0%         38.0%         26.8%           6071         Long-tailed duck (Clangula hyemalis) East Asian wintering grounds         33.3%         32.3%         26.0%           7078         L3.12. Notron Sound Inner shelf         105.1%         86.8%         25.4%           2045         Spotted seal foraging areas         199.9%         39.4%         25.3%           4005         Spawning areas of the Pacific Capelin (Mallotus villosus catervarius) in Alaska (F11)         122.7%         32.2%         23.4%           6007         Glaucous guil (Larus hyerboreus barrovianus) breeding grounds         176.3%         54.5%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla tridactyla) breeding colonies         56.8%         38.4%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla breeding colonies         57.4%         32.2%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla tri	3017	Marginal Ice Zone distribution in April in the Eastern Bering Sea LME	227.8%	73.5%	28.4%
Bearded seal whelping areas in the Bering Sea as predicted by MIZ   107.0%   38.0%   26.8%   107.0%   38.0%   26.8%   107.0%   38.0%   26.8%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   38.0%   26.0%   107.0%   39.4%   26.0%   107.0%   39.4%   26.0%   107.0%   39.4%   26.0%   107.0%	2060	Ribbon seal whelping areas as predicted by MIZ distribution	223.1%	76.4%	27.6%
107.0%   30.0%   20.	2063	Spotted seal whelping areas as predicted by MIZ distribution	144.7%	80.7%	27.0%
7088         1.3.1.2. Norton Sound inner shelf         105.1%         86.8%         25.4%           2054         Spotled seal foraging areas         199.9%         394.4%         25.3%           4005         Spawing areas of the Pacific Capelin (Maliclus villosus catervarius) in Alaska (F11)         122.7%         32.2%         23.4%           4006         Glaucous guil (Larus hyperboreus barrovianus) breeding grounds         176.3%         54.5%         22.9%           4026         Feeding/migration area of the Coho Salmon (Oncorhynchus kisutch) (F 25)         358.6%         38.4%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla tridactyla) breeding colonies         34.9%         25.3%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla tridactyla) breeding colonies         34.9%         25.3%         22.8%           6048         Black-legged kittiwake (Rissa tridactyla tridactyla) breeding colonies         34.9%         25.3%         22.8%           4026         Feeding/migration area of the Chinock Salmon (Oncorhynchus tridatyla)         311.1%         33.9%         19.8%           4051         Feeding/migration area of the Chinock Salmon (Oncorhynchus tridatyla)         311.1%         33.9%         19.8%           4052         Feeding area of the Ceping Caperatyla speciallish specially specially specially special	2058		107.0%	38.0%	26.8%
2054   Spotted seal foraging areas   199.9%   39.4%   25.3%   23.4%	6017	Long-tailed duck (Clangula hyemalis) East Asian wintering grounds	33.3%	32.3%	26.0%
Spawning areas of the Pacific Capelin (Mallotus villosus catervarius) in Alaska (F11)	7088	I.3.1.2. Norton Sound inner shelf	105.1%	86.8%	25.4%
in Alaska (F11)  6027 Glaucous gull (Larus hyperboreus barrovianus) breeding grounds  Feeding/migration area of the Coho Salmon (Oncorhynchus kisutch)  (F 25)  6048 Black-legged kittiwake (Rissa tridactyla iridactyla) breeding colonies  4026 Feeding/migration area of the Chinook Salmon (Oncorhynchus  4027 Seeding/migration area of the Chinook Salmon (Oncorhynchus  4028 Seeding/migration area of the Chinook Salmon (Oncorhynchus  4038 Feeding/migration area of the Chinook Salmon (Oncorhynchus  4039 Seeding area of the Dolly Varden (Salvelinus malma) (F 30)  4030 Feeding area of the Dolly Varden (Salvelinus malma) (F 30)  4043 Ringed sal whelping areas in the Bering Sea  4044 Ringed sal whelping areas in the Bering Sea  4054 Seeding area of the Broad whitefish (Coregonus nasus), Euro-Asian  4064 Fish zoogeography, Paofilo Region, 6 - high-boreal and 7 - low-boreal  4075 districts of the Eastern North-Pacific Subtregion  4086 Bearded seal whelping areas in the Bering Sea  4097 Seeding area of the Broad whitefish (Coregonus nasus), Euro-Asian  4098 Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian  4099 Pacific Walrus Winter Distribution  4019 Pacific Walrus Winter Distribution  4010 Sea	2054	Spotted seal foraging areas	199.9%	39.4%	25.3%
6027 Glaucous gull (Larus hyperboreus barrovianus) breeding grounds feeding/migration area of the Coho Salmon (Oncorhynchus kisutch) (F25) 6048 Black-legged kittiwake (Rissa tridactyla breeding colonies 4026 6048 Black-legged kittiwake (Rissa tridactyla breeding colonies 4040 605 6064 Black-legged kittiwake (Rissa tridactyla breeding colonies 4065 6074 King eider (Sanwalianus) 6074 King eider (Sanwalianus) 6074 King eider (Somateria spectabilis) Pacific winetring grounds 6074 King eider (Somateria spectabilis) Pacific winetring grounds 6074 King eider (Somateria spectabilis) Pacific winetring grounds 6075 Stellers Sea Lion habitats of critical importance 6076 Fish 2009 Capraphy, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion 6107 Freeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19 6208 Bearded seal whelping areas in the Bering Sea 6107 6107 Fratercula comiculata breeding colonies 6107 Fratercula comiculata breeding colonies 6107 Fratercula comiculata breeding colonies 6107 Risstridactylpoliicarius breeding colonies 6108 Horned puffin (Fratercula comiculata) breeding colonies 610 Risstridactylpoliicarius breeding colonies 610 Risstridac	4005	Spawning areas of the Pacific Capelin (Mallotus villosus catervarius)	122.7%	32.2%	
Feeding/migration area of the Coho Salmon (Oncorhynchus kisutch)   358.6%   38.4%   22.8%	6027	• •	176.3%	54.5%	22.9%
Black-legged kittiwake (Rissa tridactyla tridactyla) breeding colonies   34.9%   25.3%   22.6%		Feeding/migration area of the Coho Salmon (Oncorhynchus kisutch)			
Feeding/migration area of the Chinook Salmon (Oncorhynchus tshawytscha) (F 26)  1015 Pacific Walrus Breeding Area  403 Feeding area of the Dolly Varden (Salvelinus malma) (F 30)  403 Feeding area of the Dolly Varden (Salvelinus malma) (F 30)  404 King elder (Somateria spectabilis) Pacific winetring grounds  405 Steller Sea Lion habitats of critical importance  406 Z7.3%  407 Steller Sea Lion habitats of critical importance  408 Fish zoogeography. Pacific Region, 6 - high-boreal and 7 - low-boreal  408 districts of the Eastern North-Pacific subregion  409 Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian  400 populations F 19  400 Bearded seal whelping areas in the Bering Sea  4017 Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian  4018 pacific Walrus Winter Distribution  4019 Pacific Walrus Winter Distribution  4019 Pacific Walrus Winter Distribution  4020 Searded seal whelping areas in the Bering Sea  403 S.5.%  404 S.5.%  405 S.2.9%  406 S.2.9%  407 S.2.9%  408 S.3.1.%  409 S.3.1.4 Middle and outer shelf of eastern Bering Sea  409 Searded Seal whelping areas in the Bering Sea  400 Searded Seal whelping areas in the Bering Sea  4010 Searded Seal whelping areas in the Bering Sea  402 Searded Seal whelping sea Searded Seal Walrus Winter Distribution  400 Searded Seal whelping Sea Searded Seal Search Se	6048	,	34.9%	25.3%	22.6%
1015 Pacific Walrus Breeding Area 4033 Feeding area of the Dolly Varden (Salvelinus malma) (F 30) 6074 King eider (Somateria spectabilis) Pacific winetring grounds 30.0% 20.7% 616.2% 2043 Ringed seal whelping areas in the Bering Sea 62.0% 62.0% 635.6% 635.6% 616.0% 62.055 Steller Sea Lion habitats of critical importance 627.3% 616.4% 615t Zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion 6406 Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19 6407 Earth Search S		Feeding/migration area of the Chinook Salmon (Oncorhynchus			
4033         Feeding area of the Dolly Varden (Salvelinus malma) (F 30)         135.2%         27.8%         17.2%           6074         King eider (Somateria spectabilis) Pacific winetring grounds         30.0%         20.7%         16.2%           2043         Ringed seal whelping areas in the Bering Sea         62.0%         35.6%         16.0%           2055         Steller Sea Lion habitats of critical importance         27.3%         16.4%         15.6%           4084         Fish zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion         179.4%         41.4%         15.6%           4016         Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19         43.8%         31.1%         15.4%           2006         Bearded seal whelping areas in the Bering Sea         54.8%         31.1%         14.2%           2017         Fractic Walrus Winter Distribution         47.8%         25.5%         12.6%           1019         Pacific Walrus Winter Distribution         47.8%         25.5%         12.6%           7247         seagrasses of the High Arctic         41.7%         22.9%         12.5%           7090         1.3.1.4. Middle and outer shelf of eastern Bering Sea         285.7%         29.1%         11.3%           6010 <td>1015</td> <td></td> <td>67.4%</td> <td>32 2%</td> <td>17 5%</td>	1015		67.4%	32 2%	17 5%
6074         King elder (Somateria spectabilis) Pacific winetring grounds         30.0%         20.7%         16.2%           2043         Ringed seal whelping areas in the Bering Sea         62.0%         35.6%         10.0%           2055         Steller Sea Lion habitats of critical importance         27.3%         16.4%         15.8%           4084         Fish zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion         179.4%         41.4%         15.6%           4016         Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19         43.8%         31.1%         15.4%           2006         Bearded seal whelping areas in the Bering Sea         54.8%         31.1%         14.2%           6107         6107 Fratercula comiculata breeding colonies         52.4%         20.4%         13.6%           1019         Pacific Walrus Winter Distribution         47.8%         25.5%         12.6%           7247         seagrasses of the High Arctic         41.7%         22.9%         12.5%           7090         1.3.1.4. Middle and outer shelf of eastern Bering Sea         285.7%         29.1%         11.3%           6100         6100 Risstridactylpollicarius breeding colonies         16.0%         16.0%         11.3%           6010		•			
2043         Ringed seal whelping areas in the Bering Sea         62.0%         35.6%         16.0%           2055         Steller Sea Lion habitats of critical importance         27.3%         16.4%         15.8%           4084         Fish zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion         179.4%         41.4%         15.6%           4016         Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19         43.8%         31.1%         14.2%           2006         Bearded seal whelping areas in the Bering Sea         54.8%         31.1%         14.2%           6107         6107 Fratercula corniculata breeding colonies         52.4%         20.4%         13.6%           1019         Pacific Waltrus Winter Distribution         47.8%         25.5%         12.6%           7247         seagrasses of the High Arctic         41.7%         22.9%         12.5%           7247         seagrasses of the High Arctic         41.7%         22.9%         12.5%           7090         13.1.4         Middle and outer shelf of eastern Bering Sea         286.7%         29.1%         11.3%           6103         floned puffin (Fratercula corniculata) breeding colonies         16.0%         16.0%         16.0%           6104					
2055         Steller Sea Lion habitats of critical importance         27.3%         16.4%         15.8%           4084         Fish zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion         179.4%         41.4%         15.6%           4016         Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19         43.8%         31.1%         14.2%           6107         6107 Fratercula corniculata breeding colonies         52.4%         20.4%         13.6%           6107         6107 Fratercula corniculata breeding colonies         52.4%         20.4%         13.6%           6107         9acific Walrus Winter Distribution         47.8%         25.5%         12.6%           7247         seagrasses of the High Arctic         41.7%         22.9%         12.5%           7090         1.3.1.4. Middle and outer shelf of eastern Bering Sea         285.7%         29.1%         11.3%           6100         6100 Risstridactylpoliicarius breeding colonies         16.0%         16.0%         11.3%           6085         Horned puffin (Fratercula corniculata) breeding colonies         28.0%         12.7%         10.9%           6010         Brent goose (Branta bernicla nigricans) American breeding&moulting grounds         42.8%         30.1%         10.8%		, , , , , , , , , , , , , , , , , , , ,			
Fish zoogeography, Pacific Region, 6 - high-boreal and 7 - low-boreal districts of the Eastern North-Pacific subregion  Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19  2006 Bearded seal whelping areas in the Bering Sea 54.8% 31.1% 14.2% 13.6% 1017 Fratercula corniculata breeding colonies 52.4% 20.4% 13.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 1019 Pacific Walrus Walrus Pacific Walrus		· · · · · · · · · · · · · · · · · · ·			
districts of the Eastern North-Pacific subregion  Feeding area of the Broad whitefish (Coregonus nasus), Euro-Asian populations F 19  Bearded seal whelping areas in the Bering Sea  54.8%  31.1%  14.2%  6107 6107 Fratercula comiculata breeding colonies  52.4%  20.4%  13.6%  1019 Pacific Walrus Winter Distribution  47.8%  25.5%  12.6%  7247 seagrasses of the High Arctic  6100 6100 Risstridactylpollicarius breeding colonies  6100 6100 Risstridactylpollicarius breeding colonies  16.0%  17.3%  6085 Horned puffin (Fratercula comiculata) breeding colonies  28.0%  12.7%  10.9%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  4019 Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)  8010 Northern Fur seal range  10.9%  10.7%  10.9%  10.7%  10.9%  10.7%  10.9%		·			
populations F 19 2006 Bearded seal whelping areas in the Bering Sea 54.8% 31.1% 14.2% 20.4% 13.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 27.2% 12.5% 12.6% 27.2% 12.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 27.2% 12.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 27.2% 12.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 27.2% 12.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 12.5% 12.6% 1019 Pacific Walrus Winter Distribution 47.8% 25.5% 12.6% 12.5	4084	districts of the Eastern North-Pacific subregion	179.4%	41.4%	15.6%
6107       6107 Fratercula corniculata breeding colonies       52.4%       20.4%       13.6%         1019       Pacific Walrus Winter Distribution       47.8%       25.5%       12.6%         7247       seagrasses of the High Arctic       41.7%       22.9%       12.5%         7090       I.3.1.4. Middle and outer shelf of eastern Bering Sea       285.7%       29.1%       11.3%         6100       6100 Risstridactylpollicarius breeding colonies       16.0%       16.0%       11.3%         6085       Horned puffin (Fratercula corniculata) breeding colonies       28.0%       12.7%       10.9%         2034       Ribbon seal foraging areas in the Bering Sea       167.5%       27.2%       10.9%         6010       Brent goose (Branta bernicla nigricans) American breeding&moulting grounds       42.8%       30.1%       10.8%         4019       Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)       41.8%       21.0%       10.7%         5054       Harbour porpoise feeding areas in the Bering Sea       160.9%       30.6%       10.7%         603       6103 Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6040       Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%	4016	populations F 19			
1019 Pacific Walrus Winter Distribution       47.8%       25.5%       12.6%         7247 seagrasses of the High Arctic       41.7%       22.9%       12.5%         7090 I.3.1.4. Middle and outer shelf of eastern Bering Sea       285.7%       29.1%       11.3%         6100 6100 Risstridactylpollicarius breeding colonies       16.0%       16.0%       11.3%         6085 Horned puffin (Fratercula corniculata) breeding colonies       28.0%       12.7%       10.9%         2034 Ribbon seal foraging areas in the Bering Sea       167.5%       27.2%       10.9%         6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds       42.8%       30.1%       10.8%         4019 Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)       41.8%       21.0%       10.7%         5054 Harbour porpoise feeding areas in the Bering Sea       160.9%       30.6%       10.7%         2032 Northern Fur seal range       139.1%       26.6%       10.7%         6103 G103 Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%         4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)       161.3%       19.1%       10.3%         5058 Humpback whale s					
7247       seagrasses of the High Arctic       41.7%       22.9%       12.5%         7090       I.3.1.4. Middle and outer shelf of eastern Bering Sea       285.7%       29.1%       11.3%         6100       6100 Risstridactylpollicarius breeding colonies       16.0%       16.0%       11.3%         6085       Horned puffin (Fratercula corniculata) breeding colonies       28.0%       12.7%       10.9%         2034       Ribbon seal foraging areas in the Bering Sea       167.5%       27.2%       10.9%         6010       Brent goose (Branta bernicla nigricans) American breeding&moulting grounds       42.8%       30.1%       10.8%         4019       Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)       41.8%       21.0%       10.7%         5054       Harbour porpoise feeding areas in the Bering Sea       160.9%       30.6%       10.7%         6103       Northern Fur seal range       139.1%       26.6%       10.7%         6103       6103 Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6087       Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%         4024       Distribution of the Chum Salmon (Oncorhynchus keta) (F24)       161.3%       19.1%       10.3% <td></td> <td>•</td> <td></td> <td></td> <td></td>		•			
7090 I.3.1.4. Middle and outer shelf of eastern Bering Sea       285.7%       29.1%       11.3%         6100 6100 Risstridactylpollicarius breeding colonies       16.0%       16.0%       11.3%         6085 Horned puffin (Fratercula corniculata) breeding colonies       28.0%       12.7%       10.9%         2034 Ribbon seal foraging areas in the Bering Sea       167.5%       27.2%       10.9%         6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds       42.8%       30.1%       10.8%         4019 Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)       41.8%       21.0%       10.7%         5054 Harbour porpoise feeding areas in the Bering Sea       160.9%       30.6%       10.7%         2032 Northern Fur seal range       139.1%       26.6%       10.7%         6103 G103 Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%         4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)       161.3%       19.1%       10.3%         5058 Humpback whale summer feeding areas in the Eastern Bering Sea       65.3%       59.2%       9.9%         1017 Pacific Walrus haulouts       12.0%       11.7%       9.8%         40					
6100 6100 Risstridactylpollicarius breeding colonies 16.0% 16.0% 11.3% 6085 Horned puffin (Fratercula corniculata) breeding colonies 28.0% 12.7% 10.9% 2034 Ribbon seal foraging areas in the Bering Sea 167.5% 27.2% 10.9% 6010 Brent goose (Branta bernicla nigricans) American breeding&moulting grounds 42.8% 30.1% 10.8% 4019 Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20) 41.8% 21.0% 10.7% 5054 Harbour porpoise feeding areas in the Bering Sea 160.9% 30.6% 10.7% 6103 6103 Urilomviarrrbreeding colonies 20.0% 15.0% 10.5% 6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds 48.3% 22.2% 10.4% 4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24) 161.3% 19.1% 10.3% 5058 Humpback whale summer feeding areas in the Eastern Bering Sea 65.3% 59.2% 9.9% 1017 Pacific Walrus haulouts 12.0% 11.7% 9.8% 4044 Range of the Pacific cod (Gadus macrocephalus) (F 38) 289.0% 19.6% 9.4% 5tellers eider (Polysticta stelleri) Pacific moulting&migration 17.1%					
6085       Horned puffin (Fratercula corniculata) breeding colonies       28.0%       12.7%       10.9%         2034       Ribbon seal foraging areas in the Bering Sea       167.5%       27.2%       10.9%         6010       Brent goose (Branta bernicla nigricans) American breeding&moulting grounds       42.8%       30.1%       10.8%         4019       Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)       41.8%       21.0%       10.7%         5054       Harbour porpoise feeding areas in the Bering Sea       160.9%       30.6%       10.7%         2032       Northern Fur seal range       139.1%       26.6%       10.7%         6103       Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6087       Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%         4024       Distribution of the Chum Salmon (Oncorhynchus keta) (F24)       161.3%       19.1%       10.3%         5058       Humpback whale summer feeding areas in the Eastern Bering Sea       65.3%       59.2%       9.9%         1017       Pacific Walrus haulouts       12.0%       11.7%       9.8%         4044       Range of the Pacific cod (Gadus macrocephalus) (F 38)       289.0%       19.6%       9.4% <td></td> <td></td> <td></td> <td></td> <td></td>					
Ribbon seal foraging areas in the Bering Sea  Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  Breeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)  5054 Harbour porpoise feeding areas in the Bering Sea  160.9%  Northern Fur seal range  139.1%  6103 Urilomviarrrbreeding colonies  6103 Urilomviarrrbreeding colonies  620.0%  10.5%  6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds  48.3%  22.2%  10.4%  4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  161.3%  161.3%  19.1%  10.9%  10					
Brent goose (Branta bernicla nigricans) American breeding&moulting grounds  42.8%  30.1%  10.8%  4019 Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)  5054 Harbour porpoise feeding areas in the Bering Sea  160.9%  30.6%  10.7%  2032 Northern Fur seal range  139.1%  6103 Urilomviarrrbreeding colonies  20.0%  515.0%  10.5%  6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds  48.3%  22.2%  10.4%  4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  5058 Humpback whale summer feeding areas in the Eastern Bering Sea  1017 Pacific Walrus haulouts  4048 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration		· · · · · · · · · · · · · · · · · · ·			
Feeding area of the Vendace, Least cisco (Coregonus sardinella), American populations (F 20)  5054 Harbour porpoise feeding areas in the Bering Sea  160.9%  30.6%  10.7%  2032 Northern Fur seal range  139.1%  6103 Urilomviarrrbreeding colonies  20.0%  15.0%  6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds  48.3%  22.2%  10.4%  4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  5058 Humpback whale summer feeding areas in the Eastern Bering Sea  65.3%  59.2%  9.9%  1017 Pacific Walrus haulouts  4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration		Brent goose (Branta bernicla nigricans) American breeding&moulting			
American populations (F 20)  5054 Harbour porpoise feeding areas in the Bering Sea  160.9%  30.6%  10.7%  2032 Northern Fur seal range  139.1%  26.6%  10.7%  6103 6103 Urilomviarrrbreeding colonies  20.0%  15.0%  10.5%  6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds  48.3%  22.2%  10.4%  4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  161.3%  19.1%  10.3%  5058 Humpback whale summer feeding areas in the Eastern Bering Sea  65.3%  59.2%  9.9%  1017 Pacific Walrus haulouts  12.0%  11.7%  9.8%  4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration		Feeding area of the Vendace, Least cisco (Coregonus sardinella),			_
2032 Northern Fur seal range       139.1%       26.6%       10.7%         6103 6103 Urilomviarrrbreeding colonies       20.0%       15.0%       10.5%         6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds       48.3%       22.2%       10.4%         4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)       161.3%       19.1%       10.3%         5058 Humpback whale summer feeding areas in the Eastern Bering Sea       65.3%       59.2%       9.9%         1017 Pacific Walrus haulouts       12.0%       11.7%       9.8%         4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)       289.0%       19.6%       9.4%         Stellers eider (Polysticta stelleri) Pacific moulting&migration       17.1%       15.0%       0.3%		, ,			
6103 Urilomviarrrbreeding colonies  20.0%  15.0%  10.5%  6087 Red-legged kittiwake (Rissa brevirostris) wintering grounds  48.3%  22.2%  10.4%  4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  5058 Humpback whale summer feeding areas in the Eastern Bering Sea  10.1%  10.5%  10.5%  10.5%  10.5%  10.4%  10.3%					
Red-legged kittiwake (Rissa brevirostris) wintering grounds 48.3% 22.2% 10.4% 4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24) 161.3% 19.1% 10.3% 19.1% 10.3% 19.1% 10.3% 19.6% 10.4% 10.3% 10.4% 10.3% 10.4% 10.3% 10.4% 10.3% 10.4% 10.3% 10.3% 10.4% 10.3% 10.3% 10.4% 10.4% 10.3% 10.4% 10					
4024 Distribution of the Chum Salmon (Oncorhynchus keta) (F24)  5058 Humpback whale summer feeding areas in the Eastern Bering Sea  65.3%  59.2%  9.9%  1017 Pacific Walrus haulouts  12.0%  11.7%  9.8%  4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration					
5058 Humpback whale summer feeding areas in the Eastern Bering Sea  1017 Pacific Walrus haulouts  12.0%  11.7%  9.8%  4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration  17.1%  15.0%  9.9%					
1017 Pacific Walrus haulouts 12.0% 11.7% 9.8% 4044 Range of the Pacific cod (Gadus macrocephalus) (F 38) 289.0% 19.6% Stellers eider (Polysticta stelleri) Pacific moulting&migration		, , , , , , , , , , , , , , , , , , , ,			
4044 Range of the Pacific cod (Gadus macrocephalus) (F 38)  Stellers eider (Polysticta stelleri) Pacific moulting&migration  17 196  289.0%  19.6%  9.4%		•			
Stellers eider (Polysticta stelleri) Pacific moulting&migration					Ξ.
		- ' ' ' ' '			_

CF	Name	Target Achieve- ment for Region	Proportion of Target Achievement in Region	Amount Proportion in Region
1018	Pacific Walrus Summer-Autumn Distribution	35.4%	15.2%	9.0%
4011	Feeding area of the Lake whitefish (Coregonus clupeaformis) (F 16)	21.0%	19.9%	8.7%
6073	King eider (Somateria spectabilis) Pacific moulting&migration stopovers	17.1%	14.8%	8.7%
7055	cold corals of the Pacific	12.8%	10. <mark>8%</mark>	8.0%
4004	Range of the Pacific Capelin (Mallotus villosus catervarius) (F11)	246.3%	18.0%	8.0%
6102	6102 Uriaalge inornatbreeding colonies	19.8%	11. <mark>8%</mark>	7.9%
4070	Local forms of the White-Sea herring (Clupea pallasii) (F8)	30.2%	17.0%	7.8%
4050	Range of the Pollock (Theragra chalcogramma) (F 43)	119.8%	17.5%	7.8%
4006	Feeding/nursery area of the Pacific rainbow smelt (Osmerus dentex) (F12)	38.6%	14 <mark>.6%</mark>	7.4%
4013	Feeding/nursery area of the Bering cisco (Coregonus laurettae), Asian populations (F17)	19.5%	18.7%	7.4%
4023	Feeding/migration area of the Pink Salmon (Oncorhynchus gorbuscha), native distribution (F23)	108.2%	16 <mark>.3%</mark>	7.0%
4058	Range of the Arctic flounder (Liopsetta glacialis) (F48)	106.9%	13 <mark>.8%</mark>	6.9%
6088	6088 Aethipusillbreeding colonies	26.2%	8.8 <mark>%</mark>	6.9%
4042	Range of the Saffron cod (Eleginus gracilis) (F 36)	106.0%	15 <mark>.6%</mark>	6.9%
6078	Common murre (Uria aalge inornata) wintering grounds	28.6%	18.1%	6.0%
6084	Thick-billed murre (Uria lomvia) Pacific wintering grounds	21.4%	18.1%	6.0%
4091	Fish zoogeography, Arctic Region, Subarctic Transitional-Pacific Province	30.2%	8.6%	4.7%
4055	Range of the Shorthorn Sculpin (Myoxocephalus scorpius) (F 46), American populations	112.6%	6.4 <mark>%</mark>	3.6%
6079	Thick-billed murre (Uria lomvia arrra) breeding colonies	4.7%	4.4%	3.4%
6039	Ivory gull (Pagophila eburnea) Pacific wintering grounds	11.6%	8.6 <mark>%</mark>	3.0%
6001	Least Auklet (Aethia pusilla) breeding colonies	8.1%	3.2%	2.9%
6066	Common eider (Somateria mollissima v-nigrum) breeding&moulting grounds	6.9%	6.8%	2.4%
4014	Feeding area of the Siberian whitefish (Coregonus pidschian) (F 18)	7.6%	4.8%	1.8%
9022	polar bear denning areas of CS (Chukchi Sea) subpopulation	2.1%	2.0%	1.8%
2040	Ringed seal molting areas in the Bering Sea	3.5%	2.7%	1.7%
5112	Arctic Cetaceans (beluga, bowhead, narwhal) winter habitats as predicterd by MIZ	3.7%	3.0%	1.5%
4053	Range of the Fourhorn Sculpin (Myoxocephalus quadricornis) (F 45), Euro-Asian populations	37.1%	2.2%	1.2%
3018	Marginal Ice Zone distribution in April in the Chukchi Sea LME	4.2%	0.9%	0.9%
9004	polar bear of the CS (Chukchi Sea) subpopulation distribution	2.0%	1.7%	0.8%
4041	Range of the Polar Cod (Boreogadus saida) (F35)	5.0%	1.9%	0.6%
6050	Spectacled eider (Somateria fischeri) moulting grounds	0.0%	0.0%	0.2%