

CHAPTER II :ROOTS AND WEIGHTS FOR $SL_n(\mathbb{R})/GL_n(\mathbb{R})$

May 30, 2025

In this chapter, we review some basis theory of roots and weight. We will first recall the general theory and compute explicitly the examples for $SL_n(\mathbb{R})/GL_n(\mathbb{R})$.

1 Structure theory

1.1 The Cartan subalgebra

First we need the notion of Cartan subalgebra

Definition 1.1. For any Lie algebra \mathfrak{g} , a subalgebra \mathfrak{h} of \mathfrak{g} is said to be Cartan algebra if it is

- \mathfrak{h} is a nilpotent subalgebra.
- It is self normalizing. In particular, we have $\mathfrak{h} = \{x \in \mathfrak{g} : [x, \mathfrak{g}] \subset \mathfrak{g}\}$.

When \mathfrak{g} is a semisimple Lie algebra, we have the following theorem

Theorem 1.2. Let \mathfrak{g} be a semisimple Lie algebra over an algebraically closed field k of characteristic 0 with a subalgebra \mathfrak{h} . Then \mathfrak{h} is a Cartan subalgebra of \mathfrak{g} if and only if it is a maximal toral subalgebra, i.e. is maximal among all subalgebras containing only semisimple elements.

add
cita-
tion.

1.2 Root space decomposition

With respect to some choice of Cartan subalgebra, we have a root space decomposition. In particular, there is a finite set $\Phi \subset \mathfrak{h}^*$ of linear forms on H , whose elements are called **roots**, such that

$$\mathfrak{g} = \mathfrak{h} \oplus \left(\bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha} \right),$$

where $\mathfrak{g}_{\alpha} = \{x \in \mathfrak{g} : [h, x] = \alpha(h)x, \forall h \in \mathfrak{h}\}$ for any $\alpha \in \Phi$.

1.3 A specific example: root space decomposition for $\mathfrak{sl}_n(\mathbb{R})$

For the semisimple Lie algebra $\mathfrak{sl}_n(\mathbb{R})$, a typical choice of the Cartan subalgebra is the set

$$\mathfrak{h} = \left\{ H = \begin{bmatrix} a_1 & 0 & \cdots & 0 \\ 0 & a_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_n \end{bmatrix}, a_1 + a_2 + \cdots + a_n = 0 \right\}$$

With respect to this Cartan subalgebra, we can define the linear function

$$L_i: \mathfrak{h} \rightarrow \mathbb{R}, \quad H \mapsto L_i(H) = a_i$$

Then the roots are given by $\alpha_{ij} := L_i - L_j$ for distinct i, j . We have the root space decomposition for $\mathfrak{sl}_n(\mathbb{R})$ as follows

$$\mathfrak{g} = \mathfrak{h} \oplus \left(\bigoplus \mathfrak{g}_{\alpha_{ij}} \right).$$

For the sake of brevity, we will denote $\alpha_{i,i+1}$ by α_i - these are called **simple roots**.

1.4 Roots at group level

Since the main object in this thesis is the Lie groups, we want to understand how the roots behave at group level. The analog for the Cartan subalgebra is the maximal torus

$$T = \left\{ t = \begin{bmatrix} a_1 & 0 & \dots & 0 \\ 0 & a_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_n \end{bmatrix} : a_i \neq 0 \right\},$$

Then T acts on \mathfrak{g} by conjugation. Explicitly, we can check that

$$\text{Ad}(t)(E_{ij}) = t_i t_j^{-1} E_{ij}$$

Therefore, at the group level, the character $\alpha_{ij}(\text{diag}(t_1, \dots, t_n)) = t_i t_j^{-1}$ is a root whenever $i \neq j$. The set

$$\Delta = \{ \alpha_i \mid i = \overline{1, n} \}$$

where

$$\alpha_i: T \mapsto \mathbb{R}, t \mapsto \frac{t_i}{t_{i+1}}$$

is the set of **simple roots**. We can decompose the set of root into to disjoint subsets, namely

$$\Phi = \{ \alpha_{ij}, i \neq j \} = \Phi_+ \coprod \Phi_-$$

where the set Φ_+ comprises of α_{ij} for $i < j$ and the remaining roots are in Φ_- . The former comprises of **positive roots** while the latter contains **negative roots**. We have the following lemma

Lemma 1.3. *Each $\alpha \in \Phi$ can be written uniquely as a linear combination*

$$\alpha = m_1 \alpha_1 + \dots + m_d \alpha_d$$

with all $m_i \in \mathbb{Z}_{\geq 0}$ or $m_i \in \mathbb{Z}_{\leq 0}$. If $\alpha \in \Phi_+$ then all $m_i \geq 0$, otherwise $m_i \leq 0$ for all i .

1.5 Weights

Another class of linear forms that we are interested in are the **fundamental weights**. For each fundamental weights λ_i , we define

$$\lambda_i: T \rightarrow \mathbb{R}, \quad \lambda_i(t) = a_1 \dots a_i$$

We have the following

Lemma 1.4. *We can write*

$$\lambda_i := r_1 \alpha_1 + r_2 \alpha_2 + \dots + r_d \alpha_d$$

where r_i 's are rational number such that $r_i \geq 0$. *add proof*

Example 1.5. *When $n = 3$, we have the following relations* *add picture*

$$\lambda_1 = \frac{2}{3} \alpha_1 + \frac{1}{3} \alpha_2, \quad \lambda_2 = \frac{1}{3} \alpha_1 + \frac{2}{3} \alpha_2$$

define
in
term
of Lie
alge-
bra

Definition 1.6. A weight λ is called **dominant** if it satisfies $\langle \lambda, \alpha^\vee \rangle \in \mathbb{Z}_{\geq 0}$ for all α

Clearly by lemma 1.3, the weight λ is dominant if and only if $\langle \lambda, \alpha_i^\vee \rangle$ for all fundamental root α_i . It is also clearly that the set of fundamental weight is given by addition of the fundamental weights, namely

$$\Lambda^+ := \{c_1\lambda_1 + \dots + c_d\lambda_d \mid c_i \in \mathbb{Z}_{\geq 0}\}$$

The set of dominant weights is denoted Λ^+ . A weight $\lambda = \sum n_i \lambda_i$ is called strongly dominant if $n_i > 0$ for all i . One important example is the minimal strongly dominant weight given by

$$\rho = \sum \lambda_i$$

This is called **Weyl vector** and is characterized in several ways:

1. $\langle \rho, \alpha_i^\vee \rangle = 1$ for all i .

- 2.

$$\rho = \frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha$$

To prove the last equation we use the action of the Weyl group W . Let $\mu = \frac{1}{2} \sum \alpha$. Apply the simple reflection s_i given by

$$s_i(x) = x - \langle x, \alpha_i^\vee \rangle \alpha_i$$

We know that s_i sends α_i to $-\alpha_i$ and permutes the other positive roots. So:

$$s_i(\mu) = \mu - \langle \mu, \alpha_i^\vee \rangle \alpha_i$$

Therefore, $(\mu, \alpha_i) = \mu(h_i) = 1$ for all i . So, $\mu = \rho$.

Unlike lemma 1.4, if we try to express the fundamental weights in term of the fundamental roots, we don't always get positive coefficients. However, it is true that all the coefficients must be integer. In particular, we have

$$\alpha_j = \sum_{n_j} \lambda_j, \quad n_j \in \mathbb{Z}.$$

To put it another way, the root lattice $\mathbb{Z}\Delta$ is contained inside the weight lattice.

1.6 Weyl group

We only define the Weyl group explicitly for the group $\mathrm{SL}_n(\mathbb{R})$ or $\mathrm{GL}_n(\mathbb{R})$. It is a fact that the Weyl groups for these two Lie groups are the same and equal to $W = S_n$ - the permutation group of n letters. We recall some basis observation about this group

1. Every $\sigma \in W$ can be written (non-uniquely) as a product of $w_{i_1} \dots w_{i_k}$ for some integer k . Such a sequence is said to have length k . If k is the minimum, over all such writings, it is called the length of σ and written $\ell(\sigma)$. Any expression of length $\ell(\sigma)$ for σ is called a reduced expression.
2. The group S_n is generated by S subject to the following two types of relations:
 - (Reflection) $w_i^2 = 1$ for $i \in I$.
 - (Braid relations) $w_i w_{i+1} w_i = w_{i+1} w_i w_{i+1}$ for $i = 1, \dots, n-2$ and $w_i w_j = w_j w_i$ for $|j - i| \geq 2$.

Note that W acts on $\mathrm{Hom}(H, \mathbb{R}^*)$ in the natural way: $w \cdot \varphi(h) = \varphi(w^{-1}h)$. More explicitly, σ sends $\alpha := \alpha_{ij}$ to $\alpha_{\sigma(i), \sigma(j)}$. Hence we find that

$$w_i \alpha_j = \begin{cases} -\alpha_i & \text{if } i = j \\ \alpha_j & \text{if } |j - i| > 1 \\ \alpha_i + \alpha_j & \text{if } |j - i| = 1 \end{cases}$$

We of course also have an action of W on the weights. For example, one can verify that

$$s_i(\lambda_i) = \lambda_i - \alpha_i \quad \text{and} \quad s_j(\lambda_i) = \lambda_i \text{ for } i \neq j .$$

Recall the definition of Weyl vector ρ , we have the following generalized action of Weyl group of ρ :

$$w\rho = \rho - \sum_{\alpha \in \Delta_{w^{-1}}} \alpha,$$

where **check this explicitly**

$$\Delta_\sigma := \{\alpha \in \Phi_+ \mid \sigma(\alpha) \in \Phi_-\}$$

1.7 Cartan matrix

We fix a set of simple roots $\Delta = \{\alpha_1, \dots, \alpha_d\}$ is define to be the matrix

$$A = [\langle \alpha_i, \alpha_j^\vee \rangle]$$

If we let $a_{ij} = \langle \alpha_i, \alpha_j^\vee \rangle$ then the Cartan matrix has the following simple properties:

Lemma 1.7.

- For any i , we have $a_{ii} = 2$.
- For any $i \neq j$, a_{ij} is a non-positive integer, i.e. $a_{ij} \in \mathbb{Z}_{\leq 0}$.

We give an explicit example for $\mathfrak{sl}_n(\mathbb{R})$, which has the root system A_n . The corresponding Cartan matrix is

$$A = \begin{bmatrix} 2 & -1 & 0 & 0 & \dots & 0 \\ -1 & 2 & -1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & -1 & 2 & -1 \\ 0 & 0 & \dots & 0 & -1 & 2 \end{bmatrix}$$

The Cartan matrix provides a linear combination presentation of the fundamental roots α_j 's in term of fundamental weights λ_i , as stated in the following lemma

Lemma 1.8. Fix a number n and consider the set of fundamental roots α_j as well as the set of weights λ_i of $SL_n(\mathbb{R})$, we have the following relations

$$\alpha_i = \begin{cases} 2\lambda_1 - \lambda_2, & \text{if } i = 1 \\ -\lambda_{i-1} + 2\lambda_i - \lambda_{i+1}, & \text{if } 1 < i < n . \\ -\lambda_{n-1} + 2\lambda_n, & \text{if } i = n \end{cases}$$

Proof. Note that we have

$$\langle \alpha_i, \alpha_j^\vee \rangle = \begin{cases} 2, & \text{if } i = j \\ -1, & \text{if } |i - j| = 1 \\ 0, & \text{otherwise} \end{cases} .$$

So if we let $\alpha_j = \sum_{i=1}^n a_{ij} \lambda_i$ and compute $\langle \alpha_i, \alpha_j^\vee \rangle$, the result follows immediately. \square

2 Parabolic subgroups

We shall provide two equivalent viewpoints on parabolic subgroups. They will play different roles in defining different notions of semi-stability in the next chapter.

2.1 Parabolic subgroups I: An explicit description

For our purpose, it is enough to define the standard parabolic subgroups. There exists a bijection between each parabolic subgroup of $SL_n(\mathbb{R})$ and each partition of n . We can therefore define the parabolic subgroup explicitly as follows:

Definition 2.1. *The standard parabolic subgroup associated to the partition $n = n_1 + n_2 + \cdots + n_r$ is denoted P_{n_1, \dots, n_r} and is defined to be the group of all matrices of the form*

$$\begin{bmatrix} \mathbf{m}_1 & 0 & \cdots & 0 \\ 0 & \mathbf{m}_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \mathbf{m}_k \end{bmatrix},$$

where $\mathbf{m}_{n_i} \in GL(n_i, \mathbb{R})$ for $1 \leq i \leq r$. The integer r is called the rank of the parabolic subgroup P_{n_1, \dots, n_r} .

Definition 2.2. *The maximal standard parabolic subgroups in $GL_n(k)$ corresponds to the stabilizer of the flag of type $\rho_i = (i, n - i)$, where $i = 1, \dots, n - 1$ of n . We will further denote $Q_i = P_{\rho_i}$ and **MaxParSt** the collection of such maximal parabolic subgroups.*

Example 2.3. *Below we list all the standard parabolic subgroup in $GL_3(\mathbb{R})$ and $GL_4(\mathbb{R})$.*

- For $GL_3(\mathbb{R})$, there are three standard parabolic subgroups corresponding to three partitions of 3, namely

$$3 = 1 + 1 + 1, \quad 3 = 1 + 2, \quad 3 = 2 + 1$$

For a partition (r_1, \dots, r_{s+1}) , we denote $P_{(r_1, \dots, r_{s+1})}$ the corresponding parabolic subgroups. Thus we have

$$P_{1,1,1} = \left\{ \begin{bmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & * \end{bmatrix} \right\}, \quad P_{1,2} = \left\{ \begin{bmatrix} * & * & * \\ 0 & * & * \\ 0 & * & * \end{bmatrix} \right\}$$

$$P_{2,1} = \left\{ \begin{bmatrix} * & * & * \\ * & * & * \\ 0 & 0 & * \end{bmatrix} \right\}$$

Clearly **MaxParSt** = $\{P_{2,1}, P_{1,2}\}$.

- For $GL_4(\mathbb{R})$, there are seven standard parabolic subgroups for seven partitions

$$4 = 1 + 1 + 1 + 1, \quad 4 = 1 + 1 + 2, \quad 4 = 1 + 2 + 1$$

$$4 = 2 + 1 + 1, \quad 4 = 1 + 3, \quad 4 = 2 + 1, \quad 4 = 3 + 1$$

Explicitly, we have the following subgroups

$$\begin{aligned}
 P_{1,1,1,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\}, & P_{1,1,2} &= \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{bmatrix} \right\} \\
 P_{1,2,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\}, & P_{2,1,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\} \\
 P_{1,2,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\}, & P_{2,1,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\} \\
 P_{1,3} &= \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{bmatrix} \right\}, & P_{3,1} &= \left\{ \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\} \\
 P_{2,2} &= \left\{ \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{bmatrix} \right\}
 \end{aligned}$$

Clearly $\mathbf{MaxParSt} = \{P_{1,3}, P_{3,1}, P_{2,2}\}$.

2.2 Parabolic subgroups II: Using BN-pairs

We first introduce the BN-pairs

Definition 2.4 (BN-pairs). A **BN-pairs** is a 4-tuple (G, B, N, R) where G is a group generated by subgroups B and N . The subgroup $H = B \cap N$, R is a finite set of involutions which generate the Weyl group $W = N/H$. Moreover the following theorem holds

- If $r \in R$ and $w \in W$, then $rBw \subset BwB \cup BrwB$.
- If $r \in R$, $rBr \neq B$.

For our purpose, it is enough to concentrate on the following example

Example 2.5. Let $G = GL_n(\mathbb{R})$, then the sets B, N, R are given explicitly as follows

- $B =$ upper triangular matrices
- $N =$ monomial matrices, namely, matrices that have exactly one non-zero entry in each row and column
- From the above, it is clearly that $H = B \cap N$ is the diagonal group, and this group is normal in N .
- It can be shown that $W = N/H \cong S_n$, thus $R = \{(i, i-1)\}$ - the set of transpositions.

Let $J \subset R$, we define W_J to be the subgroup of W generated by the involutions $r \in R$. We call it **standard parabolic subgroup** of W . Set $P_J = BW_JB$ as in the notation of BN-pairs. We have the following theorem

Theorem 2.6.

- P_J is a subgroup of G . In particular, we have

$$G = BWB,$$

which is called **Bruhat decomposition** of G .

- If $P_I = P_J$ then we have $I = J$.
- All subgroups of G containing B arises in this way. *add proofs?*

The above theorem leads to the following definition of parabolic subgroups

Definition 2.7 (Parabolic subgroups). *Using the same notation in the previous theorem, we call the subgroups P_J with $J \subset R$ standard parabolic subgroups of group G .*

We would like to explicitly describe the parabolic subgroup for $SL_n(\mathbb{R})$.

Example 2.8. *As introduced in previous section, the set $\Phi = \{\alpha_{ij}\}$ forms a root system for $SL_n(\mathbb{R})$. The Borel subgroup is just $B \cap SL_n(\mathbb{R})$, the group of upper triangular matrices with determinant 1. We also consider the set N of monomial matrices with determinant 1. Then it is easy to check that N/H is the set of all permutation matrices. That is, N/H is generated by the matrices of the form *see Cambridge**

$$r_i := \begin{bmatrix} I_{i-1} & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & I_{n-i-1} \end{bmatrix}$$

Via the identification $s_i \mapsto (i, i+1)$, we identify the Weyl group $W = N/H \cong S_n$. So $R = \{(i, i+1) | i = 1, 2, \dots, n\}$. We consider the set $I = R \setminus \{r_i\}$ for some $i < n$. The associated parabolic subgroup of W is

$$W_I = \langle r_1, r_2, \dots, r_{i-1}, r_{i+1}, \dots, r_{n-1} \rangle \cong S_i \times S_{n-i}$$

The corresponding parabolic subgroup is

$$P_{r_i} := P_I = \left\{ \begin{bmatrix} A & * \\ & B \end{bmatrix} \in SL_n(\mathbb{R}) : A \in GL_i, B \in GL_{n-i} \right\}$$

2.3 Parabolic sets and parabolic subalgebras

Definition 2.9. *Given a root system Δ . A **parabolic subset** Δ_P is a subset of Δ such that it satisfies the following conditions:*

1. *For any $\alpha \in \Delta$, at most one of the two elements $\alpha, -\alpha$ is contained in Δ_P .*
2. *It is closed, in the sense that, for any two root $\alpha, \beta \in \Delta_P$ such that $\alpha + \beta$ is a root, then $\alpha + \beta \in \Delta_P$.*

The parabolic set parametrizes the parabolic subalgebra with the root system Δ , as given in the following theorem

Theorem 2.10. *Given a semisimple Lie algebra \mathfrak{g} with the root system Δ . There exists a correspondence between parabolic subset of Δ_P of Δ and the subalgebra of \mathfrak{g} containing the Borel subalgebra \mathfrak{b} . The correspondence is given by*

$$\Delta_P \longleftrightarrow \mathfrak{p} := \mathfrak{b} \oplus \bigoplus_{\alpha \in \Delta \setminus \Delta_P} g_\alpha$$

Proof. We refer to [?] for a proof of this fact. □

Example 2.11. We consider the case $\mathfrak{g} = \mathfrak{sl}_3(\mathbb{R})$. Let's denote $\Pi = \{\alpha, \beta\}$ a base for the root system of \mathfrak{g} . It is clear that the set of positive root is $\Delta_+ = \{\alpha, \beta, \gamma\}$. There are 4 parabolic sets, corresponding to 4 parabolic subalgebra given as follows

$$\begin{aligned}\Delta_P &= \Delta_+ \longleftrightarrow \mathfrak{p} = \mathfrak{b} \\ \Delta_P &= \Delta \cup \{-\alpha\} \longleftrightarrow \mathfrak{p} = \mathfrak{b} \oplus \mathfrak{g}_{-\alpha} \\ \Delta_P &= \Delta \cup \{-\beta\} \longleftrightarrow \mathfrak{p} = \mathfrak{b} \oplus \mathfrak{g}_{-\beta} \\ \Delta_P &= \Delta \longleftrightarrow \mathfrak{p} = \mathfrak{g}\end{aligned}$$

2.4 Langlands decomposition

We fix a partition of n as

$$n = n_1 + n_2 + \dots + n_k$$

and consider the parabolic subgroup of this type, i.e. the subgroup

$$P_{n_1, \dots, n_k} = \left\{ \begin{bmatrix} \mathbf{m}_1 & * & \dots & * \\ 0 & \mathbf{m}_2 & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{m}_k \end{bmatrix} \right\}$$

where \mathbf{m}_i is invertible of size $n_i \times n_i$

This group can be factored as

$$P_{n_1, \dots, n_k} = M_{n_1, \dots, n_k} N_{n_1, \dots, n_k}$$

where

$$N_{n_1, \dots, n_k} = \left\{ \begin{bmatrix} I_1 & * & \dots & * \\ 0 & I_2 & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & I_k \end{bmatrix} \right\} \quad (I_k \text{ is the } n_k \times n_k \text{ identity matrix})$$

and

$$M_{n_1, \dots, n_k} = \left\{ \begin{bmatrix} \mathbf{m}_1 & 0 & \dots & 0 \\ 0 & \mathbf{m}_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{m}_k \end{bmatrix} \right\}$$

The subgroup M_{n_1, \dots, n_k} is called **Levi component**. We can further factor this subgroup as

$$M_{n_1, \dots, n_k} = M'_{n_1, \dots, n_k} \cdot A_{n_1, \dots, n_k}$$

with A_{n_1, \dots, n_k} plays the role of the connected center of M_{n_1, \dots, n_k} :

$$A_{n_1, \dots, n_k} = \left\{ \begin{bmatrix} t_1 I_1 & 0 & \dots & 0 \\ 0 & t_2 I_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & t_k I_k \end{bmatrix} : t_i \neq 0 \right\}$$

and

$$M'_{n_1, \dots, n_k} = \left\{ \begin{bmatrix} \mathbf{m}'_1 & 0 & \dots & 0 \\ 0 & \mathbf{m}'_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{m}'_k \end{bmatrix} \right\},$$

where $\det(\mathbf{m}'_i) = \pm 1$.

Definition 2.12. For a given parabolic subgroup P , the factorization

$$P = M_P \times A_P \times N_P$$

as above is called **Langlands decomposition**.

2.5 On the function H_P