

Why Schemes

Maria Iakerson

August 11, 2024

1 Summary of affine varieties

Let k be an algebraic closed field. The main idea of classical algebraic geometry is that we have a correspondence

$$\begin{aligned} \{\text{subsets of } k^n \text{ cutout by polynomials}\} &\leftrightarrow \{\text{finitely generated reduced } k\text{-algebras}\} \\ \text{Geometry} &\leftrightarrow \text{Algebra} \end{aligned}$$

In particular, the above correspondence can be given as follows:

- $I \subset k[x_1, \dots, x_n]$ ideal: Then we define

$$X := Z(I) = \{a \in k^n \mid f(a) = 0 \quad \forall f \in I\}$$

sometimes we use the notation $V(I)$ instead of $Z(I)$. This kind of set is an affine variety.

- \mathbb{A}^n : n -dimensional affine space. As a set, it is just k^n , but we equip this set with *Zariski* topology - where the closed subsets are generated by $Z(I)$.
- $I(X) := \{f \in k[x_1, \dots, x_n] \mid f(x) = 0 \quad \forall x \in X\}$. Then the quotient replacing $k[X] := \frac{k[x_1, \dots, x_n]}{I(X)}$ is called *coordinate ring* of X .
- $k[X]$ parametrizes functions on X .
- Hilbert's weak Nullstellensatz:

$$\{\text{points of } X\} \leftrightarrow \{\text{maximal ideals of } k[X]\}$$

- Hilbert Nullstellensatz: $I(Z(I)) = \sqrt{I} := \{f : f^n \in I \text{ for some } n\}$
- Morphisms: given X and $Y \in \mathbb{A}^n$ a morphism between two affine varieties is given by $\varphi = (f_1, \dots, f_n)$. This morphism induces a k -algebra homomorphism

$$\varphi^* k[Y] \rightarrow k[X], \quad \varphi^*(\psi) = \psi \circ \varphi$$

2 Why varieties are not good enough?

Some possible reasons are:

1. embedding into \mathbb{A}^n shouldn't really be part of the data. It would be nice to have an intrinsic definition, since you can embed the same variety in different spaces.
2. for non-algebraic closed field, the Nullstellensatz doesn't work.

-
3. We can ask, on which topological space is $\mathbb{R}[x, y]/(x^2 + y^2 + 1)$ naturally a functions space? Or $\mathbb{Z}[x]$? Or \mathbb{Z} ? This leads to the ideal of considering all possible rings.
 4. nilpotent arises naturally when deforming varieties, so ignoring them is not a good option.