# Semi-stable lattices in higher rank

Tri Nguyen

April 25, 2025

## Outline

Introduction

2 In 2 dimensional



### Historical motivation

Serre [1977] and Quillen [see Grayson, 1982] used the notion of semistable vector bundle on an algebraic curve to study  $\mathrm{SL}_n(\mathcal{O})$  when  $\mathcal{O}$  is a Dedekind domain finitely generated over a finite field. Stuhler then realized he can used the same method to adapt some work of Harder and Narasimhan on stable vector bundles to yields new facts about lattices in a Euclidean space.

Due to [?] , it is heuristical that the semi-stable lattices are the lattices in which the successive minima are closed.

## Definition of two-dimensional lattices

### Lattice

A lattice  $L \subset \mathbb{R}^2$  is a set of the form

$$L = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2$$

where  $e_1$ ,  $e_2$  are linearly independent over  $\mathbb{R}$ .

# Example of a lattice




Figure: Example of a lattice

### Classification of lattices

Do we know all the possible 2 dimensional "lattice shape"?

### Classification of lattices

Do we know all the possible 2 dimensional "lattice shape"?

**Answer:** Up to maginification, rotation and change of basis, the answer is yes.

## Fundamental domain

Up to rorations and magnifications, we can reduce a lattice

$$L = \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$$

to a lattice of the form

$$L_z = \mathbb{Z}z \oplus \mathbb{Z}, \quad \Im(z) > 0$$

So the upper half-plane parametrizes the 2 dimensional lattices.

#### Classification of unit lattices

The map  $z\mapsto \mathbb{Z}z\oplus \mathbb{Z}$  induces a bijection

$$\mathsf{SL}_2(\mathbb{Z})\backslash\mathbb{H}\cong\{\ \mathsf{lattices}\}/\mathbb{C}^\times$$



#### Fundamental domain

So we reduce to study the space of lattices by looking the action of  $SL_2(\mathbb{Z})$  on the upper half plane. Geometrically, the domain is given by

$$\mathfrak{D} = \{ z = x + iy \in \mathbb{H} : |z| \ge 1, -1/2 \le x \le 1/2 \}$$


,

### Fundamental domain

So we reduce to study the space of lattices by looking the action of  $SL_2(\mathbb{Z})$  on the upper half plane. Geometrically, the domain is given by

$$\mathfrak{D} = \{ z = x + iy \in \mathbb{H} : |z| \ge 1, -1/2 \le x \le 1/2 \}$$

,



Grayson associated each lattice L to some kind of **Newton polygon**.

Grayson associated each lattice L to some kind of **Newton polygon**.

The process is as follows:

- $\bullet$  Put (0,0) to the plot.
- ② For each primitive vector  $v \in L$ , he assigns the point  $(1, \log(||v||))$  to the plot.
- **3** Put the point  $(2, \log(vol(L)))$  to the plot.




Figure: The figure on the left corresponds to z=-2/5+3i/2 and on the right corresponds to z=7/16+15i/16

Since the lattice is discrete, there is a shortest primitive vector - on the plot we have the lowest point.

Since the lattice is discrete, there is a shortest primitive vector - on the plot we have the lowest point.

Grayson called the set of points plotted above as **Canonical plot**. The convex hull of the collection of the plot point is called **profile**.

For any  $z \in \mathbb{H} = \{ \text{Im}(z) > 0 \}$ , we can assign to it a unit lattices

$$z \mapsto L_z = \mathbb{Z} \frac{e_1}{\sqrt{y}} + \mathbb{Z} \left( \frac{x}{\sqrt{y}} e_1 + \sqrt{y} e_2 \right)$$

The shortest vector is then  $e_1/\sqrt{y}$ , with length  $\frac{1}{\sqrt{y}}$ . So for y < 1, the lowest point is below the horizontal axis.

For any  $z \in \mathbb{H} = \{ \text{Im}(z) > 0 \}$ , we can assign to it a unit lattices

$$z \mapsto L_z = \mathbb{Z} \frac{e_1}{\sqrt{y}} + \mathbb{Z} \left( \frac{x}{\sqrt{y}} e_1 + \sqrt{y} e_2 \right)$$

The shortest vector is then  $e_1/\sqrt{y}$ , with length  $\frac{1}{\sqrt{y}}$ . So for y < 1, the lowest point is below the horizontal axis.

The element z corresponds to the lattice  $L_z$  such that its lowest point on the vertical line x=1 lies below the x-axis is called **semi-stable**, otherwise it is called **unstable**.

Since the semi-stability is preserved under the action of  $SL_2(\mathbb{Z})$ , the semi-stable locus in the upper half plane  $\mathbb{H}$  is as follows




Figure: Semi-stable locus over  $\mathbb H$ 

# In higher dimensional

THANK YOU FOR YOUR ATTENTION.