

Normal Families and the Riemann Mapping theorem

Tri Nguyen

March 11, 2024

This note is used to list every theorems in chapter 9 of the book **Complex made simple**.

1 Quasi-metrics

Definition 1. A function $d: X \times X \rightarrow [0, \infty]$ satisfying the condition

- $d(x, x) = 0 (x \in X)$
- $d(x, y) = d(y, x) (x, y \in X)$
- $d(x, z) \leq d(x, y) + d(y, z) (x, y, z)$

then it is called a quasi-metric on space X . We can see that d is almost the same as a metric except that $d(x, y)$ can be zero for distinct x, y .

One can construct a metric \bar{d} from quasi-metric d , noting that d is an equivalence relation on X . Now we introduction the notion of *concave function*: The function $\psi: I \rightarrow \mathbb{R}$ is said to be *concave* if

$$\psi(tx + (1 - t)y) \leq t\psi(x) + (1 - t)\psi(y),$$

for all $x, y \in X$ and $0 \leq t \leq 1$. It can be inferred from the definition that ψ is concave iff $-\psi$ is convex. Now we have the following lemma

Lemma 1. Suppose that $\psi: I \rightarrow \mathbb{R}$ is concave and $a_1, a_2, b_1, b_2 \in I$ such that $a_1 < b_1, a_2 < b_2, a_2 \geq a_1, b_2 \geq b_1$. Then

$$\frac{\psi(b_2) - \psi(a_2)}{b_2 - a_2} \leq \frac{\psi(b_1) - \psi(a_1)}{b_1 - a_1}$$

Geometrically speaking, the slope of the segment joining two points on the graph of ψ decreases as the points moves to the right.

Applying this lemma for $a_1 = 0, b_1 = x = a_2, b_2 = x + y$, we get the following result:

Lemma 2. Suppose that $\psi: [0, \infty] \rightarrow \mathbb{R}$ is concave and $\psi(0) = 0$. Then

$$\psi(x + y) \leq \psi(x) + \psi(y),$$

for all $x, y \geq 0$.

Lemma 3. Suppose that $\psi: [0, \infty] \rightarrow \mathbb{R}$ is concave and

$$\psi(0) = 0.$$

Suppose further that ψ is nondecreasing, $\psi(t) > 0$ for $t > 0$ and this function is continuous at 0.

If d is a quasi-metric on X then $\tilde{d} = \psi \circ d$ is also a quasi-metric on X such that

- $\tilde{d}(x, y) = 0$ iff $d(x, y) = 0$.
- $\tilde{d}(x_n, y_n) \rightarrow 0$ iff $d(x_n, y_n) \rightarrow 0$.

Now choosing ψ to be a bounded function, we can further assume that \tilde{d} is bounded. Two traditional choices are:

$$\psi_1(t) = \frac{t}{t + 1}$$

and

$$\psi_2(t) = \begin{cases} t & (0 \leq t \leq 1) \\ 1 & (t > 1) \end{cases}$$