

Normal Families and the Riemann Mapping theorem

Tri Nguyen

March 11, 2024

This note is used to list every theorems in chapter 9 of the book **Complex made simple**.

1 Quasi-metrics

Definition 1. A function $d: X \times X \rightarrow [0, \infty]$ satisfying the condition

- $d(x, x) = 0 (x \in X)$
- $d(x, y) = d(y, x) (x, y \in X)$
- $d(x, z) \leq d(x, y) + d(y, z) (x, y, z)$

then it is called a quasi-metric on space X . We can see that d is almost the same as a metric except that $d(x, y)$ can be zero for distinct x, y .

One can construct a metric \bar{d} from quasi-metric d , noting that d is an equivalence relation on X .

Now we introduction the notion of *concave function*: The function $\psi: I \rightarrow \mathbb{R}$ is said to be *concave* if

$$\psi(tx + (1 - t)y) \leq t\psi(x) + (1 - t)\psi(y),$$

for all $x, y \in X$ and $0 \leq t \leq 1$. It can be inferred from the definition that ψ is concave iff $-\psi$ is convex. Now we have the following lemma

Lemma 1. Suppose that $\psi: I \rightarrow \mathbb{R}$ is concave and $a_1, a_2, b_1, b_2 \in I$ such that $a_1 < b_1, a_2 < b_2, a_2 \geq a_1, b_2 \geq b_1$. Then

$$\frac{\psi(b_2) - \psi(a_2)}{b_2 - a_2} \leq \frac{\psi(b_1) - \psi(a_1)}{b_1 - a_1}$$

Geometrically speaking, the slope of the segment joining two points on the graph of ψ decreases as the points moves to the right.