



BIGFOOT SIGHTINGS

AN EXAMINATION

GROUP 11 - FULBRIGHT UNIVERSITY VIETNAM

Pham Nhat Anh | Pham Minh Hoang | Bui Quang Minh |
Nguyen Phan Quynh Nhu | Phan Le Na

INTRODUCTION:


This dataset, sourced from Kaggle, details Bigfoot sightings across the U.S., including geographic locations, sighting classifications, and environmental factors like weather conditions. Sightings are categorized by state, year, and season.

Author: Tim Renner
Uploaded by: The Devastator (@thedevastator)
Source: Kaggle

ABSTRACT:


This infographic analyzes Bigfoot sightings in the U.S., exploring geographic distribution, seasonal patterns, and how weather affects visibility. Washington has the most sightings, and summer sees the highest frequency. Environmental factors like precipitation and humidity reduce visibility, impacting sighting credibility. Sightings spiked in the 1970s and 1980s but have since declined.

WHAT IS THE GEOGRAPHIC DISTRIBUTION OF BIGFOOT SIGHTINGS?





Class A Class B Class C


BIGFOOT SIGHTINGS BY STATE




HOW DOES THE FREQUENCY OF BIGFOOT SIGHTINGS BY CLASSIFICATION VARY ACROSS DIFFERENT SEASONS?




HOW DOES VISIBILITY AFFECT BIGFOOT SIGHTING?



WHAT IS THE FREQUENCY OF BIG FOOT SIGHTINGS OVER THE YEARS



HOW DOES WEATHER (PRECIPITATION, CLOUD COVER, HUMIDITY, AND WINDSPEED) AFFECT VISIBILITY?



CONCLUSION: Bigfoot sightings are concentrated in specific states, with Washington reporting the most. Summer sightings are most common, and poor weather (rain, humidity) possibly reduces visibility. Sightings peaked in the late 20th century but have declined, with Class A sightings being most frequent. Weather and geography influence sighting patterns significantly.