

Advanced Mathematics 2 - Linear Algebra

Chapter 2: Matrix algebra

Department of Mathematics
The FPT university

Chapter 2 Introduction

Chapter 2 Introduction

Topics:

Topics:

2.1 Matrix operations: Addition, Scalar multiplication, Transposition

Topics:

2.1 Matrix operations: Addition, Scalar multiplication, Transposition

2.2 Matrix multiplication

Topics:

- 2.1 Matrix operations: Addition, Scalar multiplication, Transposition
- 2.2 Matrix multiplication
- 2.3 Matrix inverses

Chapter 2 Introduction

Topics:

- 2.1 Matrix operations: Addition, Scalar multiplication, Transposition
- 2.2 Matrix multiplication
- 2.3 Matrix inverses
- 2.5 Matrix transformations

Chapter 2 Introduction

Topics:

- 2.1 Matrix operations: Addition, Scalar multiplication, Transposition
- 2.2 Matrix multiplication
- 2.3 Matrix inverses
- 2.5 Matrix transformations
- 2.7 Applications

2.1 Matrix addition, Scalar multiplication, and Transposition

2.1 Matrix addition, Scalar multiplication, and Transposition

A matrix is an array consisting of m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & a_{ij} & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

2.1 Matrix addition, Scalar multiplication, and Transposition

A matrix is an array consisting of m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & a_{ij} & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- We say A is an $m \times n$ matrix, or A is a matrix of size $m \times n$.

2.1 Matrix addition, Scalar multiplication, and Transposition

A matrix is an array consisting of m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & a_{ij} & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- We say A is an $m \times n$ matrix, or A is a matrix of size $m \times n$. We usually write $A = [a_{ij}]$, $i = 1, \dots, m$, $j = 1, \dots, n$.

2.1 Matrix addition, Scalar multiplication, and Transposition

A matrix is an array consisting of m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & a_{ij} & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- We say A is an $m \times n$ matrix, or A is a matrix of size $m \times n$. We usually write $A = [a_{ij}]$, $i = 1, \dots, m$, $j = 1, \dots, n$.
- The number a_{ij} is called the (i, j) -entry of the matrix.

2.1 Matrix addition, Scalar multiplication, and Transposition

A matrix is an array consisting of m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & a_{ij} & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- We say A is an $m \times n$ matrix, or A is a matrix of size $m \times n$. We usually write $A = [a_{ij}]$, $i = 1, \dots, m$, $j = 1, \dots, n$.
- The number a_{ij} is called the (i, j) -entry of the matrix.
- If $m = n$ then we say A is a **square matrix**.

Matrix Addition and Scalar multiplication

Matrix Addition and Scalar multiplication

Addition and Subtraction

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} + \begin{bmatrix} A & B \\ X & Y \end{bmatrix} = \text{Undefined}$$

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} + \begin{bmatrix} A & B \\ X & Y \end{bmatrix} = \text{Undefined}$$

Scalar multiplication

$$k \begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} = \begin{bmatrix} ka & kb & kc \\ kx & ky & kz \end{bmatrix}, \quad \text{where } k \text{ is a real number}$$

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} + \begin{bmatrix} A & B \\ X & Y \end{bmatrix} = \text{Undefined}$$

Scalar multiplication

$$k \begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} = \begin{bmatrix} ka & kb & kc \\ kx & ky & kz \end{bmatrix}, \quad \text{where } k \text{ is a real number}$$

Example.

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} + \begin{bmatrix} A & B \\ X & Y \end{bmatrix} = \text{Undefined}$$

Scalar multiplication

$$k \begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} = \begin{bmatrix} ka & kb & kc \\ kx & ky & kz \end{bmatrix}, \quad \text{where } k \text{ is a real number}$$

Example. Find X : $2X + \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 3 & 1 \end{bmatrix}$

Matrix Addition and Scalar multiplication

Addition and Subtraction

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \pm \begin{bmatrix} A & B & C \\ X & Y & Z \end{bmatrix} = \begin{bmatrix} a \pm A & b \pm B & c \pm C \\ x \pm X & y \pm Y & z \pm Z \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} + \begin{bmatrix} A & B \\ X & Y \end{bmatrix} = \text{Undefined}$$

Scalar multiplication

$$k \begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} = \begin{bmatrix} ka & kb & kc \\ kx & ky & kz \end{bmatrix}, \quad \text{where } k \text{ is a real number}$$

Example. Find X : $2X + \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 3 & 1 \end{bmatrix}$

Let A, B, C be $m \times n$ matrices. Let 0 denote the $m \times n$ matrix whose entries are 0 . Let k, p be real numbers.

Let A, B, C be $m \times n$ matrices. Let 0 denote the $m \times n$ matrix whose entries are 0. Let k, p be real numbers.

Theorem

- $A + B = B + A$
- $A + 0 = 0 + A = A$
- $A + (B + C) = (A + B) + C$
- $k(A + B) = kA + kB$
- $kA + pA = (k + p)A$
- $k(pA) = (kp)A$
- $1.A = A$

Transpose

Transpose

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example.

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example.
$$\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$$

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example. $\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$

Properties

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example. $\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$

Properties

- $(A^T)^T = A$

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example. $\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$

Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example.
$$\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$$

Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example.
$$\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$$

Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Example. Find A such that:

Transpose

The **transpose** of the matrix A , denoted by A^T , is the matrix whose i th-row is the i th-column of A .

Example. $\begin{bmatrix} 1 & 2 & 3 \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} 1 & x \\ 2 & y \\ 3 & z \end{bmatrix}$

Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$

Example. Find A such that:

$$\left(2A^T + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right)^T = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$$

Symmetric matrices

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Example.

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Example. The matrix $\begin{bmatrix} 1 & 3 \\ 3 & 5 \end{bmatrix}$ is symmetric,

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Example. The matrix $\begin{bmatrix} 1 & 3 \\ 3 & 5 \end{bmatrix}$ is symmetric, but $\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$ is not.

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Example. The matrix $\begin{bmatrix} 1 & 3 \\ 3 & 5 \end{bmatrix}$ is symmetric, but $\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$ is not.

Question.

Symmetric matrices

A square matrix A is **symmetric** if $A = A^T$.

Equivalently, a square matrix A is symmetric if and only if the entries that are symmetric across the main diagonal are equal.

Example. The matrix $\begin{bmatrix} 1 & 3 \\ 3 & 5 \end{bmatrix}$ is symmetric, but $\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$ is not.

Question. Is the sum of two symmetric matrices symmetric?

2.2 Matrix multiplication

2.2 Matrix multiplication

Let A be a $m \times n$ matrix. Let B be an $n \times p$ matrix. Then the product of A and B , denoted by AB , is a matrix of size $m \times p$.

2.2 Matrix multiplication

Let A be a $m \times n$ matrix. Let B be an $n \times p$ matrix. Then the product of A and B , denoted by AB , is a matrix of size $m \times p$.

Examples.

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \begin{bmatrix} M & N \\ P & Q \\ U & V \end{bmatrix} = \begin{bmatrix} aM + bP + cU & aN + bQ + cV \\ xM + yP + zU & xN + yQ + zV \end{bmatrix}$$

2.2 Matrix multiplication

Let A be a $m \times n$ matrix. Let B be an $n \times p$ matrix. Then the product of A and B , denoted by AB , is a matrix of size $m \times p$.

Examples.

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \begin{bmatrix} M & N \\ P & Q \\ U & V \end{bmatrix} = \begin{bmatrix} aM + bP + cU & aN + bQ + cV \\ xM + yP + zU & xN + yQ + zV \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 2a + 3b + 4c \end{bmatrix}$$

2.2 Matrix multiplication

Let A be a $m \times n$ matrix. Let B be an $n \times p$ matrix. Then the product of A and B , denoted by AB , is a matrix of size $m \times p$.

Examples.

$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \begin{bmatrix} M & N \\ P & Q \\ U & V \end{bmatrix} = \begin{bmatrix} aM + bP + cU & aN + bQ + cV \\ xM + yP + zU & xN + yQ + zV \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 2a + 3b + 4c \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} \begin{bmatrix} a & b & c \end{bmatrix} = \begin{bmatrix} 2a & 2b & 2c \\ 3a & 3b & 3c \\ 4a & 4b & 4c \end{bmatrix}$$

The **identity** matrix I_n is the $n \times n$ matrix with 1 on the diagonal and 0 everywhere else.

The **identity** matrix I_n is the $n \times n$ matrix with 1 on the diagonal and 0 everywhere else.

Theorem

Let A, B, C be matrices of sizes such that the indicated operations can be performed. I denotes an identity matrix, and k a real number.

- $AI = IA = A$
- $AB \neq BA$, in general
- $A(BC) = (AB)C$
- $A(B + C) = AB + AC$
- $(B + C)A = BA + CA$
- $k(AB) = (kA)B = A(kB)$
- $(AB)^T = B^T A^T$

Block multiplication

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} =$$

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ \hline 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ \hline 0 & 0 \\ 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} A & 0 \\ I & B \end{bmatrix} \begin{bmatrix} C \\ 0 \end{bmatrix} =$$

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ \hline 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ \hline 0 & 0 \\ 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} A & 0 \\ I & B \end{bmatrix} \begin{bmatrix} C \\ 0 \end{bmatrix} = \begin{bmatrix} AC \\ C \end{bmatrix}$$

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ \hline 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ \hline 0 & 0 \\ 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} A & 0 \\ I & B \end{bmatrix} \begin{bmatrix} C \\ 0 \end{bmatrix} = \begin{bmatrix} AC \\ C \end{bmatrix} = \begin{bmatrix} 31 & 34 \\ 71 & 78 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$$

Block multiplication

To multiply two matrices of large sizes, we can divide each matrix into blocks and treat these as entries.

Example 1.

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ \hline 1 & 0 & 5 & 6 \\ 0 & 1 & 7 & 8 \end{bmatrix} \begin{bmatrix} 9 & 10 \\ 11 & 12 \\ \hline 0 & 0 \\ 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} A & 0 \\ I & B \end{bmatrix} \begin{bmatrix} C \\ 0 \end{bmatrix} = \begin{bmatrix} AC \\ C \end{bmatrix} = \begin{bmatrix} 31 & 34 \\ 71 & 78 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$$

Example 2.

Example 2.

$$\left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 \\ 0 & 1 & 3 & 4 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]^{10}$$

Example 2.

$$\left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 \\ 0 & 1 & 3 & 4 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]^{10} = \begin{bmatrix} I & A \\ 0 & I \end{bmatrix}^{10}$$

Example 2.

$$\left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 \\ 0 & 1 & 3 & 4 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]^{10} = \begin{bmatrix} I & A \\ 0 & I \end{bmatrix}^{10} = \begin{bmatrix} I & 10A \\ 0 & I \end{bmatrix}$$

Example 2.

$$\left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 \\ 0 & 1 & 3 & 4 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]^{10} = \begin{bmatrix} I & A \\ 0 & I \end{bmatrix}^{10} = \begin{bmatrix} I & 10A \\ 0 & I \end{bmatrix} = \begin{bmatrix} 1 & 0 & 10 & 20 \\ 0 & 1 & 30 & 40 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Matrices and Linear systems

Matrices and Linear systems

A system of linear equations

Matrices and Linear systems

A system of linear equations

$$\left\{ \begin{array}{cccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ & & & & \cdots & & & & \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array} \right.$$

Matrices and Linear systems

A system of linear equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be written as a matrix equation $AX = B$, where

Matrices and Linear systems

A system of linear equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \cdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

can be written as a matrix equation $AX = B$, where

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ & & \cdots & \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

2.3 Matrix inverses

2.3 Matrix inverses

2.3 Matrix inverses

Let A be an $n \times n$ matrix.

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

Notes.

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

Notes.

- A matrix may not have an inverse.

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

Notes.

- A matrix may not have an inverse. For example, the matrix $\begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$ does not have an inverse.

2.3 Matrix inverses

Let A be an $n \times n$ matrix. An $n \times n$ matrix B is called an **inverse** of A if $AB = BA = I_n$

Notes.

- A matrix may not have an inverse. For example, the matrix $\begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$ does not have an inverse.
- The inverse matrix of a matrix A , if exists, is unique and is denoted by A^{-1} . In this case we say A is **invertible**.

Inverses of 2×2 matrices

Inverses of 2×2 matrices

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The **determinant** of $A = \det(A) := ad - bc$.

Inverses of 2×2 matrices

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The **determinant** of $A = \det(A) := ad - bc$.

Theorem

The matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is invertible if and only if $\det(A) \neq 0$. In that case,

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Inverses of 2×2 matrices

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The **determinant** of $A = \det(A) := ad - bc$.

Theorem

The matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is invertible if and only if $\det(A) \neq 0$. In that case,

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example.

Inverses of 2×2 matrices

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The **determinant** of $A = \det(A) := ad - bc$.

Theorem

The matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is invertible if and only if $\det(A) \neq 0$. In that case,

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example. Find inverses of the matrices

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix},$$

Inverses of 2×2 matrices

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The **determinant** of $A = \det(A) := ad - bc$.

Theorem

The matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is invertible if and only if $\det(A) \neq 0$. In that case,

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example. Find inverses of the matrices

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 4 \\ 3 & 6 \end{bmatrix}$$

Application in solving linear systems

Consider a system of linear equations $AX = B$. If the matrix A is invertible, then $X = A^{-1}B$ is the solution.

Application in solving linear systems

Consider a system of linear equations $AX = B$. If the matrix A is invertible, then $X = A^{-1}B$ is the solution.

Example. Solve the system

$$\begin{cases} 2x_1 - x_2 = 3 \\ 3x_1 + 4x_2 = -1 \end{cases}$$

Application in solving linear systems

Consider a system of linear equations $AX = B$. If the matrix A is invertible, then $X = A^{-1}B$ is the solution.

Example. Solve the system

$$\begin{cases} 2x_1 - x_2 = 3 \\ 3x_1 + 4x_2 = -1 \end{cases}$$

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.
- To find A^{-1} , perform $[A|I] \xRightarrow{\text{reduced-row echelon}} [I|B]$. Then B is the inverse of A .

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.
- To find A^{-1} , perform $[A|I] \xRightarrow{\text{reduced-row echelon}} [I|B]$. Then B is the inverse of A .

Example.

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.
- To find A^{-1} , perform $[A|I] \xRightarrow{\text{reduced-row echelon}} [I|B]$. Then B is the inverse of A .

Example. Find the inverse of the matrix $A = \begin{bmatrix} 2 & 7 & 1 \\ 1 & 4 & -1 \\ 1 & 3 & 0 \end{bmatrix}$.

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.
- To find A^{-1} , perform $[A|I] \xRightarrow{\text{reduced-row echelon}} [I|B]$. Then B is the inverse of A .

Example. Find the inverse of the matrix $A = \begin{bmatrix} 2 & 7 & 1 \\ 1 & 4 & -1 \\ 1 & 3 & 0 \end{bmatrix}$.

$$\left[\begin{array}{ccc|ccc} 2 & 7 & 1 & 1 & 0 & 0 \\ 1 & 4 & -1 & 0 & 1 & 0 \\ 1 & 3 & 0 & 0 & 0 & 1 \end{array} \right] \xRightarrow{\text{reduced}} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -3/2 & -3/2 & 11/2 \\ 0 & 1 & 0 & 1/2 & 1/2 & -3/2 \\ 0 & 0 & 1 & 1/2 & -1/2 & -1/2 \end{array} \right]$$

Matrix Inversion Algorithm

- Let A be invertible. Then $A \xRightarrow{\text{reduced}} I$, and with the same operations $I \xRightarrow{\text{reduced}} A^{-1}$.
- To find A^{-1} , perform $[A|I] \xRightarrow{\text{reduced-row echelon}} [I|B]$. Then B is the inverse of A .

Example. Find the inverse of the matrix $A = \begin{bmatrix} 2 & 7 & 1 \\ 1 & 4 & -1 \\ 1 & 3 & 0 \end{bmatrix}$.

$$\left[\begin{array}{ccc|ccc} 2 & 7 & 1 & 1 & 0 & 0 \\ 1 & 4 & -1 & 0 & 1 & 0 \\ 1 & 3 & 0 & 0 & 0 & 1 \end{array} \right] \xRightarrow{\text{reduced}} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -3/2 & -3/2 & 11/2 \\ 0 & 1 & 0 & 1/2 & 1/2 & -3/2 \\ 0 & 0 & 1 & 1/2 & -1/2 & -1/2 \end{array} \right]$$

$$\text{Then } A^{-1} = \begin{bmatrix} -3/2 & -3/2 & 11/2 \\ 1/2 & 1/2 & -3/2 \\ 1/2 & -1/2 & -1/2 \end{bmatrix}$$

Theorem

Theorem

- $I^{-1} = I$, where I is an identity matrix.
- $(A^{-1})^{-1} = A$
- If A and B are invertible then $(AB)^{-1} = B^{-1}A^{-1}$.
- If A is invertible then $(A^n)^{-1} = (A^{-1})^n$.
- If A is invertible and $c \neq 0$ then $(cA)^{-1} = \frac{1}{c}A^{-1}$
- If A is invertible then $(A^T)^{-1} = (A^{-1})^T$.

Theorem

- $I^{-1} = I$, where I is an identity matrix.
- $(A^{-1})^{-1} = A$
- If A and B are invertible then $(AB)^{-1} = B^{-1}A^{-1}$.
- If A is invertible then $(A^n)^{-1} = (A^{-1})^n$.
- If A is invertible and $c \neq 0$ then $(cA)^{-1} = \frac{1}{c}A^{-1}$
- If A is invertible then $(A^T)^{-1} = (A^{-1})^T$.

Theorem

- $I^{-1} = I$, where I is an identity matrix.
- $(A^{-1})^{-1} = A$
- If A and B are invertible then $(AB)^{-1} = B^{-1}A^{-1}$.
- If A is invertible then $(A^n)^{-1} = (A^{-1})^n$.
- If A is invertible and $c \neq 0$ then $(cA)^{-1} = \frac{1}{c}A^{-1}$
- If A is invertible then $(A^T)^{-1} = (A^{-1})^T$.

Example.

Theorem

- $I^{-1} = I$, where I is an identity matrix.
- $(A^{-1})^{-1} = A$
- If A and B are invertible then $(AB)^{-1} = B^{-1}A^{-1}$.
- If A is invertible then $(A^n)^{-1} = (A^{-1})^n$.
- If A is invertible and $c \neq 0$ then $(cA)^{-1} = \frac{1}{c}A^{-1}$
- If A is invertible then $(A^T)^{-1} = (A^{-1})^T$.

Example. Find the matrix A that satisfy:

Theorem

- $I^{-1} = I$, where I is an identity matrix.
- $(A^{-1})^{-1} = A$
- If A and B are invertible then $(AB)^{-1} = B^{-1}A^{-1}$.
- If A is invertible then $(A^n)^{-1} = (A^{-1})^n$.
- If A is invertible and $c \neq 0$ then $(cA)^{-1} = \frac{1}{c}A^{-1}$
- If A is invertible then $(A^T)^{-1} = (A^{-1})^T$.

Example. Find the matrix A that satisfy:

$$(A^T - 2I)^{-1} = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$$

Theorem

Let A be an $n \times n$ matrix. The following are equivalent:

- *A is invertible.*
- *The homogeneous system $AX = 0$ has only the trivial solution $X = 0$.*

Theorem

Let A be an $n \times n$ matrix. The following are equivalent:

- *A is invertible.*
- *The homogeneous system $AX = 0$ has only the trivial solution $X = 0$.*
- *$A \xrightarrow{\text{reduced}} I$.*

Theorem

Let A be an $n \times n$ matrix. The following are equivalent:

- *A is invertible.*
- *The homogeneous system $AX = 0$ has only the trivial solution $X = 0$.*
- *$A \xrightarrow{\text{reduced}} I$.*
- *The system $AX = B$ has at least one solution for every choice of column B .*

Theorem


Let A be an $n \times n$ matrix. The following are equivalent:

- *A is invertible.*
- *The homogeneous system $AX = 0$ has only the trivial solution $X = 0$.*
- *$A \xrightarrow{\text{reduced}} I$.*
- *The system $AX = B$ has at least one solution for every choice of column B .*
- *There is an $n \times n$ matrix C such that $AC = I$.*


2.5 Matrix transformation

2.5 Matrix transformation


2.5 Matrix transformation



2.5 Matrix transformation




2.5 Matrix transformation



T is reflection in the x -axis

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

2.5 Matrix transformation




T is reflection in the x -axis

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

2.5 Matrix transformation




T is reflection in the x -axis

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

2.5 Matrix transformation




T is reflection in the x -axis

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

Question.

2.5 Matrix transformation




T is reflection in the x -axis

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

Question. Find the matrix of the following transformation in the plane:

2.5 Matrix transformation



T is reflection in the x -axis


$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

Question. Find the matrix of the following transformation in the plane:

(a) Reflection in the y -axis

2.5 Matrix transformation




T is reflection in the x -axis


$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

Question. Find the matrix of the following transformation in the plane:

- (a) Reflection in the y -axis
- (b) Projection on the x -axis






T is rotation through an angle θ

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


T is induced by the matrix $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$



T is rotation through an angle θ

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

T is induced by the matrix $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$



- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.

- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.
- This matrix A is of size $m \times n$.

- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.
- This matrix A is of size $m \times n$. We say the matrix of T is A , or T is induced by the matrix A .

- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.
- This matrix A is of size $m \times n$. We say the matrix of T is A , or T is induced by the matrix A .

Example.

- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.
- This matrix A is of size $m \times n$. We say the matrix of T is A , or T is induced by the matrix A .

Example. $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$, $T[x, y, z]^T = [2x + y, x - y + z]^T$, is a matrix transformation; it is induced by the matrix $\begin{bmatrix} 2 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$

- A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **matrix transformation** if there is a matrix A such that $T(\vec{v}) = A\vec{v}$ for all $\vec{v} \in \mathbb{R}^n$.
- This matrix A is of size $m \times n$. We say the matrix of T is A , or T is induced by the matrix A .

Example. $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$, $T[x, y, z]^T = [2x + y, x - y + z]^T$, is a matrix transformation; it is induced by the matrix $\begin{bmatrix} 2 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$

Linear transformations

Linear transformations

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example.

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example. Check if the transformation is linear:

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example. Check if the transformation is linear:

(a) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = ab$

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example. Check if the transformation is linear:

(a) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = ab$

(b) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = a + 2b$

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example. Check if the transformation is linear:

(a) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = ab$

(b) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = a + 2b$

(c) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = a + 2b + 1$

Linear transformations

A map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if for all \vec{u}, \vec{v} in \mathbb{R}^n :

- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- $T(k\vec{u}) = kT(\vec{u})$, where k is a real number

Example. Check if the transformation is linear:

(a) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = ab$

(b) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = a + 2b$

(c) $T : \mathbb{R}^2 \rightarrow \mathbb{R}, T \begin{bmatrix} a \\ b \end{bmatrix} = a + 2b + 1$

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1.

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Example 2.

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Example 2. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Example 2. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}, T \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}.$$

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Example 2. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}, T \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}.$$

Find $T \begin{bmatrix} 4 \\ 3 \end{bmatrix}$.

Property of Linear Transformation

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then

$$T(a_1 \vec{x}_1 + a_2 \vec{x}_2 + \cdots + a_k \vec{x}_k) = a_1 T(\vec{x}_1) + a_2 T(\vec{x}_2) + \cdots + a_k T(\vec{x}_k)$$

for all $a_i \in R$ and $\vec{x}_i \in \mathbb{R}^n$.

Example 1. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T(u) = \begin{bmatrix} -2 \\ 3 \end{bmatrix}, T(v) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Find $T(2u - 5v)$.

Example 2. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation satisfying

$$T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}, T \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}.$$

Find $T \begin{bmatrix} 4 \\ 3 \end{bmatrix}$.

Linear Transformation and Matrix Linear Transformation

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then T is a matrix transformation induced by the following $m \times n$ matrix

$$A = [T(E_1) \ T(E_2) \ \dots \ T(E_n)]$$

$$\text{where } E_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, E_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, E_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}.$$

Theorem


Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then T is a matrix transformation induced by the following $m \times n$ matrix

$$A = [T(E_1) \ T(E_2) \ \dots \ T(E_n)]$$


$$\text{where } E_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, E_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, E_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}.$$

Composite of linear transformations

Composite of linear transformations




Composite of linear transformations




The **composite** of S and T , denoted by $S \circ T$, is a transformation from \mathbb{R}^n to \mathbb{R}^p .

Composite of linear transformations



The **composite** of S and T , denoted by $S \circ T$, is a transformation from \mathbb{R}^n to \mathbb{R}^p . It is defined by

Composite of linear transformations




The **composite** of S and T , denoted by $S \circ T$, is a transformation from \mathbb{R}^n to \mathbb{R}^p . It is defined by

$$S \circ T(\vec{x}) = S(T(\vec{x}))$$

for all $\vec{x} \in \mathbb{R}^n$.

Composite of linear transformations



The **composite** of S and T , denoted by $S \circ T$, is a transformation from \mathbb{R}^n to \mathbb{R}^p . It is defined by

$$S \circ T(\vec{x}) = S(T(\vec{x}))$$

for all $\vec{x} \in \mathbb{R}^n$.

Theorem

Theorem

If the matrix of T is A and the matrix of S is B then the matrix of $S \circ T$ is BA .

Theorem

If the matrix of T is A and the matrix of S is B then the matrix of $S \circ T$ is BA .

Example. Let T be reflection in the x -axis followed by rotation through $\pi/2$. Find the matrix of T .

Theorem

If the matrix of T is A and the matrix of S is B then the matrix of $S \circ T$ is BA .

Example. Let T be reflection in the x -axis followed by rotation through $\pi/2$. Find the matrix of T .

The matrix of rotation $R_{\pi/2}$ through $\pi/2$ is $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

Theorem

If the matrix of T is A and the matrix of S is B then the matrix of $S \circ T$ is BA .

Example. Let T be reflection in the x -axis followed by rotation through $\pi/2$. Find the matrix of T .

The matrix of rotation $R_{\pi/2}$ through $\pi/2$ is $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

The matrix of reflection S in x -axis is $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$.

Theorem

If the matrix of T is A and the matrix of S is B then the matrix of $S \circ T$ is BA .

Example. Let T be reflection in the x -axis followed by rotation through $\pi/2$. Find the matrix of T .

The matrix of rotation $R_{\pi/2}$ through $\pi/2$ is $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

The matrix of reflection S in x -axis is $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$.

The matrix of the composite $T = R_{\pi/2} \circ S$ is

$$AB = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

2.7 Application: Input-Output economic models

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.
- There are three industries in the society - farming, housing, garment - that produce these commodities.

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.
- There are three industries in the society - farming, housing, garment - that produce these commodities.
- Each of these industries consumes a certain proportion of the total output of each commodity.

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.
- There are three industries in the society - farming, housing, garment - that produce these commodities.
- Each of these industries consumes a certain proportion of the total output of each commodity.

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.
- There are three industries in the society - farming, housing, garment - that produce these commodities.
- Each of these industries consumes a certain proportion of the total output of each commodity.

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Problem.

2.7 Application: Input-Output economic models

- A society has three basic needs: food, shelter, clothing.
- There are three industries in the society - farming, housing, garment - that produce these commodities.
- Each of these industries consumes a certain proportion of the total output of each commodity.

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Problem. Find the annual prices that each industry must charge for its income to equal its expenditures.

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Let p_1, p_2, p_3 be the annual prices charged by the farming, housing and garment industries.

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Let p_1, p_2, p_3 be the annual prices charged by the farming, housing and garment industries.

$$\begin{cases} 0.4p_1 + 0.2p_2 + 0.3p_3 = p_1 \\ 0.2p_1 + 0.6p_2 + 0.4p_3 = p_2 \\ 0.4p_1 + 0.2p_2 + 0.3p_3 = p_3 \end{cases}$$

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Let p_1, p_2, p_3 be the annual prices charged by the farming, housing and garment industries.

$$\begin{cases} 0.4p_1 + 0.2p_2 + 0.3p_3 = p_1 \\ 0.2p_1 + 0.6p_2 + 0.4p_3 = p_2 \\ 0.4p_1 + 0.2p_2 + 0.3p_3 = p_3 \end{cases}$$

Solution:

		OUTPUT		
		Farming	Housing	Garment
CONSUMPTION	Farming	0.4	0.2	0.3
	Housing	0.2	0.6	0.4
	Garment	0.4	0.2	0.3

Let p_1, p_2, p_3 be the annual prices charged by the farming, housing and garment industries.

$$\begin{cases} 0.4p_1 + 0.2p_2 + 0.3p_3 = p_1 \\ 0.2p_1 + 0.6p_2 + 0.4p_3 = p_2 \\ 0.4p_1 + 0.2p_2 + 0.3p_3 = p_3 \end{cases}$$

Solution: $(p_1, p_2, p_3) = (2t, 3t, 2t)$

Put

Put

$$E = \begin{bmatrix} 0.4 & 0.2 & 0.3 \\ 0.2 & 0.6 & 0.4 \\ 0.4 & 0.2 & 0.3 \end{bmatrix}, \quad P = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Put

$$E = \begin{bmatrix} 0.4 & 0.2 & 0.3 \\ 0.2 & 0.6 & 0.4 \\ 0.4 & 0.2 & 0.3 \end{bmatrix}, \quad P = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Then the equations can be rewritten as

Put

$$E = \begin{bmatrix} 0.4 & 0.2 & 0.3 \\ 0.2 & 0.6 & 0.4 \\ 0.4 & 0.2 & 0.3 \end{bmatrix}, \quad P = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Then the equations can be rewritten as

$$(I - E)P = 0$$

Put

$$E = \begin{bmatrix} 0.4 & 0.2 & 0.3 \\ 0.2 & 0.6 & 0.4 \\ 0.4 & 0.2 & 0.3 \end{bmatrix}, \quad P = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Then the equations can be rewritten as

$$(I - E)P = 0$$

The matrix E is called the **input-output** matrix for the economy.

Put

$$E = \begin{bmatrix} 0.4 & 0.2 & 0.3 \\ 0.2 & 0.6 & 0.4 \\ 0.4 & 0.2 & 0.3 \end{bmatrix}, \quad P = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Then the equations can be rewritten as

$$(I - E)P = 0$$

The matrix E is called the **input-output** matrix for the economy.

Solutions P to the equation $(I - E)P = 0$ are called **equilibrium price structures**.