

# Chapter 1 Matrix Algebra

# CONTENTS

1. Matrices
2. Special matrices
3. Matrix Addition, Scalar Multiplication, and Transposition
4. Matrix Multiplication (2.3 7e)
5. Elementary row operations
6. Rank of the matrix
7. Matrix Inverses (2.4 7e)
8. Matrix Transformations (2.2 & 2.6 7e)

# OBJECTIVES

- ★ Practicing some matrix operations
- ★ Find matrix inverse
- ★ Using matrix transformations to reveal the geometrical meaning of matrix multiplication and inverse.
- ★ The relationship of matrix algebra to linear equations

## DEFINITIONS

**Definition.** An  $m \times n$  matrix (or a matrix of size  $m \times n$ ) is a rectangular array of numbers with  $m$  rows and  $n$  columns

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}_{m \times n} = (a_{ij})_{m \times n}$$

- ★  $A$  is denoted simply as  $A = (a_{ij})$
- ★  $m \times n$  size of the matrix  $A$
- ★ The  $(i, j)$ -entry of  $A$  (denoted by  $a_{ij}$ ) lies in row  $i$  and column  $j$

## Example.

$$A = \begin{bmatrix} 7 & -3 & 1/2 \\ 3 & -5 & 0 \end{bmatrix} \rightarrow \begin{array}{l} (1,3)\text{-entry} \\ a[1,3] = 1/2 \\ a_{13} = 1/2 \end{array}$$

$A$  is a  $2 \times 3$  matrix (or a matrix of size  $2 \times 3$ ) // 2 rows, 3 columns

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 2 & 5 & -1 \end{bmatrix}$$

$3 \times 3$  matrix, a square matrix

$$C = \begin{bmatrix} 2.00 \\ -3.00 \\ 1.75 \end{bmatrix}$$

$3 \times 1$  matrix, column matrix

# ZERO MATRIX


**Definition.** The **zero matrix** (ma trận không) is a matrix in which all elements are zero, denoted by  $0_{m \times n}$  (or  $0$  ).

**Example.**

$$0_{3 \times 4} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# SQUARE MATRIX


**Definition.** An  $n \times n$  matrix is called a square matrix (ma trận vuông) of size  $n$ .

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}$$


$M_n(\mathbb{R})$  : the set of all  $n$  square matrices on  $\mathbb{R}$ .

The elements  $a_{11}; a_{22}; \dots; a_{nn}$  are called the **main diagonal** (đường chéo chính) of  $A$ .

**Example.**  $A = \begin{pmatrix} 2 & -3 & 1 \\ 0 & 4 & 1 \\ 2 & -2 & -3 \end{pmatrix}$



# TRIANGULAR MATRICES

- ★ An **upper triangular** matrix (ma trận tam giác trên) is a square matrix in which all entries **below** the **main diagonal** are zero.

$$M = \begin{pmatrix} -1 & 0 & 3 \\ 0 & 4 & 2 \\ 0 & 0 & -5 \end{pmatrix}, N = \begin{pmatrix} -2 & 3 \\ 0 & 1 \end{pmatrix}$$

- ★ A **lower triangular** matrix (ma trận tam giác dưới) is a square matrix in which all entries **above** the **main diagonal** are zero.

$$P = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 0 & 0 \\ 3 & 1 & 0 \end{pmatrix}, Q = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ -4 & 2 & 0 & 0 \\ 0 & -1 & 0 & 5 \end{pmatrix}$$

- ★ Matrix A is called **triangular** if it is upper or lower triangular



# DIAGONAL MATRIX

A **diagonal matrix** (ma trận đường chéo) is a square matrix with zero entries except possibly on the main diagonal.

$$A = \begin{pmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{pmatrix} = \text{diag}(a_{11}, a_{22}, \dots, a_{nn})$$

**Example.**

$$A = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 5 \end{pmatrix}$$

# IDENTITY MATRIX

The square matrix

$$I_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}$$

is known as the **identity matrix** (ma trận đơn vị) of dimension  $n$ .

**Example.**  $I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \dots$

## EQUAL MATRIX

Two matrices  $A = (a_{ij})$  and  $B = (b_{ij})$ .

$$A = B \Leftrightarrow \begin{cases} A, B \text{ have the same size} \\ a_{ij} = b_{ij}, \forall i, j \end{cases}$$

**Example.** Given  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ ,  $B = \begin{pmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \end{pmatrix}$ ,  $C = \begin{pmatrix} 1 & 0 \\ -1 & 2 \end{pmatrix}$ , discuss the possibility that  $A = B$ ,  $B = C$ ,  $A = C$ .

**Example.** Find  $x, y, z, t$  to

$$\begin{pmatrix} 1 & x-1 & y \\ z & -3 & 5 \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 \\ 2 & -3 & 1-2t \end{pmatrix} \Leftrightarrow \begin{cases} x = 1 \\ y = -1 \\ z = 2 \\ t = -2 \end{cases}$$

# TRANPOSE

- ★ If  $A = (a_{ij})$  is any  $m \times n$  matrix, the transpose of  $A$ , written  $A^T$ , is an  $n \times m$  matrix defined by  $A^T = (a_{ji})$ .
- ★ The **row**  $i$  of  $A$  is the **column**  $i$  of  $A^T$
- ★ The **column**  $j$  of  $A$  is the **row**  $j$  of  $A^T$

**Example.**

$$A = \begin{pmatrix} 1 & -1 & 4 & 5 \\ 6 & -8 & 0 & 1 \\ 0 & 4 & -3 & 6 \end{pmatrix} \Rightarrow A^T = \begin{pmatrix} 1 & 6 & 0 \\ -1 & -8 & 4 \\ 4 & 0 & -3 \\ 5 & 1 & 6 \end{pmatrix}$$

# SYMMETRIC MATRIX - SKEW SYMMETRIC MATRIX

- ★ If  $A^T = A$ , then  $A$  is a **symmetric matrix** (ma trận đối xứng).
- ★ If  $A^T = -A$ , then  $A$  is a **skew symmetric matrix** (ma trận phản xạ).

## Example.

★  $A = \begin{pmatrix} 1 & 2 & -2 \\ 2 & 4 & 5 \\ -2 & 5 & 6 \end{pmatrix}$  is a symmetric matrix.

★  $B = \begin{pmatrix} 0 & -2 & 1 \\ 2 & 0 & -3 \\ -1 & 3 & 0 \end{pmatrix}$  is a skew symmetric matrix.

# SCALAR MULTIPLICATION (PHÉP NHÂN VÔ HƯỚNG)

★ If  $A = [a_{i,j}]$ , that is

$$kA = [ka_{i,j}]$$

★  $kA = 0 \rightarrow$  (either  $k = 0$  or  $A = 0$  )

★  $(k = 0 \text{ or } A = 0) \rightarrow kA = 0$

**Example.** Let  $A = \begin{pmatrix} 3 & 4 & 1 \\ 0 & 1 & -3 \end{pmatrix}$ . Then

$$★ \quad 2A = \begin{pmatrix} 6 & 8 & 2 \\ 0 & 2 & -6 \end{pmatrix}.$$

$$★ \quad -A = \begin{pmatrix} -3 & -4 & -1 \\ 0 & -1 & 3 \end{pmatrix}.$$

# MATRIX ADDITION OF SAME SIZE MATRICES

If  $A = (a_{ij})$  and  $B = (b_{ij})$ , this take the form

$$A \pm B = (a_{ij} \pm b_{ij})_{m \times n}$$

**Example.**

$$\star \begin{pmatrix} -1 & 0 & 2 \\ 2 & 3 & -4 \end{pmatrix} + \begin{pmatrix} 2 & 0 & 2 \\ 5 & -3 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 4 \\ 7 & 0 & -3 \end{pmatrix}$$

$$\star \begin{pmatrix} -1 & 0 & 2 \\ 2 & 3 & -4 \end{pmatrix} - \begin{pmatrix} 2 & 0 & 2 \\ 5 & -3 & 1 \end{pmatrix} = \begin{pmatrix} -3 & 0 & 0 \\ -3 & 6 & -5 \end{pmatrix}$$

### Example.

$$\begin{aligned} 5 \begin{pmatrix} 1 & -2 \\ 4 & 0 \\ -2 & 4 \end{pmatrix} - 2 \begin{pmatrix} 9 & 2 & -1 \\ -8 & 8 & 4 \end{pmatrix}^T &= \begin{pmatrix} 5 & -10 \\ 20 & 0 \\ -10 & 20 \end{pmatrix} - \begin{pmatrix} 18 & -16 \\ 4 & 16 \\ -2 & 8 \end{pmatrix} \\ &= \begin{pmatrix} -13 & 6 \\ 16 & -16 \\ -8 & 12 \end{pmatrix} \end{aligned}$$



## PROPERTIES

Let  $A, B$  and  $C$  denote arbitrary  $n \times m$  matrices where  $n$  and  $m$  are fixed. Let  $k$  and  $p$  denote arbitrary real numbers. Then

1.  $A + B = B + A$  (commutative law: giao hoán)
2.  $A + (B + C) = (A + B) + C$  (associative law: kết hợp)
3. There is an  $n \times m$  matrix  $0$ , such that  $0 + A = A$  for each  $A$ .
4. For each  $A$  there is an  $n \times m$  matrix,  $-A$ , such that  $A + (-A) = 0$
5.  $k(A + B) = kA + kB$ .
6.  $(k + p)A = kA + pA$ .
7.  $(kp)A = k(pA)$ .
8.  $1A = A$ .

# DOT PRODUCT (TÍCH VÔ HƯỚNG)


**Example.**

$$(1 \ 2 \ 3) \cdot \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = 1 \times 3 + 2 \times 2 + 3 \times 1 = 10$$

# MATRIX MULTIPLICATION

Suppose  $A = [a_{ij}]$  is an  $m \times k$  matrix and  $B = [b_{ij}]$  is a  $k \times n$  matrix, then the product  $AB = [c_{ij}]$  is an  $m \times n$  matrix whose the  $(i, j)$ -entry is the dot product of row  $i$  of  $A$  and column  $j$  of  $B$

$$c_{ij} = (\text{row } i \text{ of } A) \cdot (\text{column } j \text{ of } B) = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots a_{ik}b_{kj}$$



**Example.** Give  $A = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix}$ ;  $B = \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix}$  Find  $AB$  và  $BA$ ?

$$AB = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix} = \begin{pmatrix} 7 & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \end{pmatrix}$$

$$c_{11} = \begin{pmatrix} 2 & -1 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} = 2 \times 1 + (-1) \times 3 + 4 \times 2 = 7$$

**Example.** Give  $A = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix}$ ;  $B = \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix}$  Find  $AB$  và  $BA$ ?

$$AB = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix} = \begin{pmatrix} 7 & 12 & c_{13} \\ c_{21} & c_{22} & c_{23} \end{pmatrix}$$

$$c_{12} = \begin{pmatrix} 2 & -1 & 4 \end{pmatrix} \begin{pmatrix} -2 \\ 0 \\ 4 \end{pmatrix} = 2 \times (-2) + (-1) \times 0 + 4 \times 4 = 12$$

**Example.** Give  $A = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix}$ ;  $B = \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix}$  Find  $AB$  và  $BA$ ?

$$AB = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 2 & 4 & 3 \end{pmatrix} = \begin{pmatrix} 7 & 12 & 15 \\ 7 & -8 & 9 \end{pmatrix}$$

$$c_{23} = \begin{pmatrix} 4 & 1 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = 4 \times 2 + 1 \times 1 + 0 \times 3 = 9$$

**Example.** Compute  $AB$  if  $A = \begin{pmatrix} 2 & 3 & 5 \\ 1 & 4 & 7 \\ 0 & 1 & 8 \end{pmatrix}$  and  $B = \begin{pmatrix} 8 & 9 \\ 7 & 2 \\ 6 & 1 \end{pmatrix}$ .

**Solution.**

$$\begin{aligned} C = AB &= \begin{pmatrix} 2 & 3 & 5 \\ 1 & 4 & 7 \\ 0 & 1 & 8 \end{pmatrix} \begin{pmatrix} 8 & 9 \\ 7 & 2 \\ 6 & 1 \end{pmatrix} = \begin{pmatrix} 2 \cdot 8 + 3 \cdot 7 + 5 \cdot 6 & 2 \cdot 9 + 3 \cdot 2 + 5 \cdot 1 \\ 1 \cdot 8 + 4 \cdot 7 + 7 \cdot 6 & 1 \cdot 9 + 4 \cdot 2 + 7 \cdot 1 \\ 0 \cdot 8 + 1 \cdot 7 + 8 \cdot 6 & 0 \cdot 9 + 1 \cdot 2 + 8 \cdot 1 \end{pmatrix} \\ &= \begin{pmatrix} 67 & 29 \\ 78 & 24 \\ 55 & 10 \end{pmatrix} \end{aligned}$$

Does  $BA$  exist?

# THEOREM

Assume that  $a$  is any scalar, and that  $A$ ,  $B$ , and  $C$  are matrices of sizes such that the indicated matrix products are defined. Then:

1.  $IA = A$  and  $AI = A$  where  $I$  denotes an identity matrix.
2.  $A(BC) = (AB)C$ .
3.  $A(B + C) = AB + AC$
4.  $(B + C)A = BA + CA$
5.  $a(AB) = (aA)B = A(aB)$ .
6.  $(AB)^T = B^T A^T$ .



## KTH POWER OF A

For a square matrix  $A$  and positive integer  $k$ , the  $k$  th power of  $A$  is defined by multiplying this matrix by itself repeatedly; that is

$$A^k = \underbrace{AA \cdots A}_{k \text{ times}}$$

where there are  $k$  copies of the matrix  $A$ .

**Example.** For  $A = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$ . Calculate  $A^2$  and  $A^3$

**Solution.**

$$A^2 = AA = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 6 \\ 0 & 1 \end{pmatrix}$$

$$A^3 = A^2A = \begin{pmatrix} 1 & 6 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 9 \\ 0 & 1 \end{pmatrix}$$

# ROW-ECHELON MATRIX

$$\begin{pmatrix} 0 & 1 & * & * & * & * & * \\ 0 & 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

A **row-echelon matrix** has 3 properties

- ★ All the **zero rows** are at the bottom
- ★ The first nonzero entry from the left in each nonzero row is a 1, called the **leading 1** for that row
- ★ Each leading 1 is **to the right** of all leading 1's in the rows above it

# ROW-ECHELON MATRIX

The row-echelon matrix has the “staircase” form

leading ones

$$\begin{pmatrix} 0 & 1 & * & * & * & * & * \\ 0 & 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

( for any choice in \*-position )

# ROW-ECHELON MATRIX

$$\begin{pmatrix} 1 & * & * \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix}$$

~~$$\begin{pmatrix} 1 & * & * & * \\ 0 & 1 & * & * \\ 0 & 1 & * & * \end{pmatrix}$$~~

$$\begin{pmatrix} 0 & 1 & * & * \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

~~$$\begin{pmatrix} 1 & * & * \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$~~

# A REDUCED ROW-ECHELON MATRIX

A **reduced row-echelon matrix** (ma trận bậc thang theo dòng thu gọn) has the properties

- ★ It is a row-echelon matrix
- ★ Each leading 1 is the **only nonzero** entry in its column

$$\begin{pmatrix} 1 & 0 & * & 0 \\ 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The matrix is shown with annotations: red arrows point up to the leading 1s in the first and second columns, and a blue arrow points up to the leading 0 in the third column. The entries in the first and second columns are red, and the entries in the third and fourth columns are blue.

# REDUCED ROW- ECHELON MATRIX

$$\begin{pmatrix} 1 & * & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

~~$$\begin{pmatrix} 1 & * & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$~~

$$\begin{pmatrix} 1 & 0 & * & 0 \\ 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

~~$$\begin{pmatrix} 1 & * & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$~~

**Example.** Find  $x, y$  so that the matrix

$$\begin{pmatrix} 1 & x & y \\ 0 & x & 1 \end{pmatrix}$$

a) is row-echelon matrix

b) is reduced row-echelon matrix

# ELEMENTARY ROW OPERATIONS

The three elementary row operations are:

- ★ (Row Swap) Exchange any two rows:  $d_i \leftrightarrow d_j$
- ★ (Scalar Multiplication) Multiply any row by a constant:  $d_i := \alpha d_i, \alpha \neq 0$
- ★ (Row Sum) Add a multiple of one row to another row:  $d_i := d_i + \beta d_j, \beta \neq 0$

**Example.** Use elementary row operations to transform A to B.

$$A = \begin{pmatrix} 2 & 1 & -1 \\ 1 & -2 & 3 \\ 3 & -1 & 2 \end{pmatrix}; B = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 1 & -7/5 \\ 0 & 0 & 0 \end{pmatrix}.$$



# GAUSSIAN ALGORITHM

**Theorem.** Every matrix can be brought to (reduced) row-echelon form by a series of elementary row operations

- ★ **Step 1.** If all row are **zeros**, stop
- ★ **Step 2.** Otherwise, find the **first column** from the left containing a **nonzero entry** (call it **a**) and **move** the row containing a to the **top position**
- ★ **Step 3.** **Multiply** that row by  **$1/a$**  to creat the **leading 1**
- ★ **Step 4.** By subtracting multiples of that row from the rows below it, make each entry below the leading 1 **zero**
- ★ **Step 5.** Repeat step 1-4 on the matrix consisting of the remaining rows

**Example.** Carry the matrix

$$A = \begin{pmatrix} 2 & 6 & -2 & 2 \\ -2 & -3 & 11 & 4 \\ 3 & 11 & 3 & 0 \end{pmatrix}$$

- ★ to row-echelon matrix
- ★ to reduced row-echelon matrix

## Solution.

$$\begin{pmatrix} 2 & 6 & -2 & 2 \\ -2 & -3 & 11 & 4 \\ 3 & 11 & 3 & 0 \end{pmatrix} \xrightarrow{r_1 \rightarrow \frac{1}{2}r_1} \begin{pmatrix} 1 & 3 & -1 & 1 \\ -2 & -3 & 11 & 4 \\ 3 & 11 & 3 & 0 \end{pmatrix} \xrightarrow[r_3 \rightarrow r_3 - 3r_1]{r_2 \rightarrow r_2 + 2r_1} \begin{pmatrix} 1 & 3 & -1 & 1 \\ 0 & 3 & 9 & 6 \\ 0 & 2 & 6 & -3 \end{pmatrix}$$

$$\xrightarrow{r_2 \rightarrow \frac{1}{3}r_2} \begin{pmatrix} 1 & 3 & -1 & 1 \\ 0 & 1 & 3 & 2 \\ 0 & 2 & 6 & -3 \end{pmatrix} \xrightarrow{r_3 \rightarrow r_3 - 2r_2} \begin{pmatrix} 1 & 3 & -1 & 1 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & -7 \end{pmatrix}$$

$$\xrightarrow{r_3 \rightarrow -\frac{1}{7}r_3} \begin{pmatrix} 1 & 3 & -1 & 1 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

row-echelon matrix

$$\begin{pmatrix} 1 & 3 & -1 & 1 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow[r_2 \rightarrow r_2 - 2r_3]{r_1 \rightarrow r_1 - 2r_3} \begin{pmatrix} 1 & 3 & -1 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{r_1 \rightarrow r_1 - 3r_2} \begin{pmatrix} 1 & 0 & -10 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

reduced row-echelon matrix

# RANK OF THE MATRIX

- ★ The **reduced row-echelon form** of a matrix  $A$  is **uniquely determined** by  $A$ , but the row-echelon form of  $A$  is not unique
- ★ The number  $r$  of leading 1's is the same in each of the different row-echelon matrices
- ★ As  $r$  depends only on  $A$  and not on the row-echelon forms, it is called **the rank of the matrix  $A$** , and written  $\text{rank}(A) = r$

**Example.** If the a matrix A has the row-echelon matrix is

$$\begin{pmatrix} 0 & 1 & * & * & * & * & * \\ 0 & 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

then  $\text{rank}(A) = 4$

**Example.** Compute the rank of

$$A = \begin{pmatrix} 1 & 1 & -1 & 4 \\ 2 & 1 & 3 & 0 \\ 0 & 1 & -5 & 8 \end{pmatrix}$$

**Solution.** The reduction of  $A$  to row - echelon form is

$$A = \begin{pmatrix} 1 & 1 & -1 & 4 \\ 2 & 1 & 3 & 0 \\ 0 & 1 & -5 & 8 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & -1 & 4 \\ 0 & -1 & 5 & -8 \\ 0 & 1 & -5 & 8 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & -1 & 4 \\ 0 & 1 & -5 & 8 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Because this row - echelon matrix has two leading 1 's,  $\text{rank}(A) = 2$ .

**Exercise.** Determine the rank of matrix

$$B = \begin{pmatrix} 1 & 2 & -1 \\ 2 & 4 & -3 \\ 3 & -1 & 0 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 6 & 9 \\ 3 & 6 & 9 & 11 \\ 4 & 8 & 14 & 10 \end{pmatrix}$$
$$D = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 2 & 5 & 1 & 3 \\ 5 & 12 & 2 & 7 \end{pmatrix} \quad E = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{pmatrix}$$

# INVERTIBLE MATRIX

If  $A$  is a square matrix, a matrix  $B$  is called an **inverse** (nghịch đảo) of  $A$  if and only if

$$AB = I \quad \text{and} \quad BA = I$$

A matrix  $A$  that has an inverse is called an **invertible matrix** (ma trận khả nghịch).

The inverse of  $A$  is denoted by  $A^{-1}$ .

$$AA^{-1} = A^{-1}A = I_n$$



**Example.** Show that  $B = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix}$  is an inverse of  $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ .

**Solution.** Compute  $AB$  and  $BA$

$$AB = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad BA = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Hence  $AB = I = BA$ , so  $B$  is indeed an inverse of  $A$ .

## THE INVERSE OF AN $2 \times 2$ MATRIX

Consider the matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

The **determinant** of  $A$  is  $\det(A) = ad - bc$

The **adjugate matrix** (ma trận liên hợp) of  $A$  is defined by  $\text{adj } A = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

$$A^{-1} = \frac{1}{\det(A)} \text{adj } A = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

**Example.**  $A = \begin{pmatrix} 3 & 2 \\ -1 & -1 \end{pmatrix} \implies A^{-1} = \frac{1}{-1} \begin{pmatrix} -1 & -2 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ -1 & -3 \end{pmatrix}$

**Exercise.** Find the inverse of the following matrices

a)  $A = \begin{pmatrix} 4 & 3 \\ 6 & 5 \end{pmatrix}$

b)  $B = \begin{pmatrix} -2 & 1 \\ 7 & 5 \end{pmatrix}$

c)  $C = \begin{pmatrix} 1 & 2 \\ 4 & 8 \end{pmatrix}$

d)  $D = \begin{pmatrix} a & 2 \\ 1 & a \end{pmatrix}$

## THEOREM

All the following matrices are square matrices of the same size.

1.  $I$  is invertible and  $I^{-1} = I$ .
2. If  $A$  is invertible, so  $A^{-1}$ , and  $(A^{-1})^{-1} = A$ .
3. If  $A$  and  $B$  are invertible, so is  $AB$ , and  $(AB)^{-1} = B^{-1}A^{-1}$ .
4. If  $A_1, A_2, \dots, A_k$  are all invertible, so is their product  $A_1A_2 \dots A_k$ , and  $(A_1A_2 \dots A_k)^{-1} = A_k^{-1} \dots A_2^{-1}A_1^{-1}$ .
5. If  $A$  is invertible, so is  $A^k$  for any  $k \geq 1$ , and  $(A^k)^{-1} = (A^{-1})^k$ .
6. If  $A$  is invertible and  $a \neq 0$  is a number, then  $aA$  is invertible and  $(aA)^{-1} = \frac{1}{a}A^{-1}$ .
7. If  $A$  is invertible, so is its transpose  $A^T$ , and  $(A^T)^{-1} = (A^{-1})^T$ .

# FIND AN INVERSE MATRIX BY USING ELEMENTARY ROW OPERATIONS

$$(A \mid I_n) \xrightarrow{\varphi_1} (A_1 \mid B_1) \rightarrow \dots \xrightarrow{\varphi_p} (I_n \mid A^{-1})$$

**Example.** Find the inverse of the following matrix

$$A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ -6 & 2 & 3 \end{pmatrix}$$

$$\begin{aligned}
 (A \mid I_3) &= \left( \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ -6 & 2 & 3 & 0 & 0 & 1 \end{array} \right) \xrightarrow[r_3 \rightarrow r_3 + 6r_1]{r_2 \rightarrow r_2 - r_1} \left( \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & -4 & 3 & 6 & 0 & 1 \end{array} \right) \\
 &\xrightarrow[r_3 \rightarrow r_3 + 4r_2]{r_1 \rightarrow r_1 + r_2} \left( \begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & -1 & 2 & 4 & 1 \end{array} \right) \\
 &\xrightarrow{r_3 \rightarrow -r_3} \left( \begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -4 & -1 \end{array} \right) \\
 &\xrightarrow[r_2 \rightarrow r_2 + r_3]{r_1 \rightarrow r_1 + r_3} \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & -3 & -1 \\ 0 & 1 & 0 & -3 & -3 & -1 \\ 0 & 0 & 1 & -2 & -4 & -1 \end{array} \right)
 \end{aligned}$$

$$A^{-1} = \begin{pmatrix} -2 & -3 & -1 \\ -3 & -3 & -1 \\ -2 & -4 & -1 \end{pmatrix}$$

# MATRIX EQUATIONS

Let  $A, A' \in M_n(\mathbb{R})$  be invertible and  $B \in M_{n \times p}(\mathbb{R}), C \in M_{m \times n}(\mathbb{R}), D \in M_n(\mathbb{R})$ . Then

1.  $AX = B \iff X = A^{-1}B$

2.  $XA = C \iff X = CA^{-1}$

3.  $AXA' = D \iff X = A^{-1}DA'^{-1}$

### Example.

1. Solve the equation  $\begin{pmatrix} 3 & 1 \\ 5 & 2 \end{pmatrix} X = \begin{pmatrix} -2 & 3 \\ 2 & 5 \end{pmatrix}$

2. Solve the equation  $X \begin{pmatrix} 3 & 1 \\ 5 & 2 \end{pmatrix} = \begin{pmatrix} -2 & 3 \\ 2 & 5 \end{pmatrix}$

3. Solve the equation  $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{pmatrix} X \begin{pmatrix} 3 & 2 \\ 3 & 3 \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ 3 & 1 \\ 2 & -1 \end{pmatrix}$



# LINEAR TRANSFORMATION

A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear transformation** (biến đổi tuyến tính) if it satisfies the following two conditions for all vectors  $x$  and  $y$  in  $\mathbb{R}^n$  and all scalars  $a$ :

1.  $T(x + y) = T(x) + T(y)$

2.  $T(ax) = aT(x)$

**Theorem.** If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear transformation, then for each  $k = 1, 2, \dots$

$$T(a_1x_1 + a_2x_2 + \dots + a_kx_k) = a_1T(x_1) + a_2T(x_2) + \dots + a_kT(x_k)$$

for all scalars  $a_i$  and all vectors  $x_i$  in  $\mathbb{R}^n$ .

**Example.** If  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a linear transformation,  $T \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ -3 \end{pmatrix}$  and  $T \begin{pmatrix} 1 \\ -2 \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix}$ , find  $T \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ .

**Solution.** Write  $z = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ ,  $x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ , and  $y = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$ .


★ We want to find numbers  $a$  and  $b$  such that  $z = ax + by$ .

★ The solution is,  $a = \frac{11}{3}$  and  $b = \frac{1}{3}$ , so  $z = \frac{11}{3}x + \frac{1}{3}y$ .

Thus

$$T(z) = T\left(\frac{11}{3}x + \frac{1}{3}y\right) = \frac{11}{3}T(x) + \frac{1}{3}T(y) = \frac{11}{3} \begin{pmatrix} 2 \\ -3 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 5 \\ 1 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 27 \\ -32 \end{pmatrix}$$

# ROTATIONS



Let

$$R_\theta : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

denote the transformation that rotates any vector counterclockwise about the origin through an angle of  $\theta$ .

**Theorem.** The rotation  $R_\theta : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation, and is induced by the matrix


$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

**Example.** We denote by  $R_\pi : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  counterclockwise rotation about the origin through an angle of  $\pi$ . We see that

$$R_\pi \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} -a \\ -b \end{pmatrix}$$

so  $R_\pi$  is a matrix transformation

## REFLECTION IN THE LINE



Let

$$Q_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

denote **reflection in the line  $y = mx$** .

**Theorem.** The transformation  $Q_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ , reflection in the line  $y = mx$ , is a linear transformation and is induced by the matrix


$$\frac{1}{1+m^2} \begin{pmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{pmatrix}$$

**Example.** We denote by  $Q_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  reflection in the line  $y = x$ . We see that

$$Q_m \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} b \\ a \end{pmatrix}$$

so  $Q_m$  is a matrix transformation

## PROJECTION ON THE LINE



Let

$$P_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

denote **projection on the line  $y = mx$** .

**Theorem.** The transformation  $P_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ , projection on the line  $y = mx$ , is a linear transformation and is induced by the matrix

$$\frac{1}{1+m^2} \begin{pmatrix} 1 & m \\ m & m^2 \end{pmatrix}$$

**Example.** Let  $P_m : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  be projection on the line  $x - 2y = 0$ . Find  $P_m \begin{pmatrix} -5 \\ 6 \end{pmatrix}$ .

**Solution.** We have

$$P_m \begin{pmatrix} -5 \\ 6 \end{pmatrix} = \begin{pmatrix} 0.8 & 0.4 \\ 0.4 & 0.2 \end{pmatrix} \begin{pmatrix} -5 \\ 6 \end{pmatrix} = \begin{pmatrix} -1.6 \\ -0.8 \end{pmatrix}$$



**Thank you for your attention.**

**Prob 1.** Let  $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$ ,  $B = \begin{bmatrix} -1 & 1 \\ 3 & 2 \end{bmatrix}$  and  $C = \begin{bmatrix} 1 & 3 & -4 \\ -1 & 2 & 1 \end{bmatrix}$ . Compute the matrix

a)  $2A - B^T$

b)  $AB$

c)  $BA$

d)  $AC$

e)  $CC^T$ .

f)  $C^TC$

g)  $A^3$

h)  $B^2A^T$

**Prob 2.** Suppose that  $A$  and  $B$  are  $n \times n$  matrices. Simplify the expression

a)  $(A + B)^2 - (A - B)^2$

b)  $A(BC - CD) + A(C - B)D - AB(C - D)$

**Prob 3.** Let  $A = \begin{bmatrix} 3 & 1 & 2 \\ 4 & 8 & 0 \\ 0 & 1 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 & 5 & 2 & 1 \\ 1 & 8 & 0 & -6 \\ 1 & 4 & 3 & 7 \end{bmatrix}$ . Compute  $AB$

**Prob 4.** Find the inverse of each of the following matrices.

a)  $\begin{bmatrix} 1 & 5 \\ 2 & -1 \end{bmatrix}$

b)  $\begin{bmatrix} 1 & -1 & 2 \\ -5 & 7 & -11 \\ -2 & 3 & -5 \end{bmatrix}$

**Prob 5.** Given  $A^{-1} = \begin{bmatrix} 1 & -1 & 3 \\ 2 & 0 & 5 \\ -1 & 1 & 0 \end{bmatrix}$ . Find a matrix  $X$  such that

a)  $AX = \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix}$

b)  $AX = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

c)  $XA = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 1 & 1 \end{bmatrix}$

**Prob 6.** Find  $A$  when

a)  $(3A)^{-1} = \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$

b)  $(I + 2A)^{-1} = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}$

c)  $(A^{-1} - 2I)^T = -2 \begin{bmatrix} 1 & 4 \\ 3 & 11 \end{bmatrix}$

**Prob 7.** Find  $A^{-1}$  if

a)  $A^2 - 6A + 5I = 0$

b)  $A^2 + 3A - I = 0$

c)  $A^4 = I$

**Prob 8.** Compute  $\begin{bmatrix} -1 & 3 \\ 0 & 1 \end{bmatrix}^{101}$

**Prob 9.** Solve for  $X$

a)  $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} X = \begin{bmatrix} 1 & -1 \\ 3 & 3 \end{bmatrix}$

b)  $ABXC = B^T$

c)  $AX^TBC = B$  where  $A, B$  and  $C$  are  $n \times n$  invertible matrices.

**Prob 10.** Find the  $(2, 1)$ -entry of the product

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 2 & 5 & 1 \\ 4 & -1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 & 2 & 1 \\ 2 & 3 & 2 \\ 5 & 1 & 0 \\ 0 & 4 & 3 \end{bmatrix}$$

**Prob 11.** Compute the rank of each of the following matrices.

a) 
$$\begin{bmatrix} 1 & 1 & 2 \\ 3 & -1 & 1 \\ -1 & 3 & 4 \end{bmatrix}$$

b) 
$$\begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix}$$

**Prob 12.** Determine the values of  $m$  such that the rank of the matrix is 2.

a) 
$$\begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 5 \\ 1 & 2 & m \end{bmatrix}$$

b) 
$$\begin{bmatrix} 1 & 2 & 1 & 4 \\ 2 & 1 & 1 & 5 \\ -3 & 6 & 1 & m \end{bmatrix}$$

**Prob 13.** Find all values of  $m$  for which the matrix is invertible 
$$\begin{bmatrix} m & 1 & 3 \\ 1 & 3 & 2 \\ 1 & 4 & 5 \end{bmatrix}$$

**Prob 14.** Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation, and assume that  $T(1, 2) = (-1, 1)$  and  $T(0, 3) = (-3, 3)$ .

- a) Compute  $T(11, -5)$       b) Compute  $T(1, 11)$       c) Find the matrix of  $T$

**Prob 15.** Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation such that the matrix of  $T$  is  $\begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$ . Find  $T(3, 2)$