

Compressed random and self-avoiding walks

Nicholas Beaton

Department of Mathematics and Statistics
University of Saskatchewan


Eleventh Prairie Discrete Mathematics Workshop
Banff International Research Station
7–9 August 2015

Collaborators

Tony Guttmann, Iwan Jensen, Greg Lawler

Introduction

A **self-avoiding walk** (SAW) is a walk on a lattice which cannot revisit vertices.



For a given lattice, c_n is the number of n -step SAWs (up to translation).
eg. square lattice:

$$c_0 = 1$$

$$c_1 = 4$$

$$c_2 = 12$$

$$c_3 = 36$$

$$c_4 = 100, \dots$$

For regular lattices in $d \geq 2$, no known expression for c_n . But we still know something!

For regular lattices in $d \geq 2$, no known expression for c_n . But we still know something!

Any SAW of length $m + n$ can be split into two smaller SAWs, of lengths m and n . So

$$c_{m+n} \leq c_m c_n.$$

So $\{c_n\}$ is a sub-multiplicative sequence. Then

$$\log c_{m+n} \leq \log c_m + \log c_n,$$

so $\{\log c_n\}$ is a sub-additive sequence. It follows that the limit

$$\log \mu = \lim_{n \rightarrow \infty} \frac{1}{n} \log c_n$$

exists. $\log \mu$ is called the connective constant of the lattice. Then

$$c_n \sim \theta_n \mu^n,$$

where μ is called the growth constant (sometimes connective constant) and $\theta_n = e^{o(n)}$. By submultiplicativity, we know that $\theta_n \geq 1$.

On the square lattice, the best estimate for μ is 2.63815853031.

On the square lattice, the best estimate for μ is 2.63815853031.

Subexponential factors unproven, but

Conjecture (Nienhuis 1982)

$$c_n \sim A n^{\gamma-1} \mu^n$$

for A, μ, γ constant. A and μ are lattice-dependent, γ depends only on dimension. In two dimensions, $\gamma = 43/32$.

On the square lattice, the best estimate for μ is 2.63815853031.

Subexponential factors unproven, but

Conjecture (Nienhuis 1982)

$$c_n \sim A n^{\gamma-1} \mu^n$$

for A, μ, γ constant. A and μ are lattice-dependent, γ depends only on dimension. In two dimensions, $\gamma = 43/32$.

In high dimensions, can do a bit better:

Theorem (Hara and Slade 1992)

On the hypercubic lattice in five or more dimensions,

$$c_n \sim A \mu^n.$$

Also interested in the **size** and **shape** of SAWs. eg. let $\langle R_e^2 \rangle_n$ be the **mean-squared end-to-end distance** of SAWs of length n .

Conjecture (Nienhuis 1982; Lawler, Schramm and Werner 2004)

$$\langle R_e^2 \rangle_n \sim C n^{2\nu}$$

with C lattice-dependent and ν dimension-dependent. In two dimensions, $\nu = 3/4$.

The exponents γ and ν are also connected to the **scaling limit** of SAWs:

Conjecture (Lawler, Schramm and Werner 2004)

Self-avoiding walks have a conformally invariant scaling limit, namely $SLE_{8/3}$.

Generating functions

The (ordinary) generating function for $\{c_n\}$ is

$$C(z) = \sum_{n \geq 0} c_n z^n$$

Then $z_c = 1/\mu$ is the radius of convergence of $C(z)$. In general, expect the behaviour near z_c to be

$$C(z) \sim \text{const.} (1 - z/z_c)^{-\gamma},$$

which leads to $c_n \sim An^{\gamma-1}\mu^n$.

Generating functions

The (ordinary) generating function for $\{c_n\}$ is

$$C(z) = \sum_{n \geq 0} c_n z^n$$

Then $z_c = 1/\mu$ is the radius of convergence of $C(z)$. In general, expect the behaviour near z_c to be

$$C(z) \sim \text{const.} (1 - z/z_c)^{-\gamma},$$

which leads to $c_n \sim A n^{\gamma-1} \mu^n$.

Because $c_n \geq \mu^n$,

$$C(z) \geq \sum_{n \geq 0} \mu^n z^n = \frac{1}{1 - z\mu}$$

So

$$C(z) \rightarrow \infty \quad \text{as } z \rightarrow z_c^-.$$

Generating functions

The (ordinary) generating function for $\{c_n\}$ is

$$C(z) = \sum_{n \geq 0} c_n z^n$$

Then $z_c = 1/\mu$ is the radius of convergence of $C(z)$. In general, expect the behaviour near z_c to be

$$C(z) \sim \text{const.} (1 - z/z_c)^{-\gamma},$$

which leads to $c_n \sim A n^{\gamma-1} \mu^n$.

Because $c_n \geq \mu^n$,

$$C(z) \geq \sum_{n \geq 0} \mu^n z^n = \frac{1}{1 - z\mu}$$

So

$$C(z) \rightarrow \infty \quad \text{as } z \rightarrow z_c^-.$$

Expect that $C(z)$ is **non-D-finite**, ie. does not satisfy a linear ODE with polynomial coefficients.

Polymer models

SAWs are an important model in statistical mechanics of **linear polymers** in a solvent: chains of monomers, connected by bonds of fixed length and at fixed angles.


Unlike random walks (another, simpler model), SAWs encapsulate the **excluded volume principle**: two different monomers can't occupy the same point in space.

Monomers in a polymer can interact with each other, other polymers, surfaces (both penetrable and impenetrable) or with other external agents. Usually, these interactions are either **attractive** or **repulsive**.

Can also model **forces** applied to the polymer at various points/directions.


Compressive force

The model we are interested in here is a polymer between two impenetrable plates, being compressed together.



Compressive force


The model we are interested in here is a polymer between two impenetrable plates, being compressed together.



Take walks in the upper half-plane, starting at the origin. For a walk γ which reaches maximum height $h(\gamma)$ above the surface, associate a **Boltzmann weight** $e^{-f \cdot h(\gamma)}$.

Compressive force

The model we are interested in here is a polymer between two impenetrable plates, being compressed together.



Take walks in the upper half-plane, starting at the origin. For a walk γ which reaches maximum height $h(\gamma)$ above the surface, associate a **Boltzmann weight** $e^{-f \cdot h(\gamma)}$.

So the walk above receives weight e^{-3f} .

The **partition function** of walks of length n is then

$$Z_n(f) = \sum_{|\gamma|=n} e^{-f \cdot h(\gamma)}$$

and the **free energy** is

$$\lambda(f) = \lim_{n \rightarrow \infty} \frac{1}{n} \log Z_n(f)$$

which can be shown to exist for all f .

The **partition function** of walks of length n is then

$$Z_n(f) = \sum_{|\gamma|=n} e^{-f \cdot h(\gamma)}$$

and the **free energy** is


$$\lambda(f) = \lim_{n \rightarrow \infty} \frac{1}{n} \log Z_n(f)$$

which can be shown to exist for all f .

$f > 0$ represents a force pushing down towards the surface. $f < 0$ represents a force pulling away from the surface. When f is large and positive, walks with small $h(\gamma)$ dominate the partition function. When f is large and negative, walks with large $h(\gamma)$ dominate.


$\lambda(f)$ is a continuous, convex function of f , and is almost-everywhere differentiable.

It has been proven [NRB 2015] that $\lambda(f)$ has a point of non-analyticity at $f = 0$: it is strictly decreasing for $f < 0$, but $\lambda(f) = \log \mu$ for $f \geq 0$:



$\lambda(f)$ is a continuous, convex function of f , and is almost-everywhere differentiable.

It has been proven [NRB 2015] that $\lambda(f)$ has a point of non-analyticity at $f = 0$: it is strictly decreasing for $f < 0$, but $\lambda(f) = \log \mu$ for $f \geq 0$:



This indicates a **phase transition** at $f = 0$.

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

Might expect that as $f \rightarrow \infty$, only the walks which stick to the surface “survive”, which would mean $\lambda(f) \rightarrow 0$. So what's going on?

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

Might expect that as $f \rightarrow \infty$, only the walks which stick to the surface “survive”, which would mean $\lambda(f) \rightarrow 0$. So what’s going on?

$\lambda(f)$ only captures the **exponential behaviour** of $Z_n(f)$. ie.

$$Z_n(f) \sim \theta_n(f) \exp\{\lambda(f) \cdot n\}$$

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

Might expect that as $f \rightarrow \infty$, only the walks which stick to the surface “survive”, which would mean $\lambda(f) \rightarrow 0$. So what’s going on?

$\lambda(f)$ only captures the **exponential behaviour** of $Z_n(f)$. ie.

$$Z_n(f) \sim \theta_n(f) \exp\{\lambda(f) \cdot n\}$$

It is the subexponential term $\theta_n(f)$ where the interesting stuff happens!

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

Might expect that as $f \rightarrow \infty$, only the walks which stick to the surface “survive”, which would mean $\lambda(f) \rightarrow 0$. So what’s going on?

$\lambda(f)$ only captures the **exponential behaviour** of $Z_n(f)$. ie.

$$Z_n(f) \sim \theta_n(f) \exp\{\lambda(f) \cdot n\}$$

It is the subexponential term $\theta_n(f)$ where the interesting stuff happens!

Note:

Conjecture

$$Z_n(0) \sim C n^{-3/64} \mu^n$$

for a constant C .

However, this behaviour is slightly puzzling. No matter how large we make f (favouring walks which stay close to the surface, and punishing those which wander away), $\lambda(f)$ remains constant and equal to its value at $f = 0$.

Might expect that as $f \rightarrow \infty$, only the walks which stick to the surface “survive”, which would mean $\lambda(f) \rightarrow 0$. So what’s going on?

$\lambda(f)$ only captures the **exponential behaviour** of $Z_n(f)$. ie.

$$Z_n(f) \sim \theta_n(f) \exp\{\lambda(f) \cdot n\}$$

It is the subexponential term $\theta_n(f)$ where the interesting stuff happens!

Note:

Conjecture

$$Z_n(0) \sim C n^{-3/64} \mu^n$$

for a constant C .

For $f < 0$, expect that $\theta_n(f)$ does not depend on n .

Random walks

Figuring out what $\theta_n(f)$ looks like is hard – uses lots of SLE, probability, etc...

Random walks

Figuring out what $\theta_n(f)$ looks like is hard – uses lots of SLE, probability, etc...

But we can instead look at **random** (not self-avoiding) walks, and get a similar result!

Random walks

Figuring out what $\theta_n(f)$ looks like is hard – uses lots of SLE, probability, etc...

But we can instead look at **random** (not self-avoiding) walks, and get a similar result!

Now have partition functions $R_n(f)$, summed over all random walks of length n . Then the free energy $\rho(f)$ looks similar to $\lambda(f)$, but with $\log \mu$ replaced by $\log 4$. There is still a phase transition at $f = 0$.

Random walks

Figuring out what $\theta_n(f)$ looks like is hard – uses lots of SLE, probability, etc...

But we can instead look at **random** (not self-avoiding) walks, and get a similar result!

Now have partition functions $R_n(f)$, summed over all random walks of length n . Then the free energy $\rho(f)$ looks similar to $\lambda(f)$, but with $\log \mu$ replaced by $\log 4$. There is still a phase transition at $f = 0$.

Easy to show

Theorem

$$R_n(0) \sim \frac{2}{\sqrt{\pi}} n^{-1/2} 4^n$$

In a horizontal strip of height $h - 2$, let $c_n(h, r, s)$ be the number of n -step walks which start at height r and end at height s , with $0 \leq r, s \leq h - 2$. Then, because random walks are Markovian, $c_n(h, r, s) = K_h^n(r, s)$, where K_h is the $(h - 1) \times (h - 1)$ symmetric matrix with

$$K_h(r, r) = 2, \quad K_h(r, r \pm 1) = 1.$$

In a horizontal strip of height $h - 2$, let $c_n(h, r, s)$ be the number of n -step walks which start at height r and end at height s , with $0 \leq r, s \leq h - 2$. Then, because random walks are Markovian, $c_n(h, r, s) = K_h^n(r, s)$, where K_h is the $(h - 1) \times (h - 1)$ symmetric matrix with

$$K_h(r, r) = 2, \quad K_h(r, r \pm 1) = 1.$$

Let $J_h = \frac{1}{4} K_h$. J_h is tridiagonal and Toeplitz, so it is (with a change of variables) the **Jacobi matrix for Chebyshev polynomials**.

Lemma

$$\det(J_h - \lambda I) = 4^{-h+1} U_{h-1}(1 - 2\lambda),$$

where $U_i(x)$ are Chebyshev polynomials of the second kind:

$$U_0(x) = 1$$

$$U_1(x) = 2x$$

$$U_{i+1}(x) = 2xU_i(x) - U_{i-1}(x)$$

In a horizontal strip of height $h - 2$, let $c_n(h, r, s)$ be the number of n -step walks which start at height r and end at height s , with $0 \leq r, s \leq h - 2$. Then, because random walks are Markovian, $c_n(h, r, s) = K_h^n(r, s)$, where K_h is the $(h - 1) \times (h - 1)$ symmetric matrix with

$$K_h(r, r) = 2, \quad K_h(r, r \pm 1) = 1.$$

Let $J_h = \frac{1}{4} K_h$. J_h is tridiagonal and Toeplitz, so it is (with a change of variables) the **Jacobi matrix for Chebyshev polynomials**.

Lemma

$$\det(J_h - \lambda I) = 4^{-h+1} U_{h-1}(1 - 2\lambda),$$

where $U_i(x)$ are Chebyshev polynomials of the second kind:

$$U_0(x) = 1$$

$$U_1(x) = 2x$$

$$U_{i+1}(x) = 2xU_i(x) - U_{i-1}(x)$$

The roots of $U_i(x)$ are well known:

$$x_k = \cos\left(\frac{k\pi}{i+1}\right) \quad \text{for } k = 1, \dots, i.$$

Eigenvalues and eigenvectors:

$$J_h \mathbf{v}_j = \lambda_j \mathbf{v}_j$$

with

$$\lambda_j = \frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \quad \text{and} \quad \mathbf{v}_j = \left\{ \sin \left(\frac{j(k+1)\pi}{h} \right) \right\}_{k=0, \dots, h-2}$$

Eigenvalues and eigenvectors:

$$J_h \mathbf{v}_j = \lambda_j \mathbf{v}_j$$

with

$$\lambda_j = \frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \quad \text{and} \quad \mathbf{v}_j = \left\{ \sin \left(\frac{j(k+1)\pi}{h} \right) \right\}_{k=0, \dots, h-2}$$

Eigendecomposition:

$$J_h^n(r, s) = \frac{2}{h} \sum_{j=1}^{h-1} \sin \left(\frac{j(r+1)\pi}{h} \right) \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \right]^n \sin \left(\frac{j(s+1)\pi}{h} \right)$$

Eigenvalues and eigenvectors:

$$J_h \mathbf{v}_j = \lambda_j \mathbf{v}_j$$

with

$$\lambda_j = \frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \quad \text{and} \quad \mathbf{v}_j = \left\{ \sin \left(\frac{j(k+1)\pi}{h} \right) \right\}_{k=0, \dots, h-2}$$

Eigendecomposition:

$$J_h^n(r, s) = \frac{2}{h} \sum_{j=1}^{h-1} \sin \left(\frac{j(r+1)\pi}{h} \right) \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \right]^n \sin \left(\frac{j(s+1)\pi}{h} \right)$$

If $n, h \rightarrow \infty$ with $h^2 \ll n$, then $j = 1$ term dominates, so

$$\begin{aligned} J_h^n(r, s) &\sim \frac{2}{h} \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{\pi}{h} \right) \right]^n \sin \left(\frac{(r+1)\pi}{h} \right) \sin \left(\frac{(s+1)\pi}{h} \right) \\ &\sim \frac{2}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \sin \left(\frac{(r+1)\pi}{h} \right) \sin \left(\frac{(s+1)\pi}{h} \right) \end{aligned}$$

Eigenvalues and eigenvectors:

$$J_h \mathbf{v}_j = \lambda_j \mathbf{v}_j$$

with

$$\lambda_j = \frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \quad \text{and} \quad \mathbf{v}_j = \left\{ \sin \left(\frac{j(k+1)\pi}{h} \right) \right\}_{k=0, \dots, h-2}$$

Eigendecomposition:

$$J_h^n(r, s) = \frac{2}{h} \sum_{j=1}^{h-1} \sin \left(\frac{j(r+1)\pi}{h} \right) \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{j\pi}{h} \right) \right]^n \sin \left(\frac{j(s+1)\pi}{h} \right)$$

If $n, h \rightarrow \infty$ with $h^2 \ll n$, then $j = 1$ term dominates, so

$$\begin{aligned} J_h^n(r, s) &\sim \frac{2}{h} \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{\pi}{h} \right) \right]^n \sin \left(\frac{(r+1)\pi}{h} \right) \sin \left(\frac{(s+1)\pi}{h} \right) \\ &\sim \frac{2}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \sin \left(\frac{(r+1)\pi}{h} \right) \sin \left(\frac{(s+1)\pi}{h} \right) \end{aligned}$$

In particular, we want walk to start at height 0, so

$$J_h^n(0, s) \sim \frac{2\pi}{h^2} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \sin \left(\frac{(s+1)\pi}{h} \right)$$

Can end at any height between 0 and $h - 2$, so define

$$F_n(h) = \sum_{s=0}^{h-2} J_h^n(0, s) \sim \frac{4}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \quad \text{for } h^2 \ll n$$

Can end at any height between 0 and $h - 2$, so define

$$F_n(h) = \sum_{s=0}^{h-2} J_h^n(0, s) \sim \frac{4}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \quad \text{for } h^2 \ll n$$

Now $F_n(h)$ is the probability that a SRW in \mathbb{Z}^2 of length n has ordinate between 0 and $h - 2$ for its entire length.

Can end at any height between 0 and $h - 2$, so define

$$F_n(h) = \sum_{s=0}^{h-2} J_h^n(0, s) \sim \frac{4}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \quad \text{for } h^2 \ll n$$

Now $F_n(h)$ is the probability that a SRW in \mathbb{Z}^2 of length n has ordinate between 0 and $h - 2$ for its entire length.

So $F_n(h + 2) - F_n(h + 1)$ is the probability that a SRW has ordinate between 0 and h for its entire length **and** reaches h at some point.

Can end at any height between 0 and $h - 2$, so define

$$F_n(h) = \sum_{s=0}^{h-2} J_h^n(0, s) \sim \frac{4}{h} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \quad \text{for } h^2 \ll n$$

Now $F_n(h)$ is the probability that a SRW in \mathbb{Z}^2 of length n has ordinate between 0 and $h - 2$ for its entire length.

So $F_n(h + 2) - F_n(h + 1)$ is the probability that a SRW has ordinate between 0 and h for its entire length **and** reaches h at some point.

So the partition function $R_n(f)$ is

$$\begin{aligned} R_n(f) &= 4^n \sum_{h=0}^{\infty} e^{-fh} [F_n(h + 2) - F_n(h + 1)] \\ &= 4^n e^{2f} (1 - e^{-f}) \sum_{h=2}^{\infty} e^{-fh} F_n(h) \\ &\sim 4^{n+1} e^{2f} (1 - e^{-f}) \sum_{h=2}^{\infty} e^{-fh} h^{-1} \exp \left\{ -\frac{n\pi^2}{4h^2} \right\} \end{aligned}$$

Convert sum to integral:

$$\sim 4^{n+1} e^{2f} (1 - e^{-f}) \int_0^\infty x^{-1} \exp \left\{ - \left(\frac{n\pi^2}{4x^2} + ux \right) \right\} dx$$

Convert sum to integral:

$$\sim 4^{n+1} e^{2f} (1 - e^{-f}) \int_0^\infty x^{-1} \exp \left\{ - \left(\frac{n\pi^2}{4x^2} + ux \right) \right\} dx$$

The asymptotics of this integral can be computed (with care). Get

Theorem

$$R_n(f) \sim A \cdot 4^n \cdot n^{-1/6} \cdot f^{-1/3} e^f (e^f - 1) \cdot \exp \left\{ B \cdot n^{1/3} \cdot f^{2/3} \right\} \quad \text{for } f > 0$$

for known constants A and B .

Convert sum to integral:

$$\sim 4^{n+1} e^{2f} (1 - e^{-f}) \int_0^\infty x^{-1} \exp \left\{ - \left(\frac{n\pi^2}{4x^2} + ux \right) \right\} dx$$

The asymptotics of this integral can be computed (with care). Get

Theorem

$$R_n(f) \sim A \cdot 4^n \cdot n^{-1/6} \cdot f^{-1/3} e^f (e^f - 1) \cdot \exp \left\{ B \cdot n^{1/3} \cdot f^{2/3} \right\} \quad \text{for } f > 0$$

for known constants A and B .

Exponential term is still 4^n with power-law correction $n^{-1/6}$, but now there is a “stretched exponential” term of the form $\exp\{c \cdot n^{1/3}\}$.

Convert sum to integral:

$$\sim 4^{n+1} e^{2f} (1 - e^{-f}) \int_0^\infty x^{-1} \exp \left\{ - \left(\frac{n\pi^2}{4x^2} + ux \right) \right\} dx$$

The asymptotics of this integral can be computed (with care). Get

Theorem

$$R_n(f) \sim A \cdot 4^n \cdot n^{-1/6} \cdot f^{-1/3} e^f (e^f - 1) \cdot \exp \left\{ B \cdot n^{1/3} \cdot f^{2/3} \right\} \quad \text{for } f > 0$$

for known constants A and B .

Exponential term is still 4^n with power-law correction $n^{-1/6}$, but now there is a “stretched exponential” term of the form $\exp\{c \cdot n^{1/3}\}$.

This matches series analysis.

Can use SLE to estimate the probability that a SAW stays in a strip of height h . Get a similar integral, and use the same method to compute the asymptotics.

Conjecture

$$Z_n(f) \sim A \cdot \mu^n \cdot n^{3/16} \cdot f^{67/115} \cdot \exp \left\{ B \cdot n^{3/7} \cdot f^{4/7} \right\} \quad \text{for } f > 0$$

for constants A, B .

Can use SLE to estimate the probability that a SAW stays in a strip of height h . Get a similar integral, and use the same method to compute the asymptotics.

Conjecture

$$Z_n(f) \sim A \cdot \mu^n \cdot n^{3/16} \cdot f^{67/115} \cdot \exp \left\{ B \cdot n^{3/7} \cdot f^{4/7} \right\} \quad \text{for } f > 0$$

for constants A, B .

Again, matches series analysis.

Unusual asymptotics

Similar results are conjectured for compressed self-avoiding bridges and polygons.

Unusual asymptotics

Similar results are conjectured for compressed self-avoiding bridges and polygons.

These asymptotics are unusual for models like this. Something similar is conjectured to occur in a model of polymer collapse, with a subexponential term like $\exp\{c \cdot n^{1/2}\}$.
[Owczarek, Prellberg & Brak 1993]

Unusual asymptotics

Similar results are conjectured for compressed self-avoiding bridges and polygons.

These asymptotics are unusual for models like this. Something similar is conjectured to occur in a model of polymer collapse, with a subexponential term like $\exp\{c \cdot n^{1/2}\}$.
[Owczarek, Prellberg & Brak 1993]

A term like $\exp\{c \cdot n^{1/2}\}$ has also now been observed in the asymptotics of 1324-avoiding permutations [Conway & Guttmann 2014].

Unusual asymptotics

Similar results are conjectured for compressed self-avoiding bridges and polygons.

These asymptotics are unusual for models like this. Something similar is conjectured to occur in a model of polymer collapse, with a subexponential term like $\exp\{c \cdot n^{1/2}\}$. [Owczarek, Prellberg & Brak 1993]

A term like $\exp\{c \cdot n^{1/2}\}$ has also now been observed in the asymptotics of 1324-avoiding permutations [Conway & Guttmann 2014].

Unclear why such terms appear for some models but not others.

Reference:

NRB, A J Guttmann, I Jensen and G F Lawler, *Compressed self-avoiding walks, bridges and polygons*, submitted, preprint at [arXiv:1506:00296](#)

Reference:

NRB, A J Guttmann, I Jensen and G F Lawler, *Compressed self-avoiding walks, bridges and polygons*, submitted, preprint at [arXiv:1506:00296](https://arxiv.org/abs/1506.00296)

Thank you!