

TRAUMA RESUS

AGE				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
ESTIMATED WEIGHT (kg)				10	12	14	16	18	20	22		25-30			30-40				
TUBES																			
ETT (CUFFED)				3.5	4	4-4.5	4.5	4.5-5	5	5-5.5	5.5	5.5-6	6	6-6.5	6.5			6.5-7.5	
LIP MARKING																			
ETT SIZE X 3																			
LARYNGOSCOPE BLADE					size 1		size 1-2		size 1-2	size 2		size 2		size 2			size 3		
CHEST TUBE SIZE					12-16 F		16-22 F		22-28 F			28-32 F		32-40 F			32-40 F		
RSI DRUGS																			
ETOMIDATE				0.3 MG/KG	MAX 20 MG	suggested doses	3 mg	3 mg	4 mg	5 mg	5 mg	6 mg	6 mg	8 - 10 mg		9 -12 mg		12 - 20 mg	
FENTANYL				1-2 MCG/KG	MAX 100 MCG		10-20 mcg	20 mcg	15-25 mcg	15-30 mcg	20-35 mcg	20-40 mcg	25-40 mcg	25-60 mcg	30-60 mcg	40-80 mcg		40-100mcg	
KETAMINE				1-2 MG/KG	MAX 100 MG		10 mg	15 mg	15 mg	20 mg	20 mg	20 mg	30 mg	30 mg	30-40 mg		40-70 mg		
MIDAZOLAM				0.1-0.2 MG/KG	MAX 10 MG		1 mg	1.5 mg	1.5 mg	2 mg	2 mg	2 mg	2.5 mg	2.5 - 3 mg	3-4 mg		2-5 mg		
ROCURONIUM				1 MG/KG	MAX 100 MG		10 mg	15 mg	15 mg	15 mg	20 mg	20 mg	25 mg	25-30 mg	30-40 mg		40-50 mg		
SUXAMETHONIUM				1.5 MG/KG	MAX 150 MG		15 mg	15 mg	20 mg	25 mg	30 mg	30 mg	30 mg	35-45 mg	45-60 mg		60-100 mg		
FLUID BOLUS																			
NORMAL SALINE				10-20 ML/KG	MAX 500 ML EACH BOLUS														
PCT				10-20 ML/KG	Use leucocyte reduced for < 1 year old														
OTHER DRUGS																			
TRANEXAMIC ACID				15 MG/KG	MAX 1 G	Over 15 mins													
3% SALINE				3-5 ML/KG															
KEPPRA (PROPHYLAXIS)				10 MG/KG	MAX 500 MG														
ADRENALINE 1:10 000				0.1 ML/KG	MAX 10 ML		1 ml	1.2 ml	1.4 ml	1.6 ml	1.8 ml	2 ml	2.2 ml	2.5-3 ml	3-4 ml		5-10 ml		

Note that age determines ETT size and estimated weight; Weight determines drug doses