# Complex Variables Week 1:

Introduction to Complex Numbers

Nichole

August 25, 2018

# 1 What is a Complex Number?

The encounter with complex numbers is usually presented as a solution to the equation:  $x^2=-1 \implies x=\pm \sqrt{-1}$ 

Though this is an intuitive approach for a high school algebra class, it can be misleading. Consider the following:

$$i = \sqrt{-1} = \frac{\sqrt{-1}}{1} = \frac{\sqrt{-1}}{\sqrt{1}} = \sqrt{\frac{-1}{1}} = \sqrt{\frac{1}{-1}} = \frac{1}{i}$$
  
 $\therefore i = \frac{1}{i}$ ?

As you can see from this example, the imaginary number i has properties that are different from real numbers.

#### 1.1 A Better Definition

The complex field  $\mathbb{C}$  is the set of ordered pairs of real numbers (a, b) with the following properties:

Addition: 
$$(a,b) + (c,d) = (a+c,b+d)$$
  
Multiplication:  $(a,b)(c,d) = (ac-bd,ad+bc)$ 

With this definition, we can now consider the ordered pair (0,1) which we will call i. If we multiply i by itself:

$$(0,1)(0,1) = (-1,0) = -1$$

... we can conclude that  $i = \sqrt{-1}$ .

### 1.2 Notation

The most common way to represent complex numbers is: z = (x, y) = x + iy **ALL** complex numbers can (and should) be written in this form.

There are three graphical ways to represent complex numbers. First, they can be represented as points on a 2D plane:




Figure 1: The point z = (1, 1) = 1 + i

Secondly, complex numbers can also be represented as vectors beginning at the origin.




Figure 2: The vector  $\vec{z} = (1,1) = 1 + i$ 

Thirdly, complex numbers can be represented in polar coordinates:  $z=re^{i\theta}=r\mathrm{cis}(\theta)=r(\cos\theta+i\sin\theta)$ 

 $\theta$  is known as the argument of z (sometimes written as  $\arg z$ ) and can be computed from Cartesian coordinates (x,y) by  $\theta=\arctan\frac{y}{x}$ . The radius r can be computed by  $r=|z|=\sqrt{x^2+y^2}$ 




Figure 3: Polar representation

## 1.3 Definitions of Terms

arg(z): Argument of z, also known as  $\theta$ , arg  $z = \arctan \frac{y}{x}$ 

 $\mathbf{cis}(\theta)$ : Short-hand for  $\cos \theta + i \sin \theta$ 

**Complex Conjugate**: Written as  $\bar{z}$  it is the reflection about the Re axis (i.e.  $z = x + iy \implies \bar{z} = x - iy$ )

Im(z): The imaginary part of z (i.e. the y term)

**Magnitude**: Distance from the origin, represented as the radius r (in polar coordinates) and  $|z| = \sqrt{x^2 + y^2}$  (in Cartesian coordinates)

 $\mathbf{Re}(z)$ : The real part of z (i.e. the x term)

- 2 Algebra with Complex Numbers
- 2.1 Addition
- 2.2 Subtraction
- 2.3 Multiplication
- 2.4 Division
- 3 Topological Properties of the Complex Plane
- 3.1 Definitions
- 3.2 Weierstrass M-Test
- 3.3 Stereographic Projection