

Topic 21: AUG experiment analysis meeting

N. Vianello and V. Naulin for the Topic 21 Scientific Team

15 September 2017



This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

## Summary of the campaigns




- ✓ L-Mode experiment, CW 17.
  - 1. Performed similar density ramps in an  $I_{p}$  scan at constant  $q_{95}$
  - 2. Performed similar density ramps in an  $I_{\text{p}}$  scan at constant  $B_{\text{t}}$

#### Summary of the campaigns




- √ L-Mode experiment, CW 17.
  - 1. Performed similar density ramps in an  $I_p$  scan at constant  $q_{95}$
  - 2. Performed similar density ramps in an  $I_p$  scan at constant  $B_t$
- √ H-Mode experiment, CW 21.
  - I. Compare divertor/midplane fueling effect on filamentary transport and profiles without cryo-pumps
  - 2. Compare profiles with the same fueling with/without cryopums
  - Determine an H-Mode with the cryopumps matching similar divertor pressure and SOL profiles






✓ We matched correctly the shape and the L<sub>||</sub> here shown from outer divertor plate up to X-point






- √ The scan was performed with similar puffing rate (0.8-1 MA) whereas we reduced it at lower current to avoid early disruption
- √ We have comparable edge density and divertor neutral pressure






✓ At comparable edge density Upstream profiles are different with the tendency to develop shoulder easier at lower current






- ✓ At comparable edge density Upstream profiles are different with the tendency to develop shoulder easier at lower current
- $\begin{array}{l} \checkmark \quad \text{Still need to provide detail} \\ \text{evolution of edge profiles as a} \\ \text{function of } \Lambda_{\text{div}} \text{ which are} \\ \text{different at different current} \\ \text{although same L}_{\parallel} \end{array}$






✓ We matched correctly the shape and the L<sub>||</sub> here shown from outer divertor plate up to X-point





- √ The scan was performed with similar puffing rate (0.8-1 MA) whereas we reduced it at lower current to avoid early disruption
- √ We have comparable edge density and divertor neutral pressure





✓ At comparable edge density Upstream profiles are different with the tendency to develop shoulder easier at lower current



- ✓ At comparable edge density Upstream profiles are different with the tendency to develop shoulder easier at lower current
- $\begin{array}{l} \checkmark \quad \text{Still need to provide detail} \\ \text{evolution of edge profiles as a} \\ \text{function of $\Lambda_{\text{div}}$ which are} \\ \text{different at different current} \\ \text{although same $L_{||}} \end{array}$